

Beach sand and the potential for infectious disease transmission: observations and recommendations

HELENA M. SOLO-GABRIELE^{1,2}, VALERIE J. HARWOOD³, DAVID KAY⁴, ROGER S. FUJIOKA⁵, MICHAEL J. SADOWSKY⁶, RICHARD L. WHITMAN⁷, ANDREW WITHER⁸, MANUELA CANIÇA^{9,10}, RITA CARVALHO DA FONSECA¹¹, AIDA DUARTE¹², THOMAS A. EDGE¹³, MARIA J. GARGATÉ¹⁴, NINA GUNDE-CIMERMAN¹⁵, FERRY HAGEN¹⁶, SANDRA L. MCLELLAN¹⁷, ALEXANDRA NOGUEIRA DA SILVA¹⁸, MONIKA NOVAK BABIČ¹⁵, SUSANA PRADA^{19,20}, RAQUEL RODRIGUES²¹, DANIELA ROMÃO¹⁴, RAQUEL SABINO¹⁴, ROBERT A. SAMSON²², ESTHER SEGAL²³, CHRISTOPHER STALEY⁶, HUW D. TAYLOR²⁴, CRISTINA VERÍSSIMO¹⁴, CARLA VIEGAS²⁵, HELENA BARROSO²⁶ AND JOÃO C. BRANDÃO¹⁴

¹University of Miami Center for Oceans and Human Health, Key Biscayne, FL 33149, USA, ²Department of Civil, Architectural, and Environmental Engineering, University of Miami, Coral Gables, FL 33146, USA, ³Department of Integrative Biology, University of South Florida, SCA 110, 4202 E. Fowler Ave., Tampa, FL 33620, USA, ⁴Centre for Research into Environment and Health, Institute of Geography and Earth Sciences, Aberystwyth University, Aberystwyth SY24 3DB, UK, ⁵Water Resources Research Center, University of Hawaii, Honolulu, HI 96822, USA, ⁶Department of Soil, Water, & Climate, and BioTechnology Institute, University of Minnesota, St Paul, MN 55108, USA, ⁷Former Chief, Lake Michigan Ecological Research Station, USGS, 1088 N 350 E., Chesterton, IN 46304, USA, ⁸National Oceanography Centre, Liverpool L3 5DA, UK, ⁹National Reference Laboratory for Antibiotic Resistances – Department of Infectious Diseases, National Institute of Health Dr Ricardo Jorge, Av. Padre Cruz 1649-016 Lisboa, Portugal, ¹⁰Centre for the Studies of Animal Science, Institute of Agrarian and Agri-Food Sciences and Technologies, Oporto University, Oporto, Portugal, ¹¹Department of Epidemiology, National Institute of Health Dr Ricardo Jorge, Av. Padre Cruz 1649-016 Lisboa, Portugal, ¹²Department of Microbiology and Immunology, Faculty of Pharmacy, University of Lisbon, Av. Professor Gama Pinto 1649-003 Lisboa, Portugal, ¹³Canada Centre for Inland Waters, Environment Canada, 867 Lakeshore Road, Burlington, ON L7R 4A6, Canada, ¹⁴Reference Unit for Parasitic and Fungal Infections – Department of Infectious Diseases, National Institute of Health Dr Ricardo Jorge, Av. Padre Cruz 1649-016 Lisboa, Portugal, ¹⁵Department of Biology, Biotechnical Faculty, University of Ljubljana, Jamnikarjeva 101, 1000 Ljubljana, Slovenia, ¹⁶Department of Medical Microbiology & Infectious Diseases, Canisius-Wilhelmina Hospital, Nijmegen, the Netherlands, ¹⁷School of Freshwater Sciences, University of Wisconsin–Milwaukee, Milwaukee, WI, USA, ¹⁸Lab. Microbiology, ADEIM – Faculty of Pharmacy, University of Lisbon, Av. Professor Gama Pinto 1649-003 Lisboa, Portugal, ¹⁹Center of Exact Sciences and Engineering, University of Madeira, Campus Universitário da Penteada 9000-390 Funchal, Madeira, Portugal, ²⁰Center of Volcanology and Geological Risk Assessment, University of Azores, 9501-801 Ponta Delgada, Portugal, ²¹Laboratory of Microbiology – Department of Environmental Health, National Institute of Health Dr Ricardo Jorge, Av. Padre Cruz 1649-016 Lisboa, Portugal, ²²CBS-KNAW Fungal Biodiversity Centre, Uppsalalaan 8, 3584 CT Utrecht, The Netherlands, ²³Department of Clinical Microbiology and Immunology, Sackler School of Medicine, Tel-Aviv, University, Tel-Aviv 69978, Israel, ²⁴Environment & Public Health Research Unit, School of the Environment & Technology, University of Brighton, Cockcroft Building, Lewes Road, Brighton BN2 4GJ, UK, ²⁵Environmental Health RG – Lisbon School of Health Technology – Polytechnic Institute of Lisbon, Av. D. João II, Lote 4.69.01, 1990-096 Lisboa, Portugal, ²⁶Interdisciplinary Research Centre Egas Moniz (CiiEm), Institute of Health Sciences Egas Moniz (ISCSEM) Campus, Monte de Caparica, 2829-511, Portugal

Recent studies suggest that sand can serve as a vehicle for exposure of humans to pathogens at beach sites, resulting in increased health risks. Sampling for microorganisms in sand should therefore be considered for inclusion in regulatory programmes aimed at protecting recreational beach users from infectious disease. Here, we review the literature on pathogen levels in beach sand, and their potential for affecting human health. In an effort to provide specific recommendations for sand sampling programmes, we outline published guidelines for beach monitoring programmes, which are currently focused exclusively on measuring microbial levels in water. We also provide background on spatial distribution and temporal characteristics of microbes in sand, as these factors influence sampling programmes. First steps toward establishing a sand sampling programme include identifying appropriate beach sites and use of initial sanitary assessments to refine site selection. A tiered approach is recommended for monitoring. This approach would include the analysis of samples from many sites for faecal indicator organisms and other conventional analytes, while testing for specific pathogens and unconventional indicators is reserved for high-risk sites. Given the diversity of microbes found in sand, studies are urgently needed to identify the most significant aetiological agent of disease and to relate microbial measurements in sand to human health risk.

Keywords: Beach sand, microbes, bacteria, fungi

Submitted 7 April 2015; accepted 19 May 2015

INTRODUCTION

Transmission of infectious diseases in terrestrial beach environments can occur via direct exposure to microbes found in sand or through the flux of microbes from water to sand within the swash or intertidal zone. Exposure to pathogens can include routes such as dermal contact, contact with eyes and ears, inhalation and ingestion. Recent studies suggest that direct exposure to beach sands is a risk factor for infectious disease, particularly in children. An epidemiological study found that gastrointestinal (GI) illness in beach users was associated with exposure to water and intertidal sand (Bonilla *et al.*, 2007; Pinto *et al.*, 2012a, b; Sabino *et al.*, 2014a). A separate epidemiological study (Heaney *et al.*, 2009) found that digging in the sand was positively associated with GI illness and was associated with levels of faecal indicator organisms (FIO), enterococci (Heaney *et al.*, 2012). While correlations between beach sand exposure and infectious disease exist, the specific causative mechanisms of infection are yet to be shown including identification of the aetiological agent.

In addition to direct exposure, sand can also serve as a vehicle for transferring pathogenic microbes to and from the adjacent water. Studies on southern Lake Michigan found that densities of *Escherichia coli* were highest in core samples taken from foreshore sands, often by several logs, but rapidly decreased from that maximum both landwards and lakewards (Whitman & Nevers, 2003; Kinzelman *et al.*, 2004; Whitman *et al.*, 2006a). Studies at marine beaches in Florida found that the intertidal zone (Figure 1), in particular the upper fringe of this zone, was a hot spot for the accumulation of microbes (Shibata *et al.*, 2004; Wright *et al.*, 2011). These microbes can be mobilized from this reservoir by mechanisms such as wave action (Phillips *et al.*, 2014), pore water transport (Phillips *et al.*, 2011b), and then impact adjacent water quality (Phillips *et al.*, 2011a). Microbes may also be re-deposited in sand by incoming current and waves (Ge *et al.*, 2010). Maximal FIO levels in both marine and freshwater beaches generally occur just behind the highest wave up-rush along the beach. Of particular significance is that the foreshore or intertidal zones are areas where beach goers congregate, and where children tend to play with sand. Thus, the swash zone and foreshore is a dynamic area of the shoreline in terms of microbe accumulation and potential exposures. It is also a zone where the interactions between the water and sand are strongest allowing for an interchange of microbes to and from this zone.

To evaluate the current state of knowledge and most pressing research needs in the area of beach sand microbiology, a panel session was convened as part of the TEMPH2014 (Trends in Environmental Microbiology and Public Health, 2014) conference held in Lisbon, Portugal during September 2014. The purpose of the panel was to discuss the potential inclusion of sand quality assessments in monitoring programmes for recreational beaches. The interdisciplinary group of participants held particular expertise in one of two

primary categories, recreational beach water quality and environmental mycology. The ideas presented in this article provide an interesting meld of concepts that would benefit beach sand monitoring programmes. Specifically, this work begins by reviewing documents that provide recommendations for changes to existing recreational water quality monitoring guidelines ('Call-to-Action' documents), and expands upon these documents by emphasizing the merits of including measures of sand. The manuscript then focuses on describing the spatial distribution and temporal characteristics of microbes in sand, which is necessary for developing general recommendations for sampling programmes. Recommendations for sampling programmes begin by identifying appropriate beach sites and inclusion of beach sanitary assessments. Strategies for sampling and analysis follow, including an emphasis on recommending which microbes to measure and on advances in microbe measurement techniques. We conclude with an identification of research needs and a call for the inclusion of microbial monitoring in sand as an integral part of routine beach health assessments.

OVERVIEW OF 'CALL-TO-ACTION' DOCUMENTS

Microbial contamination in recreational waters is monitored through measures of FIOs, including 'generic' (non-pathogenic) *E. coli* and enterococci (EU, 1976, 2006; USEPA, 1986). The enterococci are a group of bacterial species belonging to the genus *Enterococcus*. FIOs are seldom in themselves pathogenic, but since they are found in faeces of both human and animals they are useful indicators for faecal contamination of water. The ubiquitous distribution of FIOs in faeces stands in contrast to the relatively infrequent occurrence of pathogens, whose detection is complicated by their great diversity. Reliance on FIOs for water quality assessment is thus a matter of practicality, as a set of general targets that are highly concentrated and widely distributed in faeces provides an economical metric, while testing large volumes of water for innumerable pathogens is impractical for monitoring.

The sole reliance on FIO levels in water as a mechanism for classifying recreational waters was challenged by the Annapolis Protocol, a document prepared for the World Health Organization and authored by many of the acknowledged world experts in the field (WHO, 1999), well over a decade ago. The Annapolis Protocol first identified the value of a comprehensive sanitary inspection of recreational waters to identify all sources of potential pathogens, a concept further developed over the past decade (Boehm *et al.*, 2009a; Gooch-Moore *et al.*, 2011; Abdelzaher *et al.*, 2013). The Annapolis Protocol introduced the concept of a risk-based approach, and acknowledged that common FIO standards across all waters did not account for the unequal probability of pathogen presence in faecal contamination from different sources. In particular, it identified the probable




Fig. 1. Zones at the interface between beach sand and water. The terminology differs between tidal marine systems and non-tidal freshwater systems. Mean surface water elevations for marine systems tends to vary with tides. For freshwater systems, in particular within lakes, the mean surface water elevation tends to vary with seiche. For river systems, mean surface water elevations vary with the seasonal elevation of the groundwater table and waves tend to run parallel to the beach as opposed to the perpendicular direction observed in most marine and lake settings. (Image modified from Whitman *et al.*, 2014).

reduced health risk when FIOs were primarily from non-human sources. Many of the recommendations in the Annapolis Protocol were incorporated into the World Health Organization (WHO) guidelines for recreational waters (WHO, 2003, 2009). Implicit in the WHO approach is the notion that health risks are unacceptable when they exceed a set FIO threshold level.

As understanding of the differential risk inherent in faeces from different sources was accumulating, microbial source tracking (MST) emerged as a discipline. The goal of MST is to determine the host animals responsible for faecal contamination of water. It is accomplished by analysis of host-associated microorganisms (or host genes in the case of mitochondrial DNA) in the faeces of humans and various animals (reviewed in Stoeckel & Harwood, 2007; Harwood *et al.*, 2014). In 2005 the US Environmental Protection Agency (EPA) produced a document that outlined the uses of MST to aid in total maximum daily load (TMDL) and risk assessment for recreational waters (USEPA, 2005). Since that time, the ability to discriminate among different sources of contamination in water has steadily improved, and MST has been used to explore FIO sources in beach sand (Russell *et al.*, 2013).

The 2003 WHO report has an extensive review of the risk from microbes in sand, but concluded that there was insufficient evidence to support the establishment of a guideline value for indicator or pathogenic microorganisms in beach sand. Neither the subsequent European Union (2006) Bathing Water directive, nor the USEPA (2012) 'NEEAR' criteria consider sand contamination, other than acknowledging that it may be a source of FIOs in the adjacent water (EU, 2006; USEPA, 2012). Health Canada recreational water quality guidelines indicate that testing of sand may be warranted in circumstances such as support for sanitary surveys or disease outbreak investigations, though stated that more

research was needed before guideline values for sand could be established (Health Canada, 2012). Thus, no regulatory criteria exist currently for microbial levels in sand.

MICROBIAL CHARACTERISTICS OF BEACH SAND

Microbes found in beach sand

Numerous studies have been conducted that document the existence of pathogenic microbes in beach sands, providing evidence for sand as a potential reservoir for aetiological agents of disease (Whitman *et al.*, 2014; Sabino *et al.*, 2014a). The pathogenic organisms found in sand come from many groups, including bacteria, viruses, protozoa, helminths (worms) and fungi. For example, pathogenic bacteria detected in beach sands include *Vibrio vulnificus* (Abdelzaher *et al.*, 2010; Shah *et al.*, 2011), *Salmonella* (Yamahara *et al.*, 2012), *Campylobacter* (Yamahara *et al.*, 2012), *Pseudomonas aeruginosa* (Esiobu *et al.*, 2004) and *Staphylococcus aureus* including methicillin resistant strains (Plano *et al.*, 2013). Viruses found in sand have included enterovirus by culture (Shah *et al.*, 2011). Protozoans have included *Giardia* spp. and *Cryptosporidium* spp. in nearshore sands (Abdelzaher *et al.*, 2010). Nematode larvae and eggs have also been readily detected in beach sands (Shah *et al.*, 2011). Similarly, many species of potentially pathogenic yeasts and fungi have been found including *Aspergillus* sp., *Chrysosporium* sp., *Fusarium* sp., *Scedosporium* sp., *Scytalidium* sp., *Scopulariopsis* sp. (Sabino *et al.*, 2011), *Candida* sp. (Shah *et al.*, 2011), *Penicillium* sp., *Rhodotorula mucilaginosa* (Vogel *et al.*, 2007), *Cladosporium* sp., *Mucor* sp. and *Stachybotrys* sp. (Gonzales *et al.*, 2000; Migahed, 2003; Gomes *et al.*, 2008; Bik *et al.*, 2012). Fungi with propensity

to infect skin and nails include *Trichophyton* sp. and *Microsporum* sp. (Sabino *et al.*, 2011). The presence of black yeasts of the genera *Aureobasidium* and *Exophiala*, causing allergies, subcutaneous phaeohyphomycoses and neurotropic infections (de Hoog *et al.*, 2009), have been reported once (Efstratiou & Velegriaki, 2009), probably reflecting their slow growth and consequential difficulty of detection.

Spatial distribution and temporal characteristics

The spatial and temporal distribution of FIOs and fungi is highly variable in sand over distances as small as a few centimetres (WHO, 2003; Bonilla *et al.*, 2007; Whitman *et al.*, 2006b). The heterogeneity at this scale may be due to limited transport and mixing of microbes in sand such that once the sand is inoculated under conditions suitable for growth, microbial distribution becomes very patchy. Extreme patchiness can be due to discrete inputs from dog droppings, seabirds and onshore drift, followed by growth. Different zones of the beach may be more or less conducive to microbe growth. Research suggests that the swash zone, and in particular the area just above the maximum up-rush, may also be conducive for regrowth of FIOs through distinct wetting and drying actions and unique characteristics afforded by wrack, which is defined to include seaweed, algae, Sargassum, kelp, *Cladophora*, macrophytes and other vegetation-like accumulations (Alm *et al.*, 2003; Whitman *et al.*, 2003; Olapade *et al.*, 2006; Ishii *et al.*, 2007; Yamahara *et al.*, 2009). Although sand moisture content of approximately 8% is sufficient to permit the survival of bacteria, yeasts and nematodes (Whitman *et al.*, 2014), there is a lack of consistency in the literature over the distribution of organisms in sand and their relation to moisture content. Generally, a greater density of FIOs in wet foreshore sand has been observed compared with either submerged, backshore, sand at depth, or dry sand (Whitman *et al.*, 2014). Conversely, studies at a Florida marine beach found higher concentrations of *E. coli* and enterococci in supratidal sand (above the high water mark) than in intertidal sand (Abdelzaher *et al.*, 2010).

The accumulation of wrack in the swash zone also serves to maintain FIO populations by serving as a source of nutrients (Byappanahalli *et al.*, 2003; Imamura *et al.*, 2011), providing protection from UV light (Feng *et al.*, 2013), and regulating the temperature and moisture conditions in sands located immediately below them. Studies have shown that *E. coli* and enterococci can survive for over 6 months in sun-dried algal mats (*Cladophora*) stored at 4°C, and the residual bacteria in the dried alga readily grew upon rehydration (Whitman *et al.*, 2003). Experimental work in the UK found that FIOs are liable to persist, and possibly proliferate, in supra-littoral wrack piles on a beach (Ward, 2009; Dunhill *et al.*, 2013). In addition to the unique hydrodynamics of this area, which is conducive to the accumulation of wrack, the swash zone also attracts shorebirds that feed and roost in this area and may contribute to the microbial load through their faeces (Lévesque *et al.*, 1993; Fogarty *et al.*, 2003; Wright *et al.*, 2009; Edge & Hill, 2007; Lu *et al.*, 2011). Bird faeces may contribute directly to beach water contamination, although microbial source tracking techniques have found that beach sand (with bird-derived *E. coli*) can be a more significant secondary source of contamination to

adjacent beach water than directly from the bird droppings themselves (Edge & Hill, 2007).

The growth of microbes in sand is not limited to bacteria. It is well recognized that fungi survive, and even grow in sand (Anderson, 1979). This has been demonstrated using both culture and microscopic analyses. For example, Khiyama & Makemson (1973) reported that culturable fungi in 42 Mediterranean beaches can reach as high as $\sim 7 \times 10^6$ CFU g⁻¹ (Larrondo & Calvo, 1989). Fungi levels at beaches have been observed to vary temporally with extreme events. In the volcanic islands of Madeira and Porto Santo, an archipelago of Portugal, pathogens in the beach sands have been associated with intense rainfall events, flash floods and debris flow (Pereira *et al.*, 2013; Marzol *et al.*, 2006a, b). In a study of 15 Portuguese Atlantic Coast beaches, the highest number of viable fungal colony forming units in sand was in supratidal sand, at around 500 CFU g⁻¹ (Brandão *et al.*, 2002).

Antimicrobial resistance

Environmental reservoirs of both antibiotic resistant bacteria (Francino, 2012; Wellington *et al.*, 2013) and antifungal resistant fungi have been emerging. The causes may be associated with the release of antibiotic and antifungal residues, from agriculture, animal feeding, aquaculture and also hospital wastewater (Jiang *et al.*, 2011; Suzuki & Hoa, 2012; Diwan *et al.*, 2013). A variety of antibiotic resistant bacteria have been isolated from sand and beach water, which can be in contact with humans (Velonakis *et al.*, 2014). Examples include MRSA (methicillin-resistant *Staphylococcus aureus*) which has been detected in correlation with the quality of water and sand, showing a relationship with beach-user overcrowding, the concentration of other microorganisms, the presence of yeasts from human origin, as well as water temperature (Papadakis *et al.*, 1997; Plano *et al.*, 2011; Roberts *et al.*, 2013).

Mudryk *et al.* (2013) showed that *Vibrio* species inhabiting sand were more resistant to antibiotics than those isolated from seawater; in addition, more than 90% of planktonic and benthic *Vibrio*-like bacteria could present multiple antibiotic resistance. Also multidrug resistant *Enterococcus faecium* from beach sand were identified with similar features to those from clinical human isolates (Heikens *et al.*, 2008) indicating that enterococci can be included in the monitoring of sand, with the respective characterization of antibiotic resistance and virulence factors (Pinto *et al.*, 2012a, b).

Fungi can have an intrinsic antifungal resistance to certain antifungal substances (primary resistance) but initially susceptible microorganisms can also develop resistance (secondary resistance). In the first case, we have examples such as *Candida krusei*, resistant to fluconazole (Orozco *et al.*, 1998) or specific *Fusarium* species (Carneiro *et al.*, 2011), resistant to the majority of antifungals used in clinical practice (Alastruey-Izquierdo *et al.*, 2008). These species have coded in their genome molecular mechanisms that enable them to survive in presence of those antifungals. *Candida* spp. and *Fusarium* spp. are frequently found in sand samples and are considered as parameters to evaluate the microbiological quality of a given sample (Sabino *et al.*, 2011). Nevertheless, the number of fungi showing antifungal resistance has been rising over the years. *Aspergillus* is one of the major fungal threats showing high rates of resistance to azoles, especially in Europe (Sabino *et al.*, 2011) with the environment serving as one of the possible sources of resistant strains

(Snelders *et al.*, 2008; Verweij *et al.*, 2009; Mortensen *et al.*, 2010; Chowdhary *et al.*, 2013). In a recent study (Sabino *et al.*, 2014b), the cryptic species of *A. fumigatus*, *A. lentulus* and *Neosartorya pseudofischeri* were detected in samples from beaches. These species have been reported to be resistant *in vitro* to the azole antifungals (Balajee *et al.*, 2005; Yaguchi *et al.*, 2007; Alcazar-Fuoli *et al.*, 2008).

The rapid increase of antibiotic and antifungal resistance compromises the treatment of several bacterial (ECDC, 2013; WHO, 2014) and fungal infections (CDC, 2015). Antibiotic and antifungal resistance is considered a serious threat to human health because of the limited treatments available resulting in greater costs (Finley *et al.*, 2013) and increased morbidity and mortality (Ashbolt *et al.*, 2013; World Health Organization, 2014). MRSA is particularly notorious among the antibiotic resistant bacteria as it causes life-threatening skin ailments that are difficult to treat. *Escherichia coli*, although a natural inhabitant of the human intestine, has several pathogenic forms causing extreme gastrointestinal infection, some of which exhibit cephalosporin resistance (de Kraker *et al.*, 2011). Antibiotic resistant bacteria are found in bathing waters and studies have shown that risks are related to the type of water activity (Leonard *et al.*, 2015). Regarding fungi, invasive *Candida* infections are the fourth leading cause of hospital-acquired bloodstream infections, and they are associated with a high mortality (>40%) (Sipsas *et al.*, 2009). *Candida* and *Aspergillus* species cause a majority of serious infections in non-HIV patients. Because of the high risk of fungal infections in immunocompromised individuals, antifungal prophylaxis is often used to treat these patients. However, the expanding use of antifungal drugs has been associated with increasing incidence of antifungal drug resistance resulting from inherently less sensitive species and/or acquisition of drug class-specific resistance mechanisms (Pfaller *et al.*, 2011). Most alarming in recent years, resistant strains of certain *Candida* or *Aspergillus* species have emerged that are resistant to azoles, especially due to antifungal prophylaxis. Nevertheless, in *Aspergillus*, and considering the molecular mechanisms underlying their high rates of resistance to azoles, especially in Europe (Chowdhary *et al.*, 2013) (TR34/L98H and TR46/Y121F/T289A mutations of the CYP51A gene), it was hypothesized that one of the possible sources of resistant strains is the environment (Snelders *et al.*, 2008; Verweij *et al.*, 2009; Mortensen *et al.*, 2010; Chowdhary *et al.*, 2013).

RECOMMENDATIONS FOR SAND MONITORING PROGRAMMES

Criteria for selecting designated recreational beaches

The European Union under the 1976 directive has a specific two-part definition for bathing waters. It defines bathing waters as areas where bathing is explicitly authorized by the competent authorities of each member State, or where bathing is not prohibited and is traditionally practiced by a large number of bathers; The second criteria means that the public largely self select what will become a bathing area and it is then up to the authorities to ensure suitable microbial quality.

For purposes of this review, we consider a recreational beach to serve as a bathing water. We also consider a recreational beach to be a designated shore and water complex largely used for recreation. Waters may be marine or fresh water (fluvial or lentic), and sands may be calcareous, basaltic or siliceous in origin. Indeed, some very well-known beaches are made up of cobble or even bedrock, but here we restrict our discussion to sands. Biologically, beaches like all shorelines are ecotonal, where the terrestrial and nearshore ecosystems interact and overlap (Pennak, 1951). Here organisms interact at many different trophic levels including the microbes, which are found at the lowest levels. Little is known about obligate or specific bacteria, fungi or viruses in sand but we describe in this paper a diverse array of common indicators or pathogenic species known to occur in marine and freshwater sands.

Beach characteristics

At the local level, the beach itself is divided into zones (Figure 1) largely influenced by hydrology: the swash – where wave run-up and return occurs; intertidal – the horizontal extent of tides; the berm – a raised sand ridge of sand deposited by maximum wave run up; foreshore – area under influence of waves and tides; and backshore – landward side of the beach generally not affected by water except during storms. The berm tends to have the highest concentration of microbes due to filtering waves as they infiltrate the sand. The berm may also be a significant source of bacteria to bathing waters when waves or tides re-suspend stored material and return it to the nearshore (Whitman *et al.*, 2014). The back- and foreshore may also have significant input from animals especially seabirds. More research is needed on the transport of bacteria or fungi to or from the beach via groundwater but it is presumed that some microbes, especially the smaller ones (<5 microns) (Solo-Gabriele *et al.*, 1998), can pass through shallow groundwater relatively easily. Boehm *et al.* (2004) found that microbes could be potentially transported to the surf zone through tidally driven exchange of groundwater. de Sienes *et al.* (2008) hypothesized that the transport of nutrients via groundwater promotes the persistence and population replication of bacteria within the surf zone.

Sanitary assessments and sand remediation methods

A sanitary survey is the first step in evaluating pollution sources of a beach. This requires that the beach be viewed within the context of its beachshed. A 'beachshed' is 'a defined stretch of shoreline and the biogeochemical factors that influence it' (Whitman *et al.*, 2014). The extent of the beachshed and its potential influence on a beach should be considered before conducting a sanitary survey, developing a monitoring programme, estimating risk or conducting a microbial source tracking exercise. This would include groundwater, runoff, incoming streams, anthropogenic and natural faecal input, surface water dynamics, offshore influences and general water quality. A common mistake is to seek a single cause of poor sand quality or to finalize a survey after discovering obvious or superficial factors. Multiple sources of contamination are illustrated by the 63rd Street Beach, Chicago (Whitman *et al.*, 2001). Investigation has shown that FIOs

may be introduced into the foreshore from (1) direct defecation from birds, (2) accumulation of sand wave infiltration, (3) shoreward drift from re-suspended bacteria, (4) growth in stranded green algae *Cladophora* and (5) *in situ* growth of bacteria in moist sand (Halliday & Gast, 2011; Whitman *et al.*, 2014). Human faecal sources should be given high priority because they not only pose the greatest health risk, but may lend themselves to engineering and management solutions.

One of the obvious benefits of conducting a sanitary survey is discovering the source or factors contributing to sand contamination. Practical beach management and visitor education are good first steps. An adequate number of animal-proof rubbish receptacles will reduce disease-carrying wildlife, the spread of spoiled, discarded food and waste, and the associated vectors on the beach. Wrack management may help in minimizing the persistence of indicator microbes and some pathogens. Removal of nearby air pollution sources (in the case of airborne fungal spores) could help in the management of fungi in beach sands. As mentioned above, humans are sources of pathogenic microbes and limiting the number of people at a beach to prevent overcrowding will also avoid excessive microbial contributions within a congested environment (Brandão *et al.*, 2002). The number of beach visitors could possibly be controlled by limiting access and, in some cases, through the availability of parking. Encouraging visitors to shower before and after returning from the beach and hand washing before eating may reduce illnesses. One study showed that even just rinsing hands in the beach water greatly reduces bacterial adhering on hands (Whitman *et al.*, 2009). If birds contribute to poor sand quality an assessment would be necessary to determine whether deterrence is in line with local ecosystem preservation efforts. If so, there are techniques such as landscaping, sand grooming, and even the use of dogs and other tactics that can be used to deter birds. Nearby streams and margins may also contain high bacteria levels and swimmers should be encouraged avoid these areas. Often break-walls direct contaminants shoreward and managers may wish to have visitors avoid these areas (Byappanahalli *et al.*, 2015). Seepage or runoff on to the beach might also increase contamination. Thus, comprehensive sanitary surveys coupled with visitor education, adaptive management, and well-designed monitoring, will go far in providing safer enjoyment of recreational beaches.

In cases where a sanitary assessment and prevention is not enough, remediation may be needed. Remediation technologies include sand grooming (Kinzelman *et al.*, 2003; Kinzelman & McLellan, 2009), sand re-nourishment (Hernandez *et al.*, 2014) and treatment through chemical disinfectants and physical sterilization. Iodine spraying is one of the options currently employed in Portugal (Costa *et al.*, 2009) but theoretically other non-hazardous options exist such as sonication and high-energy light bathing (such as UV and infra-red radiation, ozone). As harmless for beach users as all these possibilities may be, the downstream pollution cleanup procedures inevitably will act both upon harmful contaminants and normal innocuous flora. Care should be taken that these methods are used only in extreme pollution events rather than as routine procedures.

General considerations for developing a monitoring programme

Designing a sampling programme begins with several preliminary questions: where should we sample, how do we sample,

and how often should we sample? Deciding where to monitor can be difficult; experience shows that contaminants arising from the water accumulate along the foreshore but substantial contamination may be occurring from surrounding areas, wildlife, pets and humans themselves higher on the beach. A programme that encompasses areas where visitors might encounter pathogens, ranging from backshore sand to the swash zone, may be appropriate at many beaches. Longitudinal transects along the beach at pre-selected intervals are preferred from a statistical standpoint, but known 'hotspots' should not be excluded from these studies.

Achieving representative sampling at beaches is difficult due to diverse inputs of microorganisms that create a heterogeneous community 'landscape'. Sand is arguably a more problematic matrix than water, as it is relatively less prone to mixing than water. Sample replication is essential since microbial distribution in sand is patchy. An alternative or companion strategy to replication is to collect many individual samples and mix them to create a composite sample, keeping in mind that pseudo-replicate sampling of composite samples should also be carried out to avoid placing undue weight on data based on a very small fraction of the sample. For example, Phillips *et al.* (2011a) collected 60 shallow core samples (each 2.5 cm diameter and 2.5 cm deep) along target transects. They combined these core samples, mixed them thoroughly, and utilized an aliquot for analysis.

Temperature and irradiation of exposed sand can vastly reduce levels of surface microbes. While deep within the sand, communities change due to more negative redox potential and lack of oxygen. Core samples are essential in order to provide an integrated survey of potential microbial pathogens. Sampling depths to 20–30 cm are the most practical and protective for the casual beach visitor. Once collected samples require extraction or elution prior to analysis (Boehm *et al.*, 2009b).

Decisions on the frequency of monitoring should be based on the amount of beach use, susceptibility to contamination, and also on cost. In temperate climates, beach use is very limited during cold weather, and sampling programmes may be minimized. It is known that sand microbial quality changes much less rapidly than the frequent temporal variability observed in water (Boehm *et al.*, 2002; Enns *et al.*, 2012). On the other hand, studies have shown correlations between water and sand FIO content, especially at the foreshore (Whitman & Nevers, 2003; Phillips *et al.*, 2011a, b). Generally, any monitoring programme should account for tradeoffs between visitor risk, budget, effectiveness, and accounting for expected variations in targeted microbes. Beach sand should be routinely sampled at least annually and whenever there has been an event such as a sewage release, major storm events or known seasonal events. Pre- and post-event monitoring where large crowds are anticipated might be considered. High-risk beaches, such as those potentially impacted by human sewage, require more frequent monitoring.

Sampling strategies based upon desired outcomes and integrating traditional approaches

Development of a monitoring programme for beach sand requires first that the desired outcomes of such a programme are clearly defined. An example of a two-part outcome is

protecting public health while allowing maximal use of the important resource represented by the beach. Another outcome might be to identify the source of the contamination so that mitigation of the contamination can occur. These outcomes require different sample collection and analysis approaches. For example, if public health is the main criteria, measurements should focus on the primary aetiological agent of disease or an indicator of the aetiological agent. Similarly if the focus is to identify the source of bacterial contamination, then MST should be the focus of sampling efforts. In most cases the aetiological agents and sources are inter-related and so monitoring programmes may be focused on identifying both.

Other factors that contribute towards different sand monitoring approaches include historical factors. When considering the two main categories of application, recreational water quality monitoring vs. environmental mycology, the approaches have been very different. In the USA, the modern era of recreational water quality monitoring was initiated by the amendment to the Federal Water Pollution Control Act of 1948, the Clean Water Restoration Act of 1966. Recreational water quality monitoring using FIO has been guided by a National Academy of Science initiative (NAS, 1972) with its inclusion within regulatory language by 1976 (EU, 1976; USEPA, 1976). The original purpose of beach monitoring during the 1970s was to detect whether the beach was impacted by sewage. This focus was initiated during an era when sewage was disposed to coastal areas with minimal treatment, resulting in direct impacts on local beach water quality from sewage outfalls. In this case, the concept of an indicator microbe, one found in human sewage, made perfect logical sense as indicator microbes are suited to track the aetiological agents of disease from human sewage, especially if the contamination is nearby. However, with improvements to sanitary infrastructures in most developed countries, direct impacts from human sewage have become less significant and other sources of contamination now make a larger contribution to overall beach water quality. These other sources include faeces from animals, wash-off of microbes from human bathers (Elmir *et al.*, 2007, 2009) and potential regrowth of indicator microbes in beach sands. As a result, the recreational water quality research community argues that the FIO concept is not meeting its originally intended purpose, particularly when the source of contamination is not human sewage. The scientific community has widely acknowledged the need to expand assessment to include additional measures of potential disease-causing agents due to the recognition that there are many other sources of microbes besides human sewage (Fujioka *et al.*, submitted). These additional measures can include alternative indicators such as *C. perfringens* and Bacteroidales which, unlike the traditional FIO, are unable to multiply in the environment under aerobic conditions.

In contrast, applications in environmental mycology have taken a much different approach. Environmental mycology is generally based upon an underlying notion that fungi are ubiquitous and are derived from many different sources. As such, studies focused on characterizing health threats from fungi in the environment typically include analyses of many fungi, usually as many as can practically be measured. This results in very cumbersome and time-consuming analyses, which in many cases are difficult to assess in terms of human health outcomes; in particular due to the lack of dose response data

between fungal exposures and human health outcomes. Ultimately, each individual has its exposure limit to fungi, which is dependent in part upon immune status. As a result, the environmental mycology community recognizes the need to simplify its approach for assessing the potential for transmission of infectious disease within beach settings. The 'tiered' approach described below represents a compromise between the recreational water quality community and those whose primary focus area is in environmental mycology.

A tiered approach for beach sand monitoring programmes

The 'first pass' of most monitoring programmes is culturable microbes. From the recreational monitoring community these microbes are normally FIOs. But from the environmental mycology perspective these may include total culturable fungi. The methods for analysing culturable microbes are relatively inexpensive, can be performed in laboratories with minimal specialized equipment and expertise, and in many cases have extensive historical use, providing context to new measurements. However, enumerating traditional FIOs, in particular, provides no information about contamination source in sand or water (recently reviewed in Harwood *et al.*, 2014), and the dearth of sand-related epidemiology studies leaves a substantial knowledge gap about human health risks (Whitman *et al.*, 2014).

In contrast to traditional regulatory approaches, the classic approach in environmental mycology has been to measure all fungi and FIO present in sand together with total counts of viable colony-forming units, as described in Brandão *et al.* (2002). More recently, however, environmental mycologists have started to focus on measures of specific microorganisms, more relevant in terms of public health (i.e. black moulds and keratinophilic fungi) (Sabino *et al.*, 2014a). This change is due to the lengthy and expensive practice of identifying all possible organisms. Thus, Environmental Mycology is gradually migrating towards the concept of measuring representative microbes or a fungal indicator microbe. As such the first tiered approach for monitoring the microbiological quality of water and/or beach sand should focus on measures of indicator microbes. This is consistent with current regulatory approaches used to assess recreational water for faecal contamination and is also consistent with the more recent evolving approaches in environmental mycology. In the case of fungi, the first tier analysis approach could include measures of total culturable fungi.

A second-tier approach using source-specific testing (microbial source tracking; MST) may be undertaken if information on the source of faecal contamination or fungi is required, either for mitigation or for risk assessment. Recent studies have estimated very different human health risks from faecal contamination originating from different host animals (Soller *et al.*, 2010, 2014) and so acceptable levels of indicator microbes should consider the potential source of the FIO. For fungi, species identification may help in identifying sources. For example, some fungi are associated with superficial infection, like the contagious *Tinea corporis* (ringworm). The *Trichophyton* and *Microsporum* genera include species of human, animal or soil origin (Badillet, 1973). Thus, identification of species may point to the contamination source for fungi in particular.

A third-tier approach would generally be most useful where many people are likely to be exposed (e.g. a crowded beach with human densities greater than 1 per square metre of beach sand area), and where MST has indicated the likely presence of high-risk faecal or fungal sources. The information provided by MST could be used to target particular pathogens that are known to be shed by the indicated host types. It can also require a great number of separate tests if several host types contribute to contamination. Alternatively, one can use a microarray approach, where hundreds of nucleic acid sequences representing FIOs, pathogens and MST markers can be simultaneously queried (Weidhaas *et al.*, 2014). An issue with microarray is that sample sizes are very small, so very efficient concentration methods that do not result in interference with nucleic acid hybridization are necessary, and these are still under development.

The second tier and third tier of analyses may be left for reference laboratories, laboratories with the technical expertise to analyse MST markers and pathogens from environmental samples, including capabilities for molecular analyses. These reference laboratories should assess regional specificities of indicator microorganisms and support or validate laboratories capable of basic analysis. For this level of reference, accreditation by ISO 17025 (ISO/IEC 17025, 2005) will ensure the technical proficiency of a laboratory and technical personnel. Inter-laboratory quality assessment schemes will capacitate the laboratories at this level to the point where they can validate first tier analytical approaches and provide expertise for non-standardized analytical methods. In this case, reproducibility and repeatability will ensure results independent of laboratories and technicians.

What should be measured?

The rapid pace of technological advances in the environmental detection and quantification of microbial targets has created what might be considered an embarrassment of riches. It engenders questions such as, should we test for a representative pathogen or two, or a broad suite of pathogens? Does testing need to be quantitative, or do binary results (plus/minus) (presence/absence) suffice? Should the focus be on one microbial type, such as viruses, or should the group of targets be broadened? Ideally, monitoring methods for beach sand monitoring should be inexpensive, provide instant, or at least same-day results, and be directly connected with human health outcomes (Figure 2). The current reality of monitoring methods is that there is no ready protocol that leads precisely to such an elegant outcome. Instead,




Fig. 2. Conceptual triangle for ideal characteristics of an indicator organism used for the first tier of screening sand quality at beaches.

compromise on one or more aspects of the ideal indicator is necessary and for this reason a tiered approach is recommended by the authors of this review as described above.

Over the past 60 years FIOs have proved a useful surrogate for measuring pathogens. The FIOs most commonly used for regulatory purposes are enterococci for marine waters (WHO, 2003) and *E. coli* for fresh water (USEPA, 2012). Other alternative indicator microbes that have been recommended include *Clostridium perfringens* (Fujioka & Shizumura, 1985; Roll & Fujioka, 1997; Boehm *et al.*, 2009a), Bacteroidales (Boehm *et al.*, 2009a) and coliphage (Havelaar *et al.*, 1993; Luther & Fujioka, 2004; Boehm *et al.*, 2009a). Although FIOs are utilized extensively worldwide, their limitations have been recognized (EU, 2006). Limitations include the fact that commonly used FIOs are invariably bacterial species, whereas the majority of the reported illnesses are believed to be caused by viruses (particularly norovirus), and analytical techniques suitable for routine use are poor at distinguishing between human and animal sources of bacterial FIOs. Data from the USA identified the following eight faecal pathogens as dominating waterborne illness: norovirus, rotavirus, adenovirus, *Cryptosporidium* spp., *Giardia lamblia*, *Campylobacter jejuni*, *Salmonella enterica* and *E. coli* O157:H7 (Mead *et al.*, 1999; Vital *et al.*, 2008). Investigations at beaches in Miami, FL, USA support the hypothesis that existing indicator microbes indirectly monitor several pathogens through common factors, at least in sand. However, for pathogens such as *Cryptosporidium* spp., *Giardia* spp. and enterovirus, generalizations about the predictive ability of indicator microbes must be treated with caution owing to the sparseness of data.

Soller *et al.* (2010) concluded that in fresh water, enteric viruses and *Giardia* appear to account for the vast majority of the observed swimming-associated GI illnesses, and when treated sewage effluent predominates, norovirus alone may represent the primary concern. The pre-eminence of noroviruses is supported by work by Public Health England, where norovirus dominated the identified cause of illness from consuming sewage-contaminated shellfish (Figure 3). Evaluating trends from 1991 through 2011, the aetiological agent most frequently identified as the cause of an outbreak was norovirus. This is particularly apparent for more recent years where detection technologies have been capable to identify the aetiological agent responsible for the outbreaks (David Lees, personal communication, CEFAS UK).

The European Union has sponsored an investigation (Virobathe) into analytical methods for viruses for possible incorporation into the 2020 revision of the European Union's (2006) Bathing Water Directive. The report of this work concluded that whilst adenoviruses were a possible control parameter, noroviruses were encountered too infrequently to be considered (EU, 2009; Wyer *et al.*, 2012). However the authors of Virobathe have since recommended to the European Commission that a viral pathogen standard should not be adopted on both analytical and public health grounds. They recommend instead that future risk continues to be managed through demonstrating connectivity to faecal sources, rather than proving that a pathogen is being excreted by the contributing population (Kay, 2015). Overall, because of their link to gastrointestinal disease and/or detection in recreational waters, viruses that should be considered when evaluating potential aetiological agents in sand include norovirus, adenovirus, rotavirus, enterovirus and hepatitis.




Fig. 3. Aetiology of bivalve shellfish associated infections as reported by Public Health England. Data are unpublished. The upper image provides the distribution of aetiological agents of disease for England and Wales Health Protection Agency, UK as provided by Craig Baker-Austin from the Centre for Environment, Fisheries and Aquaculture Science (CEFAS) Laboratory in Weymouth. DSP is Diarrhetic Shellfish Poisoning. The lower image provides the distribution of aetiological agents from 1991 through 2011 showing that among the aetiological agents identified, the one responsible for the majority of the outbreaks is norovirus. The data for the lower plot was provided by David Lees, CEFAS, UK.

The main fungi pathogenic to man and other mammals are found within the anamorphic (asexual reproductive phase) group. These fungi are saprophytic and occasionally pathogenic, and can be isolated from water, soil, animals and humans (Gomes *et al.*, 2008). The presence of certain ubiquitous fungal genera, such as *Alternaria*, *Acremonium*, *Aspergillus*, *Candida*, *Chaetomium*, *Cladosporium*, *Fusarium*, *Mucor*, *Penicillium*, *Phoma*, *Rhodotorula* and *Trichoderma* can be clinically important (Velonakis *et al.*, 2014; Wang *et al.*, 2014) due to their involvement in human diseases (Dolenc-Voljč, 2005; The Fungal Research Trust, 2011; Kaštelan *et al.*, 2014). Among these *Aspergillus*, *Candida*, *Fusarium* and dermatophytes like *Microsporum* and *Trichophyton* were identified as representing the majority of fungal isolates from clinical samples important for human health, and that these were appropriate for inclusion in beach sand quality legislation (Rees *et al.*, 1998). These genera should be included in beach sand quality legislation, as should allergenic airborne spore releasing moulds to protect those with respiratory disorders such as cystic fibrosis, asthma and reactive bronchitis. Beach specific studies in Egypt suggest that *Candida* and *Scopulariopsis* may be widespread (Migahed, 2003). Melanized fungi, such as black yeasts from the family *Herpotrichiellaceae* are also recognized as new emerging pathogens (de Hoog *et al.*, 2009). Fungi from genera *Cladophialophora*, *Exophiala*

and *Fonsecaea* are causative agents of chromoblastomycosis in subtropic and tropic regions and since they have been detected on wood, soil, plant material and in environments polluted with oil or creosote, their presence should also be evaluated in sand (Vicente *et al.*, 2008), especially on inland beaches. Since they are often detected in beach sand, species like *F. pedrosoi*, *F. monophora*, *C. bantiana* and *E. dermatitidis* should be included in future legislations of beach sand quality. Organisms of biosafety level 3 like *Cladophialophora bantiana* should also be considered. Figure 4 brings together data on fungi found in water and sand environments and those in clinical experiments to identify those most of concern in studies of recreational waters.

Technological advances in microbe measurement techniques

Improving technology has provided several new, but relatively expensive, methodologies for determining the safety of beach sand, e.g. quantitative PCR for specific pathogens or host-specific gene markers, or multi-target methods such as microarray or next-generation nucleic acid sequencing. However, it is important to note that molecular analysis will only reveal the presence of microbial genetic material, which does not always represent viable microbes. Culturable organisms must be capable of replication in order to be detected, a condition that is closer to an infective state than simply possessing genetic material. An example of this discrepancy is the 2012 US Environmental Protection Agency criteria for recreational water quality, which estimates 36 cases of gastroenteritis per 1000 exposed individuals in waters containing a geometric mean level of 35 culturable enterococci per 100 mL, but 470 'cell equivalents' per 100 mL by qPCR (USEPA, 2012). Conversely, some researchers have reported that FIO can enter a viable but non-culturable (VBNC) state in water from which they may be infective (Heim *et al.*, 2002; Lleò *et al.*, 2005; Boehm & Sassoubre, 2014), and these forms can be detected by molecular methods. Thus although molecular methods can detect specific microbes that are associated with human health outcomes, what they detect is different from culture-dependent methods, which adds to the complexity of adopting new methods based upon knowledge gained from older technologies. Few epidemiology studies have been conducted to determine whether new methods to detect specific pathogens are better assessments of human health outcomes than the century-old FIO paradigm. So a knowledge void exists that fosters indecision about which method(s) of assessment should be used. Although more information through epidemiology studies may be deemed better, economic constraints become quite important to regulatory agencies and the citizens who must ultimately pay for the testing.

Given the advent of new genomics approaches, measurements of the entire microbial community represent a potential new approach (see Application of Metagenomics to Assess Microbial Communities in Water and Other Environmental Matrices by Staley and Sadowsky in this issue). Many of the microbes living in sand have not been cultured and may not be culturable; thus a complete understanding of the microbial ecology of sand communities has not been possible. Furthermore, the lack of detection of unculturable and potentially infectious microorganisms has confounded monitoring efforts to protect public health. Consequently, metagenomic and 16S-amplicon-based studies to characterize microbial




Fig. 4. Presence of fungal genera in environmental and clinical studies. Blue circle includes fungal genera reported from seawater and ocean studies, orange circle presents fungi isolated from sand. In red circle there are genera reported as causative agents for human disease. The intersection of the circles includes fungi, isolated from two (seawater – beach sand, beach sand – clinical samples, clinical samples – seawater) or all three sampled sites (seawater – beach sand – clinical samples).

communities in sand, water and sediment habitats will offer great insight into the ecology of these systems (Lozupone & Knight, 2007; Staley *et al.*, 2015). In addition, characterization of these communities will provide a context for the role and relative abundance of potential pathogens, which has previously only been assessed using a relatively small number of ephemeral molecular targets (Aw & Rose, 2012). While in its infancy, this type of approach has been taken by Cui *et al.* (2013) who used 454 sequencing to find backshore sands in Hawaii had a more diverse community and contained different populations than other beach zones. Piggot *et al.* (2012) found that the Proteobacteria and Bacteroidetes dominate biofilm communities in South Florida beach sand, with microbial communities that vary by location within the tidal zones and in relation to water activity. More recently, Halliday *et al.* (2014) reported that sand at the high tide line, intertidal sand and adjacent water samples contained different overall bacterial communities, that there was some similarity in community composition between coastal water samples from two distant sites, and there was dissimilarity between bacterial communities from high tide and intertidal sands.

Recently, 16S rDNA amplicon analysis, using the Illumina HiSeq and MiSeq platforms was used to examine microbial

communities in sands obtained from an estuarine beach and a marine site in Tampa, FL; a freshwater lake in Saint Paul, MN; and Lake Michigan, near Chicago, IL (Whitman *et al.*, 2014). Among all three sites, Proteobacteria, Bacteroidetes, Firmicutes and Actinobacteria were the most abundant phyla, with families at all sites including Rhodobacteraceae, Flavobacteriaceae, Flammeovirgaceae and Campylobacteraceae. Sand from the marine sites had greater richness and higher non-parametric diversity indices than the other sites examined (Figure 5). More recently a programme ‘sands of the world’ has been initiated which utilizes, 16S amplicon sequencing and Illumina HiSeq to examine spatial and temporal diversity of bacterial as well as fungal communities in beach sands collected from fresh (Great Lakes) and salt water beaches around the world. More specifically, this project will characterize microbial diversity in sands from four beaches along the Great Lakes as well as marine beaches on both US coasts, the Gulf of Mexico, Hawaii, Japan and Korea. This information will be useful to determine what environmental factors control beach microbial communities and whether sands harbour unique or similar bacteria, archeal and fungal microbial communities that vary in some predictable manner. In addition, these data may give us insight into which microbial taxa are related to specific sand habitats.




Fig. 5. Most abundant phyla found in beach sand at (A) freshwater, temperate beaches (MN and IL) and (B) marine, tropical beaches (FL). A total of 36 phyla were identified by sequencing of the V6 hypervariable region of the 16S rDNA among all samples.

SCIENTIFIC RESEARCH NEEDS

The research community recognizes that the traditional FIO paradigm is limited. This paradigm has served the public health community well for more than a century, especially in areas directly impacted by sewage. In developed countries, direct sewage impacts on beaches are the exception rather than the rule. In beaches impacted by non-point source contamination, the relationships between FIOs and human health outcomes are not well understood; therefore it is unclear whether continued monitoring for FIOs in these cases is relevant. Regardless of the source of contamination, the incidence of illness is higher for beach bathers relative to non-bathers (Colford *et al.*, 2007; Fleisher *et al.*, 2010; Sinigalliano *et al.*, 2010) and also for beach users who play in the sand relative to those who do not play in the sand (Heaney *et al.*, 2012). It is also recognized that sand can serve as a reservoir of pathogenic microbes, including faecal pathogens of human or animal source, as well as fungi, which are generally considered to be associated with environmental sources. Thus, pathogens may be present and transmitted in the beach environment, even in the absence of sewage contamination. We recommend research to establish a cause and effect relationship for infectious disease contracted within beach environments. First and foremost is the identification of the aetiological agent of disease followed by establishing stronger links between environmental monitoring parameters and human health risk. Specific recommendations include:

- (1) *Identification of the aetiological agent(s) of disease.* A better understanding is needed of the aetiological agents responsible for the majority of disease attributed to recreational water and sand contact. Identifying the aetiological agent will provide a stronger mechanistic understanding for disease transmission in beach settings. With this understanding, effective control and monitoring programmes can be implemented. To narrow the list of possible aetiological agents, public health data should be examined. A preliminary assessment of reportable

diseases in the EU and USA (Table 1) suggest that significant pathogens that have potential sand reservoirs include GI pathogens *Salmonella*, *Shigella*, verotoxin-producing *E. coli*, *Campylobacter*, *Cryptosporidium*, *Cyclospora*, *Vibrio*, *Giardia*, hepatitis A and *Listeria*. Those associated with sediment and water reservoirs include *Yersinia*, *Leptospira* and *Tularaemia*. Although a list of reportable diseases is available, low-level self-limiting diseases (e.g. GI illness, mild skin infections and mild respiratory infections) are usually not reported and the aetiological agents are typically not measured. Wheeler *et al.* (1999) found that the incidence rate for mild gastroenteritis was under-reported by a factor of 31. Thus numbers listed in Table 1 may significantly underestimate the incidence of reportable diseases. Concerted efforts are needed to encourage clinical practitioners to more often request an evaluation of etiological agents of disease to better track them. Because of unique clinical manifestations, the tracking of fungal infections may be more easily accomplished as opposed to GI infections.

- (2) *Quantitative microbial risk assessment (QMRA).* QMRA methods should be utilized to specifically estimate public health risks from various pathogens (bacteria, fungi, viruses, protozoa, helminths) in beach sand, which can transmit diseases by various exposure routes (contact, ingestion, inhalation). QMRA methods are generally less expensive and less time consuming than epidemiological studies; however, in some cases relationships needed in calculating risks and disease rates are not available (e.g. dose-response relationships for some microbes). An assessment must be made as to which pathogens in beach sand can and cannot be evaluated by QMRA. Preliminary assessments (Shibata & Solo-Gabriele, 2012) have identified the need for dose-response estimates for fungi and helminths. Moreover, to obtain a better estimate of skin-related ailments, the impacts of wounds should be evaluated on the dose-response of various aetiological agents known to cause skin disease. In some cases, such as for helminths,

Table 1. Number of cases of reportable diseases and incidence rates in the EU and US. Data from ECDC (2011a, b, 2012, 2014a, b) and CDC (2013, 2014). Incidence rates based upon population estimates of 506–503 million and 309–314 million for the EU and US, respectively.

| Disease | Year | European Union | | United States | |
|-----------------------------|------|--------------------|--|--------------------|--|
| | | Total no. of cases | Incidence rate (per 100,000 inhabitants) | Total no. of cases | Incidence rate (per 100,000 inhabitants) |
| Salmonellosis | 2010 | 90,764 | 17.9 | 54,424 | 17.6 |
| | 2011 | 88,577 | 17.5 | 51,887 | 16.7 |
| | 2012 | 87,719 | 17.4 | 53,800 | 17.1 |
| Shigellosis | 2010 | 6839 | 1.35 | 14,786 | 4.79 |
| | 2011 | 6655 | 1.32 | 13,352 | 4.30 |
| | 2012 | 6643 | 1.32 | 15,283 | 4.86 |
| VTEC infection ^a | 2010 | 3748 | 0.74 | 5476 | 1.77 |
| | 2011 | 9661 | 1.91 | 6047 | 1.95 |
| | 2012 | 5954 | 1.18 | 6463 | 2.06 |
| Listeriosis | 2010 | 1686 | 0.33 | 821 | 0.27 |
| | 2011 | 1538 | 0.30 | 870 | 0.28 |
| | 2012 | 1692 | 0.34 | 727 | 0.23 |
| Legionellosis | 2010 | 5854 | 1.16 | 3346 | 1.08 |
| | 2011 | 4449 | 0.88 | 4202 | 1.35 |
| | 2012 | 5856 | 1.16 | 3688 | 1.17 |
| Vibriosis, non-cholera | 2010 | – | – | 846 | 0.27 |
| | 2011 | – | – | 832 | 0.27 |
| | 2012 | – | – | 1111 | 0.35 |
| Campylobacteriosis | 2010 | 218,957 | 43.28 | – | – |
| | 2011 | 227,803 | 45.02 | – | – |
| | 2012 | 218,153 | 43.37 | – | – |
| Yersiniosis | 2010 | 6614 | 1.31 | – | – |
| | 2011 | 6810 | 1.35 | – | – |
| | 2012 | 6110 | 1.21 | – | – |
| Leptospirosis | 2010 | 822 | 0.16 | – | – |
| | 2011 | 685 | 0.14 | – | – |
| | 2012 | 778 | 0.15 | – | – |
| Tularaemia | 2010 | 888 | 0.18 | 124 | 0.04 |
| | 2011 | 755 | 0.15 | 166 | 0.05 |
| | 2012 | 1003 | 0.20 | 149 | 0.05 |
| Cryptosporidiosis | 2010 | 822 | 0.16 | 8944 | 2.90 |
| | 2011 | 685 | 0.14 | 9250 | 2.98 |
| | 2012 | 778 | 0.15 | 7956 | 2.53 |
| Cyclosporiasis | 2010 | – | – | 179 | 0.06 |
| | 2011 | – | – | 151 | 0.05 |
| | 2012 | – | – | 123 | 0.04 |
| Giardiasis | 2010 | 17,130 | 3.39 | 19,811 | 6.42 |
| | 2011 | 16,475 | 3.26 | 16,747 | 5.39 |
| | 2012 | 16,424 | 3.27 | 15,178 | 4.83 |
| Hepatitis A | 2010 | 13,471 | 2.66 | 1670 | 0.54 |
| | 2011 | 12,706 | 2.51 | 1398 | 0.45 |
| | 2012 | 13,156 | 2.62 | 1562 | 0.50 |

^aVTEC, verotoxin-producing *Escherichia coli*, the Shigella-like toxin (includes toxin producing O157). Also known as shiga toxin producing *E. coli* (STEC).

larvae can penetrate the skin without requiring a wound for entry.

- (3) *Epidemiological studies.* Epidemiological studies measure disease in the exposed population and are the method of choice for establishing the link between human health (GI illness, acute febrile respiratory illness, skin ailments, ear and eye infections) and environmental factors. Since this method is expensive, work intensive and time consuming, it should be used at selected beach sites, bearing in mind the slow onset of infections caused by fungi and parasites. Epidemiologists should determine if study designs can specifically measure pathogen disease rate and measure exposure to the sand by three separate

routes (contact, ingestion, inhalation). Controlled cohort or randomized trial studies – similar in aim to those undertaken in water – are needed to better quantify disease risk from exposure to sand.

- (4) *Evaluate alternatives to FIO for beach monitoring programmes.* Although measurements of FIOs should not be discontinued because many beaches are susceptible to sewage contamination, their usefulness needs to be reassessed. The most common FIOs (*E. coli*, enterococci) fail to fulfil the following three scientifically based ideal criteria or cellular properties of the indicator bacteria, which are required to ensure that the numbers of FIO will correspond to the numbers of sewage-borne

pathogens in the water samples tested (Bonde, 1966; Yates, 2007): (a) The indicator should be consistently and exclusively associated with a source of human pathogens (e.g. human faeces/sewage); (b) FIO should not be able to multiply under environmental conditions because they would no longer track some sewage-borne pathogens (viruses, protozoa), which presumably do not multiply in the environment; (c) their resistance or survival characteristics to environmental conditions and to wastewater treatment processes should be similar to that of pathogens. Without these cellular characteristics and multiplication in human intestinal sources, the FIO would not track pathogens and subsequently human health. Research is needed to evaluate the reliability and feasibility of monitoring sand for alternative faecal indicators that meet the above three ideal criteria and thus appropriately track pathogens. Since noroviruses are one of the more likely aetiological agents for water-borne transmission of diseases during recreational uses of water, there is a pressing need to develop appropriate viral indicator(s). Viral indicators could include bacteriophages (bacterial viruses), which have similar size, chemical composition and survival characteristics as human enteric viruses. Phages, which are considered promising indicators of human enteric viruses, are F-specific RNA bacteriophages (Havelaar *et al.*, 1993; Luther & Fujioka, 2004), phages of enterococci bacteria (Santiago-Rodriguez *et al.*, 2013) or phages of *Bacteroides* (Ebdon *et al.*, 2007; McMinn *et al.*, 2014). In addition, it will be important to evaluate beach sand for the alternative FIO (*C. perfringens*) because it is a conservative indicator of sewage contamination (Fujioka & Shizumura, 1985; Roll & Fujioka, 1997). Because of their persistence as spores, the use of *C. perfringens* may be most useful in areas where currents dilute and remove existing contamination.

(5) *Develop techniques for detection and quantification of microbe levels.* There is a need for improved cultivation methods for detection of fungi, viruses, helminths, protozoa and bacteria in environmental samples. There is a need to determine if the method to detect a specific pathogen or class of pathogens is feasible and reliable for monitoring purposes. If a pathogen detection method is not feasible, then a feasible and reliable indicator monitoring method should be implemented, which should provide data on the quantity and infectivity for that pathogen or that class of pathogens. In this regard, culturable methods provide information on the theoretical infectivity of the pathogen and this kind of data can be used to determine public health risks. Currently, many molecular methods have been developed to rapidly and reliably detect specific pathogens. The limitation of this method is that it does not differentiate between dead and living pathogens. As a result, public health assessments must be based on some assumptions. The value of molecular methods is that they can be used to confirm the presence or absence of specific pathogens in beach sand, regardless of their viability. Improved molecular techniques for the detection of medically important fungi in sand are needed. Also, the ecological role of fungi in coastal reservoirs such as beach sand is little understood (Migahed, 2003) and needs to be investigated. In securing better protection, unnecessary complexity in monitoring is to be avoided. Concern has been expressed on the cost

- burden of monitoring – particularly in developing regions – of even the existing criteria (WHO, 2003). Against this must be balanced a better cost-benefit balance of targeted improvements and the avoidance of expenditure on ineffective measures undertaken simply to meet flawed criteria (Kay *et al.*, 1999).
- (6) *Pathogen levels and survival in sands.* A more complete picture of the types and levels of pathogens in sand is needed, including a focus on evaluating their geographic, spatial and temporal distribution. Multiple studies have documented *E. coli* and enterococci reservoirs in sand, but few studies have concurrently measured pathogens to determine if *E. coli* and enterococci are indicative of faecal pollution that carries pathogens, or uncoupled from their original source through prolonged survival or growth. The general consensus among researchers is that pathogens do not multiply in the environment. So prolonged survival or growth of *E. coli* and enterococci would result in their presence in the absence of pathogens. New research is needed that can provide tools to determine whether FIO are indicative of pathogens within sand environments. The sources of faecal pollution will largely determine the types of pathogens that may be present in sand. Faecal pollution can be deposited directly in sand through outfall runoff or wildlife, or may be delivered through contaminated water. Understanding how faecal pollution and its co-occurring pathogens are modulated in both the sand and water environment can guide the types of indicators or pathogens chosen. Further, gaining a more complete picture of the pathogens that persist in the beach sand and the causative agents for disease, will guide choices of indicators for monitoring and improve assessments of risk.
- (7) *Develop tools to identify sources.* Although considerable advances have been made through MST, more work is needed to identify and approve/agree methods that distinguish between human and non-human sources. There is a need to understand differences in risk among these sources. An improved understanding of the relative risk of faecal contamination from human and other sources is necessary to establish acceptable levels of FIOs in the environment.
- (8) *Regulations are to reflect microbial sources.* Through application of the Annapolis Protocol the WHO has confirmed the need to consider all potential sources of pathogens, not only those from faecal point sources, an approach endorsed by the Rotorua declaration of 2011 (IWA, 2011). The beach environment is an important contributor to water, both through the retention, and possible regrowth, of FIOs within the sand matrix and beach wrack, but also for the presence in sand of non-faecal pathogens including fungi, protozoa and parasites. Forthcoming developments of regulatory standards need to reflect this evolving understanding of microbial sources, the pathogens they contain, and the associated health risks.
- (9) *Develop reliable sand collection methods designed to recover average pathogen loads* for a given beach site or at a specific site where pathogens are suspected. Since pathogen contamination at sandy beaches is expected to be patchy, sand samples from multiple sites should be collected and pooled to determine average concentrations of pathogens in sandy areas. However, targeted sampling

(within decaying algae, bird roosts, swash zone, public showers, land-based discharges onto beach sand) should also be conducted where pathogen contamination is suspected. These contaminated patches of sand are good sources for microbial source tracking analysis.

- (10) *Determine beach sand quality at freshwater vs. marine beaches.* Fresh and marine beach sands have been reported to be contaminated by different sources of FIO and pathogens (Whitman *et al.*, 2014). For example, different species of algae have been reported to contaminate shorelines of freshwater beaches (Byappanahalli *et al.*, 2003) vs. marine beaches (Imamura *et al.*, 2011). In this regard, decay of *Cladophora* in freshwater beaches has been reported to allow for the growth of FIO and other pathogenic bacteria (Ishii *et al.*, 2006). Therefore, sand at freshwater beaches and sand at marine beaches can be expected to differ with respect to sources of contamination, types of pathogens and their survival characteristics.
- (11) *Assess beach sand quality based on contamination by land- and air-based discharges,* which are known to be major factors that determine the sources as well as persistence of microorganisms in beach sand. High rainfall patterns cause land-based discharges such as streams and storm drains and may include discharges from agricultural and animal raising facilities. Even beaches with low rainfall can receive substantial urban discharges (storm drains) that include effluents from sewage and industrial facilities as well as discharges of human faeces directly into storm drains. The impacts of these land-based discharges affect the quality of beach sand differently at different beaches and must be assessed as site-specific factors. Air transportation plays the same role for sporulating microorganisms.
- (12) *Assess standardized methods to recover and disinfect FIO and pathogens from silica-based vs. calcium carbonate-based sands.* Both silica sand and calcium carbonate sand are chemically stable sand particles. However, calcium carbonate sand is more reactive and dissolves in dilute acid more readily compared with silica sand. As a result, each may not react similarly to all reagents and may have different influences on survival of microorganisms. There is a need to determine the impact of silica-based and calcium carbonate-based sands on survival characteristics of microbes, on the use of reagents to recover microbes from sand and in the use of chemical reagents to disinfect these two types of beach sand. As a corollary to disinfection, efforts are needed to evaluate the impacts of sand disinfection on the microbial ecosystems and at upper trophic levels.

SUMMARY AND CONCLUSIONS

There is compelling scientific evidence that beaches, through their sands, are a significant contributor to the pathogen load to which beach users are exposed. Many beach epidemiological studies have focused on the impacts of bathing. At beaches that are not impacted by sewage effluent, the source of pathogens originates from the local beach site itself and includes human visitors at the beach, animals, local runoff and the release of microbes from sand. The microbes released from sand can include native microbes (autochthonous) or those that have been deposited from outside sources

(allochthonous). Studies have identified the presence of pathogenic microbes in beach sand and have identified factors other than point source pollution that contribute to their presence (e.g. moisture, wrack, wildlife, domestic animals, beach morphology, currents). More recent epidemiological studies have shown that children who play in sand are subject to higher rates of illness relative to those who do not play in the sand. Thus beach sand can serve as a vehicle for disease transmission, either through direct sand contact containing microbes or indirectly through contact with water containing microbes washed off from sand. Given the ability of sand to harbour microbes, we recommend the inclusion of sand measurements in all beach monitoring programmes.

We provide a series of recommendations for beach monitoring programmes that begin by identifying designated recreational beach areas, beach sanitary surveys inclusive of remediation methods, general considerations for monitoring programmes, and a sampling strategy based upon desired outcomes. Given the large number of potential aetiological agents of disease, a tiered approach is recommended for beach sand monitoring. The approach should begin with measures of FIOs and/or total culturable fungi followed by microbes with potential for source tracking. For microbes transmitted via faecal-oral routes, sources should be identified through microbial source tracking. For fungi, specific species can be used to help identify sources. For the third tier, the specific aetiological agent responsible for disease should be measured. No beach epidemiological study to date (whether focused on water or sand) has directly measured pathogens in human subjects to confirm the aetiological agent of disease. According to QMRA methods, the most likely aetiological agents for faecal-oral beach illnesses include norovirus and *Giardia*. So far, measurements of fungi have not been included in beach epidemiological studies. However, given their propensity in beach sands, agents that we recommend for inclusion in the third tier of measurements are pathogenic *Aspergillus* sp., *Candida* sp., *Microsporum* sp., and *Trichophyton* sp. Measurement techniques include culture-based methods and quantitative PCR. An alternative approach can include the measure of the beach metagenome as a means of assessing the microbial ecological factors that may facilitate the presence of pathogens.

Considerable evidence exists that sand can serve as a reservoir of enteric microorganisms and fungi, which can be vehicles of disease transmission at beach sites. Current policies worldwide, at both national and international levels, give scant regard to the impact of sands on the health of users of beaches. We recommend that sand quality measures should be considered with some urgency for inclusion in regulatory programmes aimed at protecting recreational beach user health. Contaminated sands present health and economic costs that can and should be known by decision makers, communities and by individuals. Available evidence should be evaluated by both scientists and regulators with a view to filling the data gaps outlined here, which should be followed by sound policy development for safeguarding public health.

ACKNOWLEDGEMENTS

The authors acknowledge the hard work and investment of *Eruditus, Relações Públicas e Serviços* for the logistics involved in organizing the round table at TEMP2014 which

originated this paper. We acknowledge Katsia Kprzybyla-Kelly for providing valuable suggestions. We thank the ECDC for providing data reported in Table 1 on notifiable infections within Europe. Data provided by ECDC extracted from The European Surveillance System – TESSy. The views and opinions of the authors expressed herein do not necessarily state or reflect those of the ECDC. The accuracy of the authors' statistical analysis and the findings they report are not the responsibility of ECDC. ECDC is not responsible for conclusions or opinions drawn from the data provided. ECDC is not responsible for the correctness of the data and for data management, data merging and data collation after provision of the data. ECDC shall not be held liable for improper or incorrect use of the data.

FINANCIAL SUPPORT

We thank the Fulbright Foundation for support of Dr Harwood. Participation by H. Solo-Gabriele is associated with the University of Miami Center for Oceans and Human Health (NSF oCE0432368/0911373/1127813 and NIEHS P50 ES12736). Raquel Sabino was financially supported by a fellowship from Fundação para a Ciência e Tecnologia (FCT) Portugal (contract SFRH/BPD/72775/2010).

REFERENCES

- Abdelzaher A., Solo-Gabriele H., Phillips M., Elmir S. and Fleming L. (2013) An alternative approach to water regulations for public health protection at bathing beaches. *Journal of Environmental and Public Health* 2013, 1–9.
- Abdelzaher A., Wright M., Ortega C., Solo-Gabriele H.M., Miller G., Elmir S., Newman X., Shih P., Bonilla J.A., Bonilla T.D., Palmer C.J., Scott T., Lukasik J., Harwood V.J., McQuaig S., Sinigalliano C., Gidley M., Plano L.R.W., Zhu X., Wang J.D. and Fleming L.E. (2010) Presence of pathogens and indicator microbes at a non-point source subtropical recreational marine beach. *Applied and Environmental Microbiology* 76, 724–732.
- Alastruey-Izquierdo A., Cuenca-Estrella M., Monzón A., Mellado E. and Rodríguez-Tudela J.L. (2008) Antifungal susceptibility profile of clinical *Fusarium* spp. isolates identified by molecular methods. *Epub* 61, 805–809. doi: 10.1093/jac/dkn022.
- Alcazar-Fuoli L., Mellado E., Alastruey-Izquierdo A., Cuenca-Estrella M. and Rodríguez-Tudela J.L. (2008) *Aspergillus* section *Fumigati*: antifungal susceptibility patterns and sequence-based identification. *Antimicrobial Agents and Chemotherapy* 52, 1244–1251.
- Alm E.W., Burke J. and Spain A. (2003) Fecal indicator bacteria are abundant in wet sand at freshwater beaches. *Water Research* 37, 3978–3982.
- Anderson J.H. (1979) *In vitro* survival of human pathogenic fungi in Hawaiian beach sand. *Sabouraudia* 17, 13–22.
- Ashbolt N.J., Amézquita A., Backhaus T., Borriello P., Brandt K.K., Collignon P., Coors A., Finley R., Gaze W.H., Heberer T., Lawrence J.R., Larsson D.G., McEwen S.A., Ryan J.J., Schönfeld J., Silley P., Snape J.R., Van den Eede C. and Topp E. (2013) Human Health Risk Assessment (HHRA) for environmental development and transfer of antibiotic resistance. *Environmental Health Perspectives* 121, 993–1001.
- Aw T.G. and Rose J.B. (2012) Detection of pathogens in water: from phylotypes to qPCR to pyrosequencing. *Current Opinion in Biotechnology* 23, 422–430.
- Badillet G. (1973) *Les Dermatophytes: atlas clinique et biologique*, 3^e ed. Ed. Paris: Editions Varia.
- Balajee S.A., Gribskov J.L., Hanley E., Nickle D. and Marr K.A. (2005) *Aspergillus lentulus* sp. nov., a new sibling species of *A. fumigatus*. *Eukaryotic Cell* 4, 625–632.
- Bik M.H., Halanaych M., Sharma J. and Thomas K. (2012) Dramatic shifts in benthic microbial eukaryote communities following the deep water horizon oil spill. *PLoS ONE* 7, 1–6.
- Boehm A., Ashbolt N., Colford J. Jr, Dunbar L., Fleming L., Gold M., Hansel J., Hunter P., Ichida A., McGee C., Soller J. and Weisberg S. (2009a) A sea change ahead for recreational water quality criteria. *Journal of Water and Health* 7, 9–20.
- Boehm A.B., Grant S.B., Kim J.H., Mowbray S.L., McGee C.D., Clark C.D., Foley D.M. and Wellman D.E. (2002) Decadal and shorter period variability of surf-zone water quality at Huntington beach, California. *Environmental Science and Technology* 36, 3885–3892.
- Boehm A.B., Griffith J., McGee C., Edge T.A., Solo-Gabriele H.M., Whitman R., Cao Y., Getrich M., Jay J.A., Ferguson D., Goodwin K.D., Lee C.M., Madison M. and Weisberg S.B. (2009b) Faecal indicator bacteria enumeration in beach sand: a comparison study of extraction methods in medium to coarse sands. *Journal of Applied Microbiology* 107, 1740–1750.
- Boehm A. and Sassoubre L. (2014) Enterococci as indicators of human faecal pollution. In Gilmore M.S., Clewell D.B., Ike Y. and Shankar N. (eds) *Enterococci: from commensals to leading causes of drug resistant infection*. Boston, MA: Massachusetts Eye and Ear Infirmary.
- Boehm A.B., Shellenbarger G.G. and Paytan A. (2004) Groundwater discharge: potential association with faecal indicator bacteria in the surf zone. *Environmental Science & Technology* 38, 3558–3566.
- Bonde G.J. (1966) Bacteriological methods for estimation of water pollution. *Health Laboratory Science* 3, 124–128.
- Bonilla T.D., Nowosielski K., Cuvelier M., Hartz A., Grenn M., Esiobu N., McCorquodale D.S., Fleisher J.M. and Rogerson A. (2007) Prevalence and distribution of faecal indicator organisms in South Florida beach sand and preliminary assessment of health effects associated with beach sand exposure. *Marine Pollution Bulletin* 54, 1472–1482.
- Brandão J., Wergikosky B., Rosado C., Noronha G., Verissimo C., Falcão M.L., Giralde A., Simões M. and Rebelo H. (2002) *Qualidade Microbiológica de areias de Praias Litoriais – Relatório Final*. Alfragide: Instituto do Ambiente, Associação Bandeira Azul da Europa. Available at: <http://repositorio.insa.pt/handle/10400.18/232>
- Byappanahalli M.N., Nevers M.B., Whitman R.L., Ge Z., Spoljaric A., Shively D. and Przybyla-Kelly K. (2015) Wildlife, urban inputs, and landscape configuration are responsible for degraded swimming water quality at an embayed beach. *Journal of Great Lakes Research* 41, 156–163.
- Byappanahalli M.N., Shively D.A., Nevers M.B., Sadowsky M.J. and Whitman R.L. (2003) Growth and survival of *Escherichia coli* and enterococci populations in the macro-alga *Cladophora* (Chlorophyta). *FEMS Microbiology Ecology* 46, 203–211.
- Carneiro H.A., Coleman J.J., Restrepo A. and Mylonakis E. (2011) *Fusarium* infection in lung transplant patients: report of 6 cases and review of the literature. *Medicine (Baltimore)* 90, 69–80.
- CDC (2013) Summary of notifiable diseases — United States, 2011. *Morbidity and Mortality Weekly Report* 60, 8–107.

- CDC (2014) Summary of notifiable diseases — United States, 2011. *Morbidity and Mortality Weekly Report* 61, 1–121.
- CDC (2015) Antifungal resistance. <http://www.cdc.gov/fungal/antifungal-resistance.html>.
- Chowdhary A., Kathuria S., Xu J. and Meis J.F. (2013) Emergence of azole-resistant *Aspergillus fumigatus* strains due to agricultural azole use creates an increasing threat to human health. *PLoS Pathogens* 9, e1003633. doi: 10.1371/journal.ppat.1003633.
- Colford J.M. Jr., Wade T.J., Schiff K.C., Wright C.C., Griffith J.F., Sandhu S.K., Burns S., Sobsey M., Lovelace G. and Weisberg S.B. (2007) Water quality indicators and the risk of illness at beaches with nonpoint sources of fecal contamination. *Epidemiology* 18, 27–35.
- Costa I., Bértolo E., Bagulho A. and Duarte A. (2009) Monitoring of fungi flora on beaches sand along the Estoril coast, Portugal. 4th Trends in Medical Mycology Meeting (TIMM), 18–21 October, Athens, Greece.
- Cui H., Yang K., Pagaling E. and Yan T. (2013) Spatial and temporal variation in enterococcal abundance and its relationship to the microbial community in Hawaii beach sand and water. *Applied and Environmental Microbiology* 79, 3601–3609.
- de Hoog G.S., Guarro J., Gene J. and Figueras M.J. (2009) *Atlas of clinical fungi*. Utrecht/Reus: Centra al bureau Voor Schimmelcultures, Universitat Rovira I Virgili.
- de Kraker M.E., Davey P.G., Grundmann H. and BURDEN Study Group (2011) Mortality and hospital stay associated with resistant *Staphylococcus aureus* and *Escherichia coli* bacteremia: estimating the burden of antibiotic resistance in Europe. *PLoS Medicine* 8, e1001104.
- de Sieyes N.R., Yamahara K.M., Layton B.A., Joyce E.H. and Boehm A.B. (2008) Submarine discharge of nutrient-enriched fresh groundwater at Stinson Beach, California is enhanced during neap tides. *Limnology and Oceanography* 53, 1434–1445.
- Diwan V., Stålsby L.C. and Tamhankar A.J. (2013) Seasonal and temporal variation in release of antibiotics in hospital wastewater: estimation using continuous and grab sampling. *PLoS ONE* 8, e68715.
- Dolenc-Voljč M. (2005) Dermatophyte infections in the Ljubljana region, Slovenia. *Mycoses* 48, 181–186.
- Dunhill I., Nicholson N., Long R., Porter J. and Rahman A. (2013) *Faecal indicator organism populations in wrack piles*. Bristol: Evidence Directorate, Environment Agency.
- Ebdon J., Muniesa M. and Taylor H.D. (2007) The application of a recently isolated strain of *Bacteroides*(GB-124) to identify human sources of faecal pollution in a temperate river catchment. *Water Research* 41, 3683–3690.
- Edge T.A. and Hill S. (2007) Multiple lines of evidence to identify the sources of fecal pollution at a freshwater beach in Hamilton Harbour, Lake Ontario. *Water Research* 41, 3585–3594.
- ECDC (2011a) *Annual report of the Director 2011*. Stockholm: European Centre for Disease Prevention and Control. doi: 10.2900/25172, 71 pp.
- ECDC (2011b) *Annual report of the Director 2010*. Stockholm: European Centre for Disease Prevention and Control. doi: 10.2900/52932, 64 pp.
- ECDC (2012) *Annual report of the Director 2012*. Stockholm: European Centre for Disease Prevention and Control. doi: 10.2900/7911, 77 pp.
- ECDC (2013) Antimicrobial resistance surveillance in Europe. *Annual report of the European Antimicrobial Resistance Surveillance Network (EARS-Net)*. Stockholm: European Centre for Disease Prevention and Control.
- ECDC (2014a) *Annual epidemiological report 2014 – Food and water-borne diseases and zoonoses. Surveillance report*. Stockholm: European Centre for Disease Prevention and Control.
- ECDC (2014b) *Annual epidemiological report 2014 – respiratory tract infections. Surveillance Report*. Stockholm: European Centre for Disease Prevention and Control.
- Efstratiou M. and Velegraki A. (2009) Recovery of melanised yeasts from Eastern Mediterranean beach sand associated with the prevailing geochemical and marine flora patterns. *Medical Mycology* 48, 413–415.
- Elmir S.M., Shibata T., Solo-Gabriele H.M., Sinigalliano C.D., Gidley M.L., Miller G., Plano L., Kish J., Withum K. and Fleming L. (2009) Quantitative evaluation of enterococci and Bacteroidales released by adults and toddlers in marine water. *Water Research* 43, 4610–4616.
- Elmir S.M., Wright M.E., Solo-Gabriele H.M., Abdelzaher A., Fleming L.E., Miller G., Rybolowik M., Shih M.-T.P., Pillai S.P., Cooper J.A. and Quaye E.A. (2007) Quantitative evaluation of bacteria released by bathers in a marine water. *Water Research* 41, 3–10.
- Enns A.A., Vogel L.J., Abdelzaher A.M., Solo-Gabriele H.M., Plano L.R.W., Gidley M.L., Phillips M.C., Klaus J.S., Piggot A.M., Feng Z., Reniers A.J.H.M., Haus B.K., Elmir S.M., Zhang Y., Jimenez N.H., Abdel-Mottaleb N., Schoor M.E., Brown A., Khan S.Q., Dameron A.S., Salazar N.C. and Fleming L.E. (2012) Spatial and temporal variation in indicator microbe sampling is influential in beach management decisions. *Water Research* 46, 2237–2246.
- Esiobu N., Mohammed R., Echeverry A., Green M., Bonilla T., Hartz A., McCorquodale D. and Rogerson A. (2004) The application of peptide nucleic acid probes for rapid detection and enumeration of eubacteria, *Staphylococcus aureus* and *Pseudomonas aeruginosa* in recreational beaches of S Florida. *Journal of Microbiological Methods* 57, 157–162.
- EU (2009) European Union FP6 Frame Work Virobathe Project reference 513648. Methods for the concentration and detection of adenoviruses and noroviruses in European bathing waters with reference to the revision of the Bathing Water Directive 76/160/EEC, accessible report. Brussels: European Union. http://cordis.europa.eu/publication/rcn/12840_en.html.
- EU 1976 Council Directive (1976) *Council Directive of 8 December 1975 concerning the quality of bathing water (76/160/EEC)*. Brussels: European Communities.
- EU 2006 European Union (EU) (2006) Directive 2006/7/EC of the European Parliament and of the Council of 15 February 2006 concerning the management of bathing water quality and repealing directive 76/160/EEC. *Official Journal of the European Union* 64, 37–51.
- Feng Z., Reniers A., Haus B.K. and Solo-Gabriele H.M. (2013) Modeling sediment-related enterococci loading, transport, and inactivation at an embayed nonpoint source beach. *Water Resources Research* 49, 693–712.
- Finley R.L., Collignon P., Larsson D.G., McEwen S.A., Li X.Z., Gaze W.H., Reid-Smith R., Timinouni M., Graham D.W. and Topp E. (2013) The scourge of antibiotic resistance: the important role of the environment. *Clinical Infectious Diseases* 57, 704–710.
- Fleisher J.M., Fleming L.E., Solo-Gabriele H.M., Kish J.K., Sinigalliano C.D., Plano L.R.W., Elmir S.M., Wang J.D., Withum K., Shibata T., Gidley M.L., Abdelzaher A., He G., Ortega C., Zhu X., Wright M., Hollenbeck J. and Backer L.C. (2010) The BEACHES study: health effects and exposures from non-point source microbial contaminants in subtropical recreational marine waters. *International Journal of Epidemiology* 39, 1291–1298.
- Fogarty L.R., Haack S.K., Wolcott M.J. and Whitman R.L. (2003) Abundance and characteristics of the recreational water quality indicator bacteria *Escherichia coli* and enterococci in gull faeces. *Journal of Applied Microbiology* 94, 865–878.

- Francino P.M. (2012) The ecology of bacterial genes and the survival of the new. *International Journal of Evolutionary Biology* Art. ID 394026, 14 pages. <http://dx.doi.org/10.1155/2012/394026>.
- Fujioka R. and Shizumura L.K. (1985) *Clostridium perfringens*, a reliable indicator of stream water quality. *Journal of the Water Pollution Control Federation* 57, 986–992.
- Ge Z., Nevers M.B., Schwab D.J. and Whitman R.L. (2010) Coastal loading and transport of *Escherichia coli* at an embayed beach in Lake Michigan. *Environmental Science & Technology* 44, 6731–6737.
- Gomes D.N.F., Cavalcanti M.A.Q., Fernandes M.J.S., Lima D.M.M. and Passavante J.Z.O. (2008) Filamentous fungi isolated from sand and water of Bairro Novo and Casa Caiada beaches. *Brazilian Journal of Biology* 68, 577–582.
- Gonzales M., Hanlin T.R., Herrera T. and Ulloa M. (2000) Fungi colonizing hair-bats from three coastal beaches of Mexico. *Mycoscience* 41, 259–262.
- Gooch-Moore J., Goodwin K., Doresey C., Ellender R., Mott J., Ornelas M., Sinigalliano C., Vincent B., Whiting D. and Wolfe S. (2011) New USEPA Water Quality criteria 2012. GOMA concerns and recommendations. *Journal of Water and Health* 9, 718–733.
- Halliday E. and Gast R.J. (2011) Bacteria in beach sands: an emerging challenge in protecting coastal water quality and bather health. *Environmental Science & Technology* 45, 370–379.
- Halliday E., McLellan S.L., Amaral-Zettler L.A., Sogin M.L. and Gast R.J. (2014) Comparison of bacterial communities in sands and water at beaches with bacterial water quality violations. *PLoS ONE* 9, e90815. doi: 10.1371/journal.pone.0090815.
- Harwood V.J., Staley C., Badgley B.D., Borges K. and Korajkic A. (2014) Microbial source tracking markers for detection of fecal contamination in environmental waters: relationships between pathogens and human health outcomes. *FEMS Microbiology Review* 38, 1–40.
- Havelaar A.H., van Olphen M. and Drost Y.C. (1993) F-specific RNA-bacteriophages are adequate model organisms for enteric viruses in fresh water. *Applied and Environmental Microbiology* 59, 2956–2962.
- Health Canada (2012) *Guidelines for Canadian Recreational Water Quality*. 3rd edition. Ottawa: Water, Air and Climate Change Bureau, Healthy Environments and Consumer Safety Branch, Health Canada. (Catalogue No H129–15/2012E). http://www.hc-sc.gc.ca/ewh-semt/pubs/water-eau/guide_water-2012-guide_eau/index-eng.php#a9.
- Heaney C.D., Sams E., Dufour A.P., Brenner K.P., Haugland R.A., Chern E., Wing S., Marshall S., Love D.C., Serre M., Noble R. and Wade T.J. (2012) Faecal indicators in sand, sand contact, and risk of enteric illness among beachgoers. *Epidemiology* 23, 95–106.
- Heaney C., Sams E., Wing S., Marshall S., Brenner K., Dufour A.P. and Wade T.J. (2009) Contact with beach sand among beachgoers and risk of illness. *American Journal of Epidemiology* 170, 164–172.
- Heikens E., van Schaik W., Leavis H.L., Bonten M.J. and Willems R.J. (2008) Identification of a novel genomic island specific to hospital-acquired clonal complex 17 *Enterococcus faecium* isolates. *Applied and Environmental Microbiology* 74, 7094–7097.
- Heim S., Del Mar Lleo M., Bonato B., Guzman C.A. and Canepari P. (2002) The viable but nonculturable state and starvation are different stress responses of *Enterococcus faecalis*, as determined by proteome analysis. *Journal of Bacteriology* 184, 6739–6745.
- Hernandez R.J., Hernandez Y., Jimenez N.H., Piggot A.M., Klaus J.S., Feng Z., Reniers A. and Solo-Gabriele H.M. (2014) Effect of full-scale beach renovation on fecal indicator levels in shoreline sand and water. *Water Research* 48, 579–591.
- Imamura G.J., Boehm A.B., Thompson R.S. and Jay J.A. (2011) Wrack promotes the persistence of fecal indicator bacteria in marine sands and seawater. *FEMS Microbiology Ecology* 77, 40–49.
- Ishii S., Hansen D.L., Hicks R.E. and Sadowsky M.J. (2007) Beach sand and sediments are temporal sinks and sources of *Escherichia coli* in Lake Superior. *Environmental Science & Technology* 41, 2203–2209.
- Ishii S.T., Yan T., Shively D.A., Byappanahalli M.N., Whitman R.L. and Sadowsky M.J. (2006) *Cladophora* (Chlorophyta) spp. harbor human bacterial pathogens in nearshore water of Lake Michigan. *Applied and Environmental Microbiology* 72, 4545–4553.
- International Organization for Standardization and the International Electrotechnical Commission (ISO/IEC) (2005) 17025, *General requirements for the competence of testing and calibration laboratories*, 2nd edn, 12 May 2005. ISBN : 0 580 34929 2
- IWA (2011) Declaration adopted by a joint meeting of the International Water Association Health Related Microbiology and Diffuse Pollution held at Rotorua, New Zealand, September. http://www.iwamicrobiology.org/uploadedFiles/Website_IWA_Microbiology/Newsletter/HRWM_Newsletter_February_2012.
- Jiang L., Hu X., Yin D., Zhang H. and Yu Z. (2011) Occurrence, distribution and seasonal variation of antibiotics in the Huangpu River, Shanghai, China. *Chemosphere* 82, 822–828.
- Kaštelan M., Utješinović-Gudelj V., Prpić-Massari L. and Brajac I. (2014) Dermatophyte infections in Primorsko-Goranska County, Croatia: a 21 year survey. *Acta Dermatovenerologica Croatica* 22, 175–179.
- Kay D. (2015) Centre for Research into Environment and Health, Aberystwyth University, UK. <http://cadair.aber.ac.uk/dspace/handle/2160/12683?locale-attribute=cy>. Last Accessed 18 March 2015.
- Kay D., Wyer M.D., Crowther J. and Fewtrell L. (1999) Faecal indicator impacts on recreational waters: budget studies and diffuse source modeling. *Journal of Applied Microbiology Symposium Supplement* 85, 70S–82S.
- Khiyama H.M. and Makemson J.C. (1973) Sand beach bacteria: enumeration and characterization. *Applied and Environmental Microbiology* 26, 293–297.
- Kinzelman J. and McLellan S. (2009) Success of science-based management practices in reducing swimming bans: a case study from Racine, WI, USA. *Aquatic Ecosystems and Health Management* 12, 187–196.
- Kinzelman J., McLellan S., Daniels A., Cashin S., Singh A., Gradus S. and Bagley R. (2004) Non-point source pollution: determination of replication versus persistence of *Escherichia coli* in surface water and sediments with correlation of levels to readily measurable environmental parameters. *Journal of Water and Health* 2, 103–114.
- Kinzelman J.L., Whitman R.L., Byappanahalli M., Jackson E. and Bagley R.C. (2003) Evaluation of beach grooming techniques on *Escherichia coli* density in foreshore sand. *Lake and Reservoir Management* 19, 349–354.
- Larrondo J.V. and Calvo M.A. (1989) Fungal density in the sands of the Mediterranean coast beaches. *Mycopathologia* 108, 185–193.
- Leonard A.F., Zhang L., Balfour A.J., Garside R. and Gaze W.H. (2015) Human recreational exposure to antibiotic resistant bacteria in coastal bathing waters. *Environment International*. doi: 10.1016/j.envint.2015.02.013. [Epub ahead of print]
- Lévesque B., Brousseau P., Simard P., Dewailly E., Meisels M., Ramsay D. and Joly J. (1993) Impact of the ring-billed gull (*Larus delawarensis*) on the microbiological quality of recreational water. *Applied and Environmental Microbiology* 59, 1228–1230.

- Lleó M., Bonato B., Benedetti D. and Canepari P. (2005) Survival of enterococcal species in aquatic environments. *FEMS Microbiology Ecology* 54, 189–196.
- Lozupone C.A. and Knight R. (2007) Global patterns in bacterial diversity. *Proceeding of the National Academy of Sciences USA* 104, 11436–11440.
- Lu J., Hodon R., Hill S., Schoen M., Ashbolt N., Edge T.A. and Santo Domingo J. (2011) Distribution and potential significance of a gull fecal marker in urban coastal and riverine areas of southern Ontario, Canada. *Water Research* 45, 3960–3968.
- Luther K. and Fujioka R. (2004) Usefulness of monitoring tropical streams for male-specific RNA coliphages. *Journal of Water and Health* 2, 171–181.
- Marzol M.V., Yanes A., Romero C., Brito de Azevedo E., Prada S. and Martins A. (2006a) Los riesgos de las lluvias torrenciales en las islas de la Macaronesia (Azores, Madeira, Canarias y Cabo Verde). In Cuadrat Prats J.M., Saz Sánchez M.A., Vicente Serrano S.M., Lanjeri S., De Luis Arrillaga M. and González-Hidalgo J.C. (eds) *Clima, Sociedad y Medio Ambiente*. Publicaciones de la Asociación Española de Climatología, Serie A, pp. 443–452.
- Marzol M.V., Yanes A., Romero C., Brito de Azevedo E., Prada S. and Martins A. (2006b) Caractéristiques des précipitations dans les îles de la Macaronesia (Açores, Madère, Canaries et Cap Vert). *Les Risques Liés au Temps et au climat. XIX Colloque de l'Association Internationale de Climatologie (AIC)*, Épernay, 415–420.
- McMinn B.R., Korajkic A. and Ashbolt N.J. (2014) Evaluation of *Bacteroides fragilis* GB-124 bacteriophages as novel human-associated faecal indicators in the United States. *Letters in Applied Microbiology* 59, 115–121.
- Mead P.S., Slutsker L., Dietz V., McCaig L.F., Bresee J.S., Shapiro C., Griffin P.M. and Tauxe R.V. (1999) Food-related illness and death in the United States. *Emerging Infectious Diseases* 5, 607.
- Migahed F. (2003) Distribution of fungi in the sandy soil of Egyptian beaches. *Mycobiology* 31, 61–67.
- Mortensen K.L., Mellado E., Lass-Florl C., Rodriguez-Tudela J.L., Johansen H.K. and Arendrup M.C. (2010) Environmental study of azole-resistant *Aspergillus fumigatus* and other aspergilli in Austria, Denmark, and Spain. *Antimicrobial Agents and Chemotherapy* 54, 4545–4549.
- Mudryk Z.J., Kosiorek A. and Perlinski P. (2013) *In vitro* antibiotic resistance of Vibrio-like organisms isolated from seawater and sand of marine recreation beach in the southern Baltic Sea. *Hydrobiologia* 702, 141–150.
- NAS U.S. National Academy of Science (1972) Water Quality Criteria 1972, a report of the committee on water quality criteria, 549, National Academy of Sciences, National Technical Advisory Committee, Federal Water Pollution Administration, Department of the Interior, Washington DC.
- Olapade O.A., Depas M.M., Jensen E.T. and McLellan S.L. (2006) Microbial communities and faecal indicator bacteria associated with *Cladophora* mats on beach sites along Lake Michigan shores. *Applied and Environmental Microbiology* 72, 1932–1938.
- Orozco A.S., Higginbotham L.M., Hitchcock C.A., Parkinson T., Falconer D., Ibrahim A.S., Ghannoum M.A. and Filler S.G. (1998) Mechanism of fluconazole resistance in *Candida krusei*. *Antimicrobial Agents and Chemotherapy* 42, 2645–2649.
- Papadakis J.A., Mavridou A., Richardson S.C., Lampiri M. and Marcelou U. (1997) Bather-related microbial and yeast populations in sand and seawater. *Water Research* 31, 799–804.
- Pennak R.W. (1951) Comparative ecology of the interstitial fauna of fresh-water and marine beaches. *Annals of Biology* 27, 449–480.
- Pereira E., Figueira C., Aguiar N., Vasconcelos R., Vasconcelos S., Silva M.G., Brandao J. and Prada S. (2013) Microbiological and mycological beach sand quality in a volcanic environment: Madeira archipelago, Portugal. *Science of the Total Environment* 461–462, 469–479.
- Pfaller M.A., Moet G.J., Messer S.A., Jones R.N. and Castanheira M. (2011) Geographic variations in species distribution and echinocandin and azole antifungal resistance rates among *Candida* bloodstream infection isolates: report from the SENTRY Antimicrobial Surveillance Program (2008 to 2009). *Journal of Clinical Microbiology* 49, 396–399. doi: 10.1128/JCM.01398-10. PMID:21068282.
- Phillips M.C., Feng Z., Vogel L.J., Reniers A.J.H.M., Haus B.K., Enns A.A., Zhang Y., Hernandez D.B. and Solo-Gabriele H.M. (2014) Microbial release from seeded beach sediments during wave conditions. *Marine Pollution Bulletin* 79, 114–122.
- Phillips M.C., Solo-Gabriele H.M., Piggot A.M., Klaus J.S. and Zhang Y. (2011a) Relationships between sand and water quality at recreational beaches. *Water Research* 45, 6763–6769.
- Phillips M.C., Solo-Gabriele H.M., Reniers A.J.H.M., Wang J.D., Kiger R.T. and Abdel-Mottaleb N. (2011b) Pore water transport of enterococci out of beach sediments. *Marine Pollution Bulletin* 62, 2293–2298.
- Piggot A.M., Klaus J.S., Johnson S., Phillips M.C. and Solo-Gabriele H.M. (2012) Relationship between enterococcal levels and sediment biofilms at recreational beaches in South Florida. *Applied and Environmental Microbiology* 78, 5973–5982.
- Pinto K., Hachich E., Sato M., Di Bari M., Coelho M., Matté M., Lamparelli C. and Razzolini M. (2012a) Microbiological quality assessment of sand and water from three selected beaches of South Coast, São Paulo State, Brazil. *Water Science and Technology* 66, 2475–2482.
- Pinto D., Ruivo M., Vandamme P. and Lopes M.F.S. (2012b) Hospital-adapted clonal complex 17 *Enterococcus faecium* found among sand enterococcal isolates. *Journal of Environmental Protection* 3, 74–82.
- Plano L.R.W., Garza A.C., Shibata T., Elmir S.M., Kish J., Sinigalliano C.D., Gidley M.L., Miller G., Withum K., Fleming L.E. and Solo-Gabriele H.M. (2011) Shedding of *Staphylococcus aureus* and methicillin-resistant *Staphylococcus aureus* from adult and pediatric bathers in marine waters. *BMC Microbiology* 11, 1–10.
- Plano L.R.W., Shibata T., Garza A.C., Kish J., Fleisher J., Sinigalliano C.D., Gidley M.L., Withum K., Elmir S.M., Hower S., Jackson C.R., Barrett J.B., Cleary T., Davidson M., Davis J., Mukherjee S., Fleming L.E. and Solo-Gabriele H.M. (2013) Human-associated methicillin-resistant *Staphylococcus aureus* from a subtropical recreational marine beach. *Microbial Ecology* 65, 1039–1051.
- Rees R.J., Pinner W.R., Hajjeh A.R., Brandt E.M. and Reingold L.A. (1998) The epidemiological features of invasive mycotic infections in the San Francisco Bay Area, 1992–1993: results of population-based laboratory active surveillance. *Clinical Infectious Diseases* 27, 1138–1147.
- Roberts M.C., Olusegun O.S. and No D. (2013) Comparison of multi-drug resistant environmental methicillin-resistant *Staphylococcus aureus* isolated from recreational beaches and high touch surfaces in built environments. *Frontiers in Microbiology* 4, 1–8.
- Roll B.M. and Fujioka R.S. (1997) Sources of faecal indicator bacteria in a brackish, tropical stream and their impact on recreational water quality. *Water Science and Technology* 35, 179–186.
- Russell T.L., Sassoubre L.M., Wang D., Masuda S., Chen H., Soetjijto C., Hassaballah A. and Boehm A.B. (2013) A coupled modeling and molecular biology approach to microbial source tracking at Cowell Beach, Santa Cruz, CA, United States. *Environmental Science & Technology* 47, 10231–10239.

- Sabino R., Rodrigues R., Costa I., Carneiro C., Cunha M., Duarte A., Faria N., Ferreira F.C., Gargaté M.J., Júlio C., Martins M.L., Nevers M.B., Oleastro M., Solo-Gabriele H., Veríssimo C., Viegas C., Whitman R.L. and Brandão J. (2014a) Routine screening of harmful microorganisms in beach sands: implications to public health. *Science of the Total Environment* 472, 1062–1069.
- Sabino R., Veríssimo C., Cunha M.A., Wergikoski B., Ferreira F.C., Rodrigues R., Parada H., Falcão L., Rosado L., Pinheiro C., Paixão E. and Brandão J. (2011) Pathogenic fungi: an unacknowledged risk at coastal resorts? New insights on microbiological sand quality in Portugal. *Marine Pollution Bulletin* 62, 1506–1511.
- Sabino R., Veríssimo C., Parada H., Brandão J., Viegas C., Carolino E., Clemons K.V. and Stevens D.A. (2014b) Molecular screening of 246 Portuguese *Aspergillus* isolates among different clinical and environmental sources. *Medical Mycology* 52, 519–529.
- Santiago-Rodriguez T.M., Marcos P., Monteiro S., Urdaneta M., Santos R. and Toranzos G.A. (2013) Evaluation of *Enterococcus*-infecting phages as indices of fecal pollution. *Journal of Water and Health* 11, 51–63.
- Shah A., Abdelzaher A., Phillips M., Hernandez R., Solo-Gabriele H., Kish J., Scorzetti G., Fell J., Diaz M., Scott T., Lukaski J., Harwood V., McQuaig S., Sinigalliano C., Gidley M., Wanless D., Ager A., Lui J., Stewart J., Plano L. and Fleming L. (2011) Indicator microbes correlate with pathogenic bacteria, yeasts and helminthes in sand at a subtropical recreational beach site. *Journal of Applied Microbiology* 110, 1571–1583.
- Shibata T. and Solo-Gabriele H. (2012) Quantitative microbial risk assessment of human illness from exposure to marine beach sand. *Environmental Science & Technology* 46, 2799–2805.
- Shibata T., Solo-Gabriele H.M., Fleming L. and Elmir S. (2004) Monitoring marine recreational water quality using multiple microbial indicators in an urban tropical environment. *Water Research* 38, 3119–3131.
- Sinigalliano C.D., Fleisher J.M., Gidley M.L., Solo-Gabriele H.M., Shibata T., Plano L.R.W., Elmir S.M., Wanless D., Bartkowiak J., Boiteau R., Withum K., Abdelzaher A.M., He G., Ortega C., Zhu X., Wright M.E., Kish J., Hollenbeck J., Scott T., Backer L.C. and Fleming L.E. (2010) Traditional and molecular analyses for fecal indicator bacteria in non-point source subtropical recreational marine waters. *Water Research* 44, 3763–3772.
- Sipsas N.V., Lewis R.E., Tarrand J., Hachem R., Rolston K.V., Raad I.I. and Kontoyiannis D.P. (2009) Candidemia in patients with hematologic malignancies in the era of new antifungal agents (2001–2007): stable incidence but changing epidemiology of a still frequently lethal infection. *Cancer* 115, 4745–4752.
- Snelders E., van der Lee A.L.H., Kuijpers J., Rijs J.M.M.A., Varga J., Samson R.A., Mellado E., Donders A.R.T., Melchers W.J.G. and Verweij P.E. (2008) Emergence of azole resistance in *Aspergillus fumigatus* and spread of a single resistance mechanism. *PLoS Medicine* 5, e219.
- Soller A., Bartrand T., Ashbolt N.J., Ravenscroft J. and Wade T.J. (2010) Estimating the primary aetiological agents in freshwater recreational waters impacted by human sources of faecal contamination. *Water Research* 44, 4473–4477.
- Soller J.A., Schoen M.E., Varghese A., Ichida A.M., Boehm A.B., Eftim S., Ashbolt N.J. and Ravenscroft J.E. (2014) Human health risk implications of multiple sources of faecal indicator bacteria in a recreational waterbody. *Water Research* 66, 254–264.
- Solo-Gabriele H.M., Neumeister S.M., Dubon J.M. and Palmer C.J. (1998) Occurrence of *Cryptosporidium* oocysts and *Giardia* cysts in water supplies of San Pedro Sula, Honduras. *Pan American Journal of Public Health* 4, 398–400.
- Staley C., Gould T.J., Wang P., Phillips J., Cotner J.B. and Sadowsky M.J. (2015) Species sorting and seasonal dynamics primarily shape bacterial communities in the Upper Mississippi River. *Science of the Total Environment* 505, 435–445.
- Stoeckel D.M. and Harwood V.J. (2007) Performance, design, and analysis in microbial source tracking studies. *Applied and Environmental Microbiology* 73, 2405–2415.
- Suzuki S. and Hoa P.T. (2012) Distribution of quinolones, sulfonamides, tetracyclines in aquatic environment and antibiotic resistance in Indochina. *Frontiers in Microbiology* 3, 67.
- The Fungal Research Trust (2011) *How common are fungal diseases?* London: Fungal Research Trust 20th Anniversary Meeting.
- USEPA (1976) *Quality criteria for water*. Washington, DC: U.S. Environmental Protection Agency. PB-263 943.
- USEPA (1986) *Ambient water quality criteria for bacteria – 1986*. Washington, DC: U.S. Environmental Protection Agency, Office of Research and Development, Microbiology and Toxicology Division and Office of Water Regulations and Standards, Criteria and Standards Division. EPA 440/5-84-002.
- USEPA (2005) *Microbial source tracking guide document*. Washington, DC: Office of Research and Development. EPA-600/R-05/064. 131 pp.
- USEPA (2012) *Recreational water criteria*. Washington, DC: US Environmental Protection Agency Office of Water. 820-F-12-058.
- Velonakis E., Dimitriadi D., Papadogiannakis E. and Vatopoulos A. (2014) Present status of effect of microorganisms from sand beach on public health. *Journal of Coastal Life Medicine* 2, 746–756.
- Verweij P.E., Snelders E., Kema G.J.H., Mellado E. and Melchers W.J.G. (2009) Azole resistance in *Aspergillus fumigatus*: a side-effect of environmental fungicide use? *Lancet Infectious Diseases* 9, 789–795.
- Vicente V., Attili-Angelis D., Pie M.R., Queiroz-Telles F., Cruz L.M., Najafzadeh M.J., de Hoog G.S., Zhao J. and Pizzirani-Kleiner A. (2008) Environmental isolation of black yeast-like fungi involved in human infection. *Studies in Mycology* 61, 137–144.
- Vital M., Hammes F. and Egli T. (2008) *Escherichia coli* O157 can grow in natural freshwaters at low carbon concentrations. *Environmental Microbiology* 10, 2387–2396.
- Vogel C., Rogerson A., Schatz S., Laubach H., Tallman A. and Fell J. (2007) Prevalence of yeast in beach sand at three bathing beaches in South Florida. *Water Research* 41, 1915–1920.
- Wang X., Singh P., Gao Z., Zhang X., Johnson I.Z. and Wang G. (2014) Distribution and diversity of planktonic fungi in the West Pacific warm pool. *PLoS ONE* 9. doi:10.1371/journal.pone.0101523. PLoS One. 2014 Jul 3;9(7):e101523. doi: 10.1371/journal.pone.0101523. eCollection 2014.
- Ward C. (2009) *Do seaweeds play a part in the survival and/or growth of faecal indicator organisms which could affect counts on bathing beaches in UK waters?* Millport Marine Station: University of London.
- Weidhaas J.L., Li X. and Harwood V.J. (2014) Microbial Source Tracking (MST) Microarray for Advanced Pathogen Detection and Source Identification in Environmental Systems. American Society for Microbiology General Meeting. 17–20, May. Boston, MA.
- Wellington E.M.H., Boxall A.B.A., Cross P., Feil E.J., William H.G., Hawkey P.M., Johnson-Rollings A.S., Jones D.L., Lee N.M., Otten W., Thomas C.M. and Williams A.P. (2013) The role of the natural environment in the emergence of antibiotic resistance in Gram-negative bacteria. *Lancet Infectious Diseases* 13, 155–165.
- Wheeler J.G., Sethi D., Cowden J.M., Wall P.G., Rodrigues L.C., Tompkins D.S., Hudson M.J. and Roderick P.J. (1999) Study of infectious intestinal disease in England: rates in the community, presenting to general practice, and reported to national surveillance.

- The Infectious Intestinal Disease Study Executive. *British Medical Journal* 318, 1046–1050.
- Whitman R., Harwood V.J., Edge T.A., Nevers M., Byappanahalli M., Vijayavel K., Brandão J., Sadowsky M.J., Alm E.W., Crowe A., Ferguson D., Zhongfu G., Halliday E., Kinzelman G., Kleinheinz G., Przybyla-Kelly K., Staley C., Staley Z. and Solo-Gabriele H.M.** (2014) Microbes in beach sands: integrating environment, ecology and public health. *Reviews in Environmental Science and Biotechnology* 13, 329–368.
- Whitman R.L., Horvath T.G., Goodrich M.L., Nevers M.B., Wolcott M.J. and Haack S.K.** (2001) Characterization of *E. coli* levels at 63rd Street Beach; City of Chicago and the Chicago Park District, Chicago. <http://www.onsetcomp.com/files/63rdStreetBeachReport.pdf>.
- Whitman R.L. and Nevers M.B.** (2003) Foreshore sand as a source of *Escherichia coli* in nearshore water of a Lake Michigan beach. *Applied and Environmental Microbiology* 69, 5555–5562.
- Whitman R.L., Nevers M.B. and Byappanahalli M.N.** (2006a) Examination of the watershed-wide distribution of *Escherichia coli* along Southern Lake Michigan: an integrated approach. *Applied and Environmental Microbiology* 72, 7301–7310.
- Whitman R., Przybyla-Kelly K., Shively D., Nevers M. and Byappanahalli M.** (2009) Hand-mouth transfer and potential for exposure to *E. coli* and F+ coliphage in beach sand, Chicago, Illinois. *Journal of Water and Health* 7, 623–629.
- Whitman R., Shively D., Pawlik H., Nevers M. and Byappanahalli M.** (2003) Occurrence of *Escherichia coli* and enterococci in *Cladophora* (Chlorophyta) in nearshore water and beach sand of Lake Michigan. *Applied and Environmental Microbiology* 69, 4714–4719.
- Whitman R.L., Shively D., Ting W., Tseng C. and Nevers M.** (2006b) Seasonal persistence and population characteristics of *Escherichia coli* and enterococci in deep backshore sand of two freshwater beaches. *Journal of Water Health* 4, 313–320.
- WHO** (1999) *Health-based monitoring of recreational waters: the feasibility of a new approach (the 'Annapolis Protocol')*. Report WHO/SDE/WSH99.1. Geneva: World Health Organization.
- WHO** (2003) *Guidelines for safe recreational water environments*. Volume 1: Coastal and fresh waters. Geneva: World Health Organization.
- WHO** (2009) *Addendum to the WHO Guidelines for safe recreational water environments*, Volume 1, Coastal and fresh waters. WHO/HSE/WSH/10.04. Geneva: World Health Organization.
- WHO** (2014) *Antimicrobial – resistance: global report on surveillance*. Geneva: World Health Organization. http://apps.who.int/iris/bitstream/10665/112642/1/9789241564748_eng.pdf
- Wright M.E., Solo-Gabriele H.M., Elmir S. and Fleming L.E.** (2009) Microbial load from animal faeces at a recreational beach. *Marine Pollution Bulletin* 58, 1649–1656.
- Wright M., Abdelzaher A., Solo-Gabriele H., Elmir S. and Fleming L.** (2011) The inter-tidal zone is the pathway of input of enterococci to a subtropical recreational marine beach. *Water Science and Technology* 63, 542–549.
- Wyer M., Wyn-Jones P., Kay D., Au-Yeung H., Gironés S., López-Pila J., de Roda Husman A., Rutjes S. and Schneider O.** (2012) Relationships between human adenoviruses and faecal indicator organisms in European recreational waters. *Water Research* 46, 4139–4141.
- Yaguchi T., Horie Y., Tanaka R., Matsuzawa T., Ito J. and Nishimura K.** (2007) Molecular phylogenetics of multiple genes on *Aspergillus* section *Fumigati* isolated from clinical specimens in Japan. *Japanese Journal of Medical Mycology* 48, 37–46.
- Yamahara K.M., Sassoubre L.M., Goodwin K.D. and Boehm A.B.** (2012) Occurrence and persistence of bacterial pathogens and indicator organisms in beach sand along the California coast. *Applied and Environmental Microbiology* 78, 1733–1745.
- Yamahara K., Walters S. and Boehm A.** (2009) Growth of enterococci in unaltered, unseeded beach sands subject to tidal wetting. *Applied and Environmental Microbiology* 75, 1517–1524.
- and
- Yates M.V.** (2007) Classical indicators in the 21st century – far and beyond coliform. *Water Environment Research* 79, 279–286.
- Correspondence should be addressed to:**
 H.M. Solo-Gabriele
 Dept. of Civil, Arch., and Environmental Engineering
 P.O. Box 248294 Coral Gables, FL, USA 33124-0630
 email: hmsolo@miami.edu