

Role of Macronutrients and Micronutrients in DNA Damage: Results From a Food Frequency Questionnaire

Carina Ladeira^{1,2,3}, Elisabete Carolino¹, Manuel C Gomes⁴ and Miguel Brito²

¹Environment and Health Research Group, Instituto Politécnico de Lisboa (IPL), Escola Superior de Tecnologia da Saúde de Lisboa (ESTeSL), Lisboa, Portugal. ²Grupo de Investigação em Genética e Metabolismo, Instituto Politécnico de Lisboa (IPL), Escola Superior de Tecnologia da Saúde de Lisboa (ESTeSL), Lisboa, Portugal. ³Centro de Investigação e Estudos em Saúde Pública, Escola Nacional de Saúde Pública (ENSP), Universidade Nova de Lisboa, Lisboa, Portugal. ⁴Faculdade de Ciências da Universidade de Lisboa, Lisboa, Portugal.

Nutrition and Metabolic Insights

1–8

© The Author(s) 2017

Reprints and permissions:

sagepub.co.uk/journalsPermissions.nav

DOI: 10.1177/1178638816684666



ABSTRACT: The links between diet and genomic instability have been under investigation for several decades, and evidence suggests a significant causal or preventive role for various dietary factors. This study investigates the influence of macronutrients (calories, protein, and glucides) and micronutrients, such as vitamins and minerals, as assessed by a food frequency questionnaire, on genotoxicity biomarkers measured by cytokinesis-blocked micronucleus assay and comet assay. The results found significant positive and negative correlations. Micronucleus frequency tends to increase with higher intake of caffeine, calcium, magnesium, zinc, and protein ($P < .05$, Spearman correlation). Calorie and omega-6 intakes are negatively correlated with DNA damage measured by the comet assay. These results are somewhat controversial because some of the correlations found are contrary to dominant views in the literature; however, we suggest that unraveling the association between diet and genetic instability requires a much better understanding of the modulating role of macronutrients and micronutrients.

KEYWORDS: macronutrients, micronutrients, food frequency questionnaire, genetic instability, CBMN assay, comet assay

RECEIVED: June 9, 2016. **ACCEPTED:** November 25, 2016.

PEER REVIEW: Four peer reviewers contributed to the peer review report. Reviewers' reports totaled 1359 words, excluding any confidential comments to the academic editor.

TYPE: Original Research

FUNDING: The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This work was partially funded by Autoridade para as Condições de Trabalho—ACT Project No. 036APJ/09.

DECLARATION OF CONFLICTING INTERESTS: The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

CORRESPONDING AUTHOR: Carina Ladeira, Environment and Health Research Group, Instituto Politécnico de Lisboa (IPL), Escola Superior de Tecnologia da Saúde de Lisboa (ESTeSL), Av. D. João II, Lote 4.69.01, 1990-096 Lisboa, Portugal. Email: carina.ladeira@estesl.ipl.pt

Introduction

Dietary habits are recognized to be an important modifiable environmental factor influencing cancer risk, depending on the foods consumed and the specific type of cancer.^{1–3} Nutrition science consists in the application of molecular biology technology in epidemiologic investigations that, starting from an individual health perspective, upgrades to the population.⁴ Nutritional genomics studies the functional interaction between nutrition (macronutrients and micronutrients) and an individual's genome at the molecular, cellular, and systemic level.^{5,6} Integrating the information regarding nutritional intake, effect biomarkers, susceptibility biomarkers (gene polymorphisms and expression), diseases, and phenotypes may lead to the development of nutritional biomarkers.⁷ Understanding the influence of nutrition on metabolic pathways and homeostatic control and the mechanisms of its (de) regulation could lead to evidence-based dietary intervention strategies aimed at restoring health and fitness for preventing diet-related diseases.⁸

Dietary patterns involve complex interactions of food and nutrients, being important to study not only each individual nutrient but also the diet as a whole. An important example is the high consumption of vegetables, legumes, fruits, nuts, minimally processed cereals, monounsaturated lipids, moderately

high consumption of fish, low consumption of dairy and meat products, and regular but moderate intake of alcohol of the well-characterized Mediterranean diet, which is linked with cancer prevention.^{9,10}

Important epidemiologic studies postulate the role of carcinogens present in food and how its exposure is linked to specific cancers and how incidence varies among countries.¹ The classical evidence of the relationship between diet and cancer is shown by the change in the incidence of certain types of cancers occurring in populations that migrate to a different geographic area and culture,¹¹ suggesting that international differences in cancer incidence can be attributed primarily to environmental or lifestyle factors rather than genetic ones.^{1,11}

There are several known mechanisms by which diet can influence cancer development: food carcinogens that act on DNA provoking its damage, blockage, or induction of enzymes involved in the activation or deactivation of carcinogenic substances, inadequate intake-promoting alterations in DNA synthesis, repair, or methylation, thereby influencing mutation rate and/or gene expression.¹² The impact of energy balance and growth rates—hormone levels and growth factors—should also be taken into account because it influences cell division, cell cycle, DNA repair, and replication rates.¹²



Creative Commons Non Commercial CC-BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 3.0 License (<http://www.creativecommons.org/licenses/by-nc/3.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

Tools to assess markers of general DNA damage include cytokinesis-blocked micronucleus cytome (CBMN) assay, a comprehensive system for measuring DNA damage which measures end points such as micronuclei (MN), a biomarker of chromosome breakage and/or whole chromosome loss; nucleoplasmic bridges (NPB), a biomarker of DNA misrepair and/or telomere end-fusions; and nuclear buds (NBUD), a biomarker of elimination of amplified DNA and/or DNA repair complexes.^{13,14}

MN are small, extranuclear bodies, originate in dividing cells from acentric chromosome/chromatid fragments or whole chromosome/chromatid that lag behind in anaphase, and are not included in the daughter nuclei in telophase.^{15–19}

NPB occur when centromeres of dicentric chromosomes are pulled to opposite poles of the cell at anaphase. In the absence of breakage of the anaphase bridge, the nuclear membrane eventually surrounds the daughter nuclei and the anaphase bridge, and a nucleoplasmic bridge is formed.²⁰ NBUD have been observed in cultures grown under strong selective conditions that induce gene amplification, as well as under moderate folic acid deficiency.^{14,17} Gene amplification plays a crucial role in the malignant transformation of human cells as it mediates the activation of oncogenes or the acquisition of drug resistance.²¹

Another useful tool is the comet assay, a simple and sensitive method for detecting DNA-strand breaks that originated from many sources, such as through oxidative DNA damage.^{22,23}

The comet assay has become one of the standard methods for assessing DNA damage, with a wide range of applications, namely, in genotoxicity testing, human biomonitoring and molecular epidemiology, ecogenotoxicology, as well as fundamental research in DNA damage and repair^{24,25}; studying the mechanisms of action of genotoxic chemicals; investigating oxidative damage as a factor in disease; monitoring oxidative stress in animals or human subjects resulting from exercise, diet, or exposure to environmental agents; studying the effects of dietary antioxidants; and monitoring environmental pollution by studying sentinel organisms.^{26,27}

Although there is a general recognition that diet influences cancer risk, information regarding precise dietary factors that determine human cancer is an ongoing debate.^{9,11,28,29} Diet is usually a complex of both nutritive (macronutrients and micronutrients) and non-nutritive food constituents; thus, the search for specific factors is bound to involve a long gathering of epidemiologic information regarding food habits. Epidemiologic studies based on food frequency questionnaires (FFQs) are tools aimed at such goals and can be very important to develop a hypothesis regarding diet and cancer.

The aim of this study was to investigate the potential influence of macronutrients (calories, protein, and glucides) and micronutrients (vitamins and minerals) by a FFQ on genotoxicity biomarkers measured by the CBMN assay and the comet assay.

Methods

Subjects and personal questionnaire

The sample studied consisted of 44 healthy subjects from the Lisbon and Tagus Valley Region, mainly whites, in Portugal with administrative job descriptions that assigned to participate in the study.

The study participants filled in a questionnaire on individual characteristics and working practices. These included questions regarding age, gender, tobacco and alcohol consumption habits, medication, hereditary diseases, cancer history, exposure to chemical agents, and characterization of professional activity. The questionnaire was validated by application of a pretest in a group of subjects, which was used as a pilot sample at the beginning of the study.

Food frequency questionnaire

Dietary intake was assessed using a self-administered FFQ.^{30,31} The FFQ included type and quantity of food intake, namely, some food items, which allowed for the quantification of different macronutrients and micronutrients. The FFQ is a 3-page booklet with a list of 92 common food and beverage items and questions relating to food preparation and dietary habits. Participants were required to indicate how often each food and beverage was usually consumed per month, week, or day. Average daily consumption was based on the participants' reports on how often a specified serving size of each food or beverage item was consumed. This information, along with the nutrient composition of the food item/unit weight taken from 92 selected items, allowed participants' daily micronutrient and macronutrient intake to be calculated using the FREQUAN dietary analysis program.³²

Laboratory Proceedings

Cytokinesis-blocked micronucleus assay

Evaluation of genotoxic effects was conducted by applying the CBMN assay in peripheral blood lymphocytes as described in Ladeira et al.^{33,34} All blood samples were collected from subjects by venipuncture with heparin between 10 and 12 AM, and coded and analyzed under blind conditions. Briefly, lymphocytes were isolated using Ficoll-Paque gradient and placed in RPMI 1640 culture medium with L-glutamine and red phenol added with 10% inactivated fetal calf serum, 50 µg/mL streptomycin + 50 U/mL penicillin, and 10 µg/mL phytohemagglutinin. Duplicate cultures from each subject were incubated at 37°C in a humidified 5% CO₂ incubator for 44 hours, and cytochalasin B 6 µg/mL was added to the cultures to prevent cytokinesis. After 28 hours of incubation, cells were spun onto microscope slides using a cytocentrifuge. Smears were air-dried and double-stained with May-Grünwald-Giemsa and mounted with Entellan. The criteria for scoring the nuclear abnormalities in lymphocytes

were the ones established and validated by the Human MicroNucleus International Collaborative Project available in <http://www.humn.org> and in Fenech et al.³⁵

Comet assay

Isolated lymphocytes were cryopreserved following the protocols of Duthie et al³⁶ and Singh and Lai.³⁷ Briefly, isolated lymphocytes suspended in RPMI medium with L-glutamine were centrifuged (600g, 10 minutes) and resuspended in freezing mix (90% v/v heat-inactivated fetal calf serum and 10% v/v dimethyl sulfoxide [DMSO]), frozen at $-1^{\circ}\text{C}/\text{min}$ in polystyrene, and stored at -80°C .

For analysis of DNA damage and oxidative damage by alkaline comet assay, a modification of the originally described technique by Singh et al³⁸ was used to measure the basal level of DNA oxidation in lymphocytes.³⁹ Briefly, cell suspension (2.0×10^4 cells/mL) was mixed with 140 μL of 1% low melting point agarose (LM Pronadisa) in a microcentrifuge tube and added to a slide previously precoated with 1% agarose (SeaKem), 2 gels per slide, and covered with a cover slip ($22\text{ mm} \times 22\text{ mm} \times 1.0\text{ mm}$) and allowed to set on a cold plate. The cover slips were removed and the slides immersed in lysis solution (2.5M NaCl, 100mM Na_2EDTA , 10mM TRIS, 1% Triton X-100 [pH 10]) for 60 minutes. Following lysis, the slides were immersed in 2 changes of Buffer F (40mM HEPES, 0.1M KCl, 0.5mM Na_2EDTA and 0.2mg/L BSA [pH 8.0]) for 5 minutes, each time at 4°C . FPG (kindly donated by Professor Andrew Collins [Department of Nutrition, University of Oslo, Norway]) was added to the gel previously diluted in Buffer F. Incubation of the slides with FPG and Buffer F was performed in a humid chamber at 37°C for 30 minutes. The reaction was stopped by placing them at 4°C . The cover slips were removed and all the slides—lysis, Buffer F, and FPG treatment—were placed on an electrophoresis platform, covered with electrophoresis buffer (1mM Na_2EDTA , 0.3M NaOH [pH 13]), and DNA was allowed to unwind for 20 minutes before electrophoresis at 1.14V/cm, 300mA for a further 20 minutes. DNA unwinding and electrophoresis were performed in a cold unit at 4°C . The slides were transferred to a Coplin jar and immersed in PBS and then in distilled water, both for 10 minutes at 4°C . After that, the slides were dehydrated in increasing ethanol concentrations (70%, 96%, and 100%), 5 minutes each. The slides dried at room temperature were stained with DAPI (1 $\mu\text{g}/\text{mL}$) and then visualized and scored by one single observer using Zeiss AxioScope.A1 fluorescence microscope according to the criteria of scoring comets from each gel described by Collins,⁴⁰ and the Comet Assay IV software from Perceptive Instruments (Bury St Edmunds, UK) was used.

Statistical analysis

Variables were compared with the normal distribution using the Shapiro-Wilk test. The association between the macronutrients and micronutrients and genotoxicity biomarkers was investigated by Spearman correlation coefficient. Statistical

analyses were performed using the SPSS package for Windows, version 21.0.

Results

Individual characteristics such as gender distribution, age, tobacco habits, and alcohol consumption were analyzed for the sample of 44 individuals. In the sample, 75% were women ($n=33$) and 25% men ($n=11$), with a mean age of 39.26 ± 1.46 years. Concerning tobacco habits, nonsmokers comprised 77.3% of the sample ($n=34$), whereas 22.7% were smokers ($n=10$).

Twenty-three items were selected from the FFQ based on the possible influence on genomic stability. Therefore, the following nutritional items were selected from the FFQ for analysis: total calories; protein; glucides; fat; niacin; calcium; copper; magnesium; manganese; zinc; iron; selenium; vitamins B12, C, D, E, and K; retinol; folate; omega-3 and omega-6; fibers; and caffeine. A statistical description of the amounts consumed daily by item in the sample, as well as their reference intake values, is shown in Table 1. All the studied items presented values between the minimum and maximum reference values, meaning no deficiencies or excesses existed.

To investigate the possible correlations between genotoxicity biomarkers and nutritional items, bivariate Spearman correlations were computed. There were 2 significant positive correlations, between genotoxicity biomarkers (MN and NBUD) and protein, and a negative correlation between calories and % DNA (Table 2).

Regarding significant correlations with micronutrients, MN were positively correlated with calcium, magnesium, zinc, and caffeine and niacin with NBUD, and none of the vitamins assessed by the FFQ exhibited a statistically significant correlation with the genotoxicity biomarkers studied.

Other positive correlations were found, however, without statistical significance, namely, niacin, copper, and caffeine with all the genotoxicity biomarkers measured by CBMN and with oxidative DNA damage. Vitamin C was also positively correlated with the CBMN genotoxicity biomarkers under study (MN, NPB, and NBUD) but not with the biomarkers measured by the comet assay (DNA damage and oxidative DNA damage).

Two statistically significant negative correlations were found between omega-6 and calorie intake and DNA damage measured by the comet assay.

Several other negative correlations were found without reaching statistical significance, namely, between vitamins B12 and D and all the genotoxicity biomarkers studied; folate was correlated with MN, NPB, NBUD, and DNA damage, and omega-3 and omega-6 were negatively correlated with NPB, NBUD, DNA damage, and oxidative DNA damage.

Discussion

Determining the intake levels of macronutrients and micronutrients required to maintain genome stability is an essential step in the definition of optimal diets for the prevention of

Table 1. Dietary parameters (macronutrients, micronutrients, and others) collected by FFQ (mean intake per day \pm SD) and respective dietary reference intakes.

DIETARY PARAMETERS		MEAN \pm SD (RANGE)	DIETARY REFERENCE INTAKES (FOOD AND NUTRITION BOARD, INSTITUTE OF MEDICINE, NATIONAL ACADEMIES)
Macronutrients	Calories	2527.40 \pm 123.07	Variable by age and gender (kcal)
	Protein	115.36 \pm 37.02 (57.89-211.31)	Variable by age and gender, g/d
	Glucides	315.00 \pm 125.56 (127.54-621.198)	Variable by age and gender, g/d
	Fat	95.75 \pm 28.49 (41.45-150.72)	Variable by age and gender
	Niacin	27.75 \pm 9.27 (12.53-50.17)	35 mg/d
	Calcium	1227.97 \pm 582.02 (538.96-3584.74)	1000 mg/d
	Copper	2.10 \pm 0.84 (0.98-4.25)	700 μ g/d
	Magnesium	417.03 \pm 159.16 (161.72-805.18)	Female: 265 mg/d Male: 350 mg/d
	Manganese	4.82 \pm 2.30 (1.24-10.79)	Female: 1.8 mg/d Male: 2.3 mg/d
	Zinc	14.89 \pm 4.69 (8.02-24.96)	Female: 6.8 mg/d Male: 9.4 mg/d
	Iron	18.82 \pm 1.07 (1.13-6.66)	Female: 8.1 mg/d Male: 6 mg/d
	Selenium	138.67 \pm 8.58 (1.13-6.66)	45 μ g/d
	Vitamin B12	12.31 \pm 0.78 (1.13-6.66)	2.0 μ g/d
	Vitamin C	163.65 \pm 97.61 (41.37-440.86)	Female: 60 mg/d Male: 75 mg/d
	Vitamin D	4.67 \pm 0.35 (1.13-6.66)	10 μ g/d
	Vitamin E	11.80 \pm 0.67 (1.13-6.66)	12 μ g/d
	Vitamin K	18.40 \pm 15.13 (2.46-81.63)	Female: 35 μ g/d Male: 45 μ g/d
	Retinol	776.51 \pm 70.10 (1.13-6.66)	Female: 500 μ g/d Male: 625 μ g/d
	Folate	401.21 \pm 26.21 (1.13-6.66)	320 μ g/d
	Omega-3	1.64 \pm 0.59 (0.67-3.63)	Variable by age and gender
	Omega-6	12.30 \pm 5.38 (5.44-27.10)	Variable by age and gender
Others	Fibers	30.01 \pm 14.05 (9.32-61.28)	Female: 25 g/d Male: 38 g/d
	Caffeine	83.22 \pm 59.16 (4.13-350.25)	200-400 mg/d

Abbreviation: FFQ, Food Frequency Questionnaire.

cancer and other diseases caused by genome damage.⁴¹ We have conducted an FFQ inquiry of 44 subjects, followed by a detailed assessment of DNA damage markers (MN, NPB, NBUD, and DNA tail by the comet assay), which found 7 significant positive correlations and 2 significant negative ones.

We have found correlations between calorie and protein intake and genotoxicity biomarkers. The average amount of calories ingested daily was negatively correlated with % DNA in tail, an indicator of DNA damage. This was somewhat unexpected, as calorie restriction reduces metabolic rate and oxidative stress, meaning lower calorie ingestion is expected to decrease DNA damage,^{42,43} as well as damage to other molecules, such as protein and lipids.⁴⁴ Regardless of the source and nature of DNA damage, DNA repair is better preserved and/or enhanced when caloric consumption decreases.⁴² The possible

mechanisms associating calorie restriction with cancer prevention evolve around the regulation of cellular proliferation and apoptosis (decrease in DNA replication) and reduction in metabolic rate, oxidative damage, and inflammation mediators (reduction in reactive oxygen species [ROS] and consequent reduction in DNA damage).⁴⁵ Therefore, it is possible that because of the self-reported nature of the FFQ, some dietary items become subjectively overrated, in a way that cannot be easily corrected.

A positive correlation was found between protein intake and micronucleus, meaning people with higher protein consumption would on average have higher micronucleus frequency. In general, high-protein diets show a protective effect that could be due to an increased plasma concentration of sulfhydryl amino acids, such as cysteine and methionine, which have

Table 2. Spearman correlations between genotoxicity biomarkers (MN, NPB, NBUD, and % DNA in tail and oxidative DNA damage—FPG) and nutrients in the studied sample.

DIETARY PARAMETERS	MN	NPB	NBUD	% DNA TAIL	FPG
Calories	−0.117	−0.037	−0.005	−0.313*	0.008
Protein	0.375*	−0.059	0.308*	−0.112	−0.130
Glucides	0.057	0.074	0.308*	−0.081	−0.114
Fat	0.259	−0.087	0.098	−0.228	−0.097
Fibers	0.251	0.087	0.343*	−0.106	−0.112
Niacin	0.210	0.104	0.300*	−0.161	0.002
Calcium	0.320*	−0.093	0.137	0.071	0.014
Copper	0.168	0.014	0.157	−0.061	0.018
Magnesium	0.337*	0.031	0.220	−0.139	−0.076
Manganese	0.218	0.178	0.164	−0.139	−0.002
Zinc	0.355*	−0.031	0.219	−0.031	0.059
Iron	−0.167	−0.042	0.019	−0.234	0.143
Selenium	−0.240	−0.149	0.109	−0.250	0.028
Vitamin B12	−0.219	−0.087	−0.093	−0.218	−0.112
Vitamin C	0.250	0.054	0.181	−0.224	−0.170
Vitamin D	−0.191	−0.116	−0.161	−0.106	−0.015
Vitamin E	0.004	−0.048	−0.216	−0.096	0.145
Vitamin K	0.172	−0.037	0.085	0.168	0.163
Retinol	−0.243	0.070	−0.073	−0.086	0.170
Folate	−0.047	−0.144	−0.086	−0.024	0.235
Omega-3	0.143	−0.082	0.215	−0.208	−0.270
Omega-6	0.256	−0.087	0.107	−0.344*	−0.040
Caffeine	0.321*	0.155	0.234	−0.082	0.068

Abbreviations: MN, micronuclei; NPB, nucleoplasmic bridges; NBUD, nuclear buds.

*Significant correlations at $P < .05$.

considerable antioxidant activity, or via increased glutamine, which increases tissue antioxidant glutathione.^{46,47}

Many studies associate increased cancer risk with diets rich in starches, such as sugar and sucrose.¹² In contrast, a diet rich in dietary fibers has been associated with low risk of cancer because fibers can have the potential to dilute carcinogens, speed up bulk transition, reduce time for carcinogen absorption, and also serve as a substrate to generate short-chain fatty acids used by colonic epithelial cells.^{48,49} In this sense, a diet with a high proportion of simple glucides that is high in dietary fibers might increase the risk of cancer development. The study of the relation between fiber and glucide consumption with biomarkers detected statistically significant correlations, in a positive sense with low intensity, between fiber consumption and NBUD ($r=.343$, $P=.023$) and glucide consumption and NBUD ($r=.308$, $P=.042$). Regarding the interaction between fiber and glucide consumption, a positive statistically significant correlation, with low intensity, with NBUD ($r=.315$, $P=.037$) was detected. These results indicate that higher intakes of fibers and glucides can lead to higher NBUD values.

We have also examined the correlation between fiber/glucide consumption and genotoxicity biomarkers and found a

statistically significant positive correlation between them ($r=.871$, $P<0.0001$).

Niacin, or nicotinic acid, is one of the few vitamins that has an intimate role in DNA synthesis, DNA repair, and cell death.^{50–52} Niacin is required as a substrate for poly(ADP-ribose) polymerase (PARP), which is involved in cleavage and rejoining of DNA and in telomere length maintenance. The consequence of its deficiency is increased DNA oxidation, DNA breaks, and an elevated chromosome damage rate.^{51–53} According to the Expert Group on Vitamins and Minerals (2003), there are no genotoxicity and carcinogenicity data available for nicotinic acid, and the impact of niacin on human carcinogenesis is therefore confounded by the effect of other micronutrients.⁵³ Our results show a positive correlation with NBUD. As already mentioned, NBUD are biomarkers of gene amplification, and they play a crucial role in the malignant transformation of human cells as they mediate the activation of oncogenes or the acquisition of drug resistance.²¹

Fenech and Bonassi⁵⁴ showed a decrease in micronucleus frequency related to niacin intake using the CBMN assay, and no results for NBUD were reported.

Calcium has not been considered of etiological importance in cancer for many years. However, several studies have suggested that dietary calcium may be associated with a reduced risk of colon cancer.⁵⁵ Calcium may bind with fats in the intestine, making fats less accessible to metabolism by bile salts and the subsequent formation of carcinogenic breakdown products.⁵⁶ Fenech and Bonassi⁵⁴ showed a decrease in MN frequency related to calcium intake. Our results are not consistent with most epidemiologic literature as we have found a positive correlation between calcium and MN, but we corroborate Smith et al⁵⁶ who also found that high calcium intake was associated with higher levels of MN.

Magnesium by itself is not genotoxic, being highly required to maintain genomic stability. Adding to its stabilizing effect on DNA and chromatin structure, magnesium is an essential cofactor in almost all enzymatic systems involved in DNA processing. As an essential cofactor in nucleotide excision repair, base excision repair, and mismatch repair, magnesium is required for the removal of DNA damage. Its relation with tumor formation is more complex; magnesium appears to be protective at early stages but promotes the growth of existing tumors at latter stages.^{57,58} Any decrease in magnesium below physiological levels should trigger Mg-deficiency-related diseases, such as cardiovascular diseases, accelerated aging, and cell cycle control apoptosis and carcinogenesis.⁵⁸ Our results showed a positive correlation between magnesium and micronucleus frequency. Fenech et al⁴⁶ showed that a mitogen response significantly and positively correlated with plasma zinc, magnesium, selenium, and folate, thus corroborating our results insofar as magnesium is concerned.

Zinc is an important element for numerous proteins, playing a pivotal role in essential cell functions such as cell proliferation and apoptosis, defense against free radicals, and DNA damage repair.^{28,50,51,59-61} Alterations in zinc status would affect DNA integrity by altering oxidative stress, antioxidant defenses, and DNA repair functions.^{60,61} Our study found a positive correlation between zinc and MN, as higher zinc intake was associated with increased MN. This result is contrary to most of the literature; namely, Sharif et al⁶² showed that supplementation in an elderly population was beneficial in reducing the micronucleus frequency and DNA damage. Changes in dietary zinc intake affect DNA single-strand breaks, appearing to be a critical factor for maintaining DNA integrity in humans.⁶³ Antioxidants, such as vitamins A, D, and E, are known to be reducing agents, and these molecules are capable of slowing or preventing the oxidation of other molecules being associated with reduced risk of several chronic diseases, particularly some cancers and heart diseases.⁶⁴⁻⁶⁷ They are at the end of oxidative chain reactions, removing free radicals and preventing the oxidation of unsaturated fats, and are clearly documented antigenotoxic and antimutagenic potential antioxidants.⁶⁷ Epidemiologic evidence indicates that intake of foods that are naturally rich in vitamin C is

associated with reduced risk of cardiovascular and neurodegenerative diseases and various cancers, but the extent to which vitamin C contributes to this effect remains unclear.⁵¹ Folate plays a key role in a number of processes related to DNA integrity, such as DNA synthesis and methylation. In vitro studies have shown that folic acid deficiency causes a dose-dependent increase in uracil incorporation into human lymphocyte DNA. Folate administration reduces DNA uracil incorporation and the occurrence of chromosome breaks in human cells.^{65,68,69} A low folate concentration has been implicated as a potential promoter of carcinogenesis, for example, in colorectal cancer, lung, breast, pancreatic, gastric, esophageal, and prostate malignancies.^{3,28,50,65,70,71}

The results achieved by the FFQ regarding these items were not significant despite the majority of negative correlations, meaning an increase in the intake of vitamins would decrease genotoxic damage; however, statistical significance was not reached.

Omega-6 polyunsaturated fatty acids (PUFAs) are considered to increase lipid peroxidation because of lipid peroxidation products, whereas omega-3 PUFAs exert a chemopreventive effect.⁷² In our study, a significant negative correlation was found between omega-6 and % DNA in tail, meaning higher intake of omega-6 will decrease DNA damage. This result is contrary to most of the literature, given omega-6 fatty acid's reputation for promoting cancer, as shown by the study of Bishop et al⁷³ on DNA damage and dietary fatty acids, where whole blood omega-6 PUFAs were positively correlated with DNA damage, also measured by the comet assay. Along with omega-3, omega-6 plays a crucial role in brain function and normal growth and development. Besides some omega-6 fatty acids promoting inflammation (linoleic acid and arachidonic acid), γ -linolenic acid actually reduces inflammation and even protects DNA.⁷³

Epidemiologic studies have suggested that people who consume a diet high in omega-3 fatty acids may experience a lower prevalence of cancer, and many small trials have attempted to assess the effects of omega-3 fatty acid in the diet, either as omega-3 fatty acid-rich foods or as dietary supplements. A systematic review made by MacLean et al⁷⁴ regarding the effects of omega-3 fatty acids on cancer risk, which compiled a large body of literature spanning numerous cohorts from many countries and with different demographic characteristics, did not provide evidence of a significant association between omega-3 fatty acids and cancer incidence. Our study did not lead to any statistically significant correlation associating omega-3 and biomarkers.

We have found a positive correlation between caffeine consumption and micronucleus, in line with Kiefer and Wiebel⁷⁵ who showed that caffeine increases the number of micronuclei in test cell lines. Also, Smith et al⁵⁶ showed a statistically significant association between MN frequency and coffee consumption. This study also reported that intake of decaffeinated

coffee was not associated with a similar increase in MN. There is no unequivocal epidemiologic evidence indicating that caffeine is carcinogenic to humans,⁷⁵ although it has been a suspected carcinogen, primarily for cancers of the pancreas, bladder, kidney, and ovary.⁵⁶ Caffeine is involved in the inhibition of p53, homologous recombination pathway, namely, Rad-51,⁷⁶ inhibition of DNA synthesis, delays in cell proliferation, reverse cell cycle checkpoint function, enhanced genotoxicity after DNA damage by ultraviolet radiation, and others.⁷⁷

Conclusions

Nowadays, the link between environment, where diet is included, and cancer is gaining even more evidence, indicative of a significant causal or preventive role for various dietary factors.⁶⁰ Therefore, our study deemed to investigate the influence that macronutrients and micronutrients could have on DNA damage, as a known precursor of mutations that can lead to disease. The use of CBMN technique and comet assay to study human nutrition and cancer⁷⁸ is an accepted method, and positive and negative correlations were found.

In general, investigations support the hypothesis that dietary antioxidants may protect against cancer as a moderate effect of long-term antioxidant supplementation on oxidative DNA damage, but significant associations are difficult to find for many reasons. On one hand, studies are usually based on relatively small samples of healthy subjects; on the other hand, synergistic effects involving more than one antioxidant, not seen for each one, should be taken into account,⁷⁸ but that requires much larger samples. To unravel the association between diet and cancer, the modulating role of macronutrients and micronutrients needs to be more clearly understood. It is possible that other technical tools for measuring DNA damage might allow for capturing more accurate associations involving the antioxidants examined. It is also possible that the range of antioxidant concentrations and/or oxidative DNA damage in this study was not wide enough to detect associations or it can be concluded that associations simply do not exist. Finally, our study was based on a self-administered FFQ, allowing for the subjectivity to which questionnaires are usually prone, and we cannot preclude the possibility that this has influenced the results.

Acknowledgements

The authors acknowledge Susana Viegas who was the coordinator of the initial project.

Author Contributions

CL, MCG, and MB conceived the idea and designed the study. CL performed the cytokinesis-blocked micronucleus cytome and comet assays for the assessment of genotoxicity biomarkers and applied the food frequency questionnaires. EC performed the statistical data analysis. All authors have read and approved the final manuscript.

REFERENCES

1. Strickland P, Groopman J. Biomarkers for assessing environmental exposure to carcinogens in the diet. *Am J Clin Nutr*. 1995;61:710S–720S.
2. Davis C, Milner J. Biomarkers for diet and cancer prevention research: potentials and challenges. *Acta Pharmacol Sin*. 2007;28:1262–1273.
3. Sutandyo N. Nutritional carcinogenesis. *Acta Med Indones*. 2010;42:36–42.
4. Go V, Butrum R, Wong D. Diet, nutrition, and cancer prevention: the postgenomic era. *J Nutr*. 2003;133:3830S–3836S.
5. Davis C, Milner J. Frontiers in nutrigenomics, proteomics, metabolomics and cancer prevention. *Mutat Res*. 2004;551:51–64.
6. Elliot R, Ong TJ. Nutritional genomics. *Br Med J*. 2002;324:1438–1442.
7. Ordovas F, Corella D. Nutritional genomics. *Annu Rev Genomics Hum Genet*. 2004;5:71–118.
8. Afman L, Müller M. Nutrigenomics: from molecular nutrition to prevention of disease. *J Am Diet Assoc*. 2006;106:569–576.
9. Couto E, Boffetta P, Lagiou P, et al. Mediterranean dietary pattern and cancer risk in the EPIC cohort. *Br J Cancer*. 2011;104:1493–1499.
10. Couto E, Sandin S, Löf M, Ursin G, Adami H-O. Mediterranean dietary pattern and risk of breast cancer. *PLoS ONE*. 2013;8:e55374.
11. Anand P, Kunnumakara A, Sundaram C, et al. Cancer is a preventable disease that requires major lifestyle changes. *Pharm Res*. 2008;25:2097–2116.
12. Willett W, Giovannucci E. Epidemiology of diet and cancer risk. In: Shils M, Shike M, Ross A, Caballer B, Cousins R, eds. *Modern Nutrition in Health and Disease*. 10th ed. Philadelphia, PA: Lippincott Williams & Wilkins; 2006;1267–1279.
13. Fenech M. Cytokinesis-block micronucleus assay evolves into a “cytome” assay of chromosomal instability, mitotic dysfunction and cell death. *Mutat Res*. 2006;600:58–66.
14. Fenech M. Cytokinesis-block micronucleus cytome assay. *Nat Protoc*. 2007;2:1084–1104.
15. Fenech M. The advantages and disadvantages of the cytokinesis-block micronucleus method. *Mutat Res*. 1997;392:11–18.
16. Fenech M. Important variables that influence base-line micronucleus frequency in cytokinesis-blocked lymphocytes—a biomarker for DNA damage in human populations. *Mutat Res*. 1998;404:155–165.
17. Fenech M. The *in vitro* micronucleus technique. *Mutat Res*. 2000;455:81–95.
18. Fenech M, Crott JW. Micronuclei, nucleoplasmic bridges and nuclear buds induced in folic acid deficient human lymphocytes—evidence for breakage-fusion-bridge cycles in the cytokinesis-block micronucleus assay. *Mutat Res*. 2002;504:131–136.
19. Mateuca R, Lombaert N, Aka PV, Decordier I, Kirsch-Volders M. Chromosomal changes: induction, detection methods and applicability in human biomonitoring. *Biochimie*. 2006;88:1515–1531.
20. Fenech M, Kirsch-Volders M, Natarajan AT, et al. Molecular mechanisms of micronucleus, nucleoplasmic bridge and nuclear bud formation in mammalian and human cells. *Mutagenesis*. 2011;26:125–132.
21. Utani K, Kawamoto J, Shimizu N. Micronuclei bearing acentric extrachromosomal chromatin are transcriptionally competent and may perturb the cancer cell phenotype. *Mol Cancer Res*. 2007;5:695–704.
22. Collins A, Dusinska M, Franklin M, et al. Comet assay in human biomonitoring studies: reliability, validation, and applications. *Environ Mol Mutagen*. 1997;30:139–146.
23. Moller P, Knudsen LE, Loft S, Wallin H. The comet assay as a rapid test in biomonitoring occupational exposure to DNA-damaging agents and effect of confounding factors. *Cancer Epidemiol Biomarkers Prev*. 2000;9:1005–1015.
24. Collins AR. The comet assay for DNA damage and repair: principles, applications, and limitations. *Mol Biotechnol*. 2004;26:249–261.
25. Collins AR. Investigating oxidative DNA damage and its repair using the comet assay. *Mutat Res*. 2009;681:24–32.
26. Dusinska M, Collins AR. The comet assay in human biomonitoring: gene–environment interactions. *Mutagenesis*. 2008;23:191–205.
27. Azqueta A, Shaposhnikov S, Collins A. Detection of oxidised DNA using DNA repair enzymes. In: Dhawan A, Hartmann A, Plewa MJ, et al, eds. *The Comet Assay in Toxicology*. London, England: Royal Society of Chemistry (www.rsc.org); 2009;58–63.
28. Ames B. DNA damage from micronutrient deficiencies is likely to be a major cause of cancer. *Mutat Res*. 2001;475:7–20.
29. Key T, Schatzkin A, Willett W, Allen N, Spencer E, Travis R. Diet, nutrition and the prevention of cancer. *Public Health Nutr*. 2004;7:187–200.
30. Lopes C. Reproducibility and validation of a Food Frequency Questionnaire. In: *Diet and Myocardial Infarction: A Community-Based Case-Control Study. A Population-Based Case Control Study* [PhD thesis]. Porto, Portugal: University of Porto; 2000;79–115 (in Portuguese).
31. Lopes C, Aro A, Azevedo A, Ramos E, Barros H. Intake and adipose tissue composition of fatty acids and risk of myocardial infarction in a male Portuguese community sample. *J Am Diet Assoc*. 2007;107:276–286.

32. Baghurst K, Record S. A computerised dietary analysis system for use with diet diaries or food frequency questionnaires. *Aust NZ J Publ Heal.* 1984;1:11–18.
33. Ladeira C, Viegas S, Carolino E, Gomes MC, Brito M. The influence of genetic polymorphisms in XRCC3 and ADH5 genes on the frequency of genotoxicity biomarkers in workers exposed to formaldehyde. *Environ Mol Mutagen.* 2013;54:213–221.
34. Ladeira C, Viegas S, Carolino E, Prista J, Gomes MC, Brito M. Genotoxicity biomarkers in occupational exposure to formaldehyde—the case of histopathology laboratories. *Mutat Res.* 2011;721:15–20.
35. Fenech M, Holland N, Chang WP, Zeiger E, Bonassi S. The HUMAN MicroNucleus Project—an international collaborative study on the use of the micronucleus technique for measuring DNA damage in humans. *Mutat Res.* 1999;428:271–283.
36. Duthie SJ, Narayanan S, Brand GM, Pirie L, Grant G. Impact of folate deficiency on DNA stability. *J Nutr.* 2002;132:2444S–2449S.
37. Singh NP, McCoy MT, Tice RR, Schneider EL. A simple technique for quantitation of low levels of DNA damage in individual cells. *Exp Cell Res.* 1988;175:184–191.
38. Collins AR. Investigating oxidative DNA damage and its repair using the comet assay. *Mutat Res.* 2009;681:24–32.
39. Collins AR. The comet assay. Principles, applications, and limitations. *Methods Mol Biol (Clifton, N.J.).* 2002;203:163–177.
40. Fenech M. The Genome Health Clinic and Genome Health Nutrigenomics concepts: diagnosis and nutritional treatment of genome and epigenome damage on an individual basis. *Mutagenesis.* 2005;20:255–269.
41. Hart RW, Dixit R, Seng J, et al. Adaptive role of caloric intake on the degenerative disease processes. *Toxicol Sci.* 1999;52:3–12.
42. Heilbronn LK, Ravussin E. Caloric restriction and aging: review of the literature and implications for studies in humans. *Am J Clin Nutr.* 2003;78:361–369.
43. Heydari A, Unnikrishnan A, Lucente LV, Richardson A. Caloric restriction and genomic stability. *Nucleic Acids Res.* 2007;35:7485–7496.
44. Masoro EJ. Overview of caloric restriction and ageing. *Mech Ageing Dev.* 2005;126:913–922.
45. Fenech M, Noakes M, Bowen J, Clifton P. Effect of protein intake, hyperglycaemia and micronutrients on DNA damage and mitogen responsiveness of peripheral blood lymphocytes. *Nutr Diet.* 2008;65:S27–S32.
46. Hassan AM, Abdel-Aziem SH, Abdel-Wahhab MA. Modulation of DNA damage and alteration of gene expression during aflatoxicosis via dietary supplementation of Spirulina (Arthrospira) and whey protein concentrate. *Ecotoxicol Environ Saf.* 2012;79:294–300.
47. Kumar V, Sinha AK, Makkar HP, de Boeck G, Becker K. Dietary roles of non-starch polysaccharides in human nutrition: a review. *Crit Rev Food Sci Nutr.* 2012;52:899–935.
48. Lagergren K, Lindam A, Lagergren J. Dietary proportions of carbohydrates, fat, and protein and risk of oesophageal cancer by histological type. *PLoS ONE.* 2013;8:1–7.
49. Ames B. Micronutrients prevent cancer and delay aging. *Toxicol Lett.* 1998;102–103:5–18.
50. Fenech M, Ferguson L. Vitamins/minerals and genomic stability in humans. *Mutat Res.* 2001;475:1–6.
51. Hageman G, Stierum R, van Herwijnen M, van der Veer M, Kleinjans J. Nicotinic acid supplementation: effects on niacin status, cytogenetic damage, and poly(ADP-ribosylation) in lymphocytes of smokers. *Nutr Cancer.* 1998;32:113–120.
52. Surjana D, Halliday GM, Damian DL. Role of nicotinamide in DNA damage, mutagenesis, and DNA repair. *J Nucleic Acids.* 2010;2010:157591.
53. Fenech M, Bonassi S. The effect of age, gender, diet and lifestyle on DNA damage measured using micronucleus frequency in human peripheral blood lymphocytes. *Mutagenesis.* 2011;26:43–49.
54. Guerreiro C, Cravo ML, Brito M, Vidal PM, Fidalgo PO, Leitão CN. The D1822V APC polymorphism interacts with fat, calcium, and fiber intakes in modulating the risk of colorectal cancer in Portuguese persons. *Am J Clin Nutr.* 2007;85:1592–1597.
55. Smith DF, MacGregor JT, Hiatt RA, et al. Micronucleated erythrocytes as an index of cytogenetic damage in humans: demographic and dietary factors associated with micronucleated erythrocytes in splenectomized subjects. *Cancer Res.* 1990;50:5049–5054.
56. Hartwing A. Role of magnesium in genomic stability. *Mutat Res.* 2001;475:113–121.
57. Anastassopoulou J, Theophanides T. Magnesium-DNA interactions and the possible relation of magnesium to carcinogenesis. Irradiation and free radicals. *Crit Rev Oncol Hematol.* 2002;42:79–91.
58. Sharif R, Thomas P, Zalewski P, Fenech M. The role of zinc in genomic stability. *Mutat Res.* 2012;733:111–121.
59. Song Y, Leonard SW, Traber MG, Ho E. Zinc deficiency affects DNA damage, oxidative stress, antioxidant defenses, and DNA repair in rats. *J Nutr.* 2009;139:1626–1631.
60. Ho E. Zinc deficiency, DNA damage and cancer risk. *J Nutr Biochem.* 2004;15:572–578.
61. Sharif R, Thomas P, Zalewski P, Fenech M. Zinc supplementation influences genomic stability biomarkers, antioxidant activity, and zinc transporter genes in an elderly Australian population with low zinc status. *Mol Nutr Food Res.* 2015;59:1200–1212.
62. Mustafa TG, Monirujjaman MD, Zabeen S, Hossain B. Effect of zinc supplementation on serum zinc level and micronucleus frequency in Bangladeshi adult females with poor socioeconomic status. *AKMMCJ.* 2010;1:4–8.
63. Hwang E-S, Bowen P. DNA damage, a biomarker of carcinogenesis: its measurement and modulation by diet and environment. *Crit Rev Food Sci Nutr.* 2007;47:27–50.
64. Ames B, Wakimoto P. Are vitamin and mineral deficiencies a major cancer risk? *Nat Rev Cancer.* 2002;2:694–704.
65. Ames B. Low micronutrient intake may accelerate the degenerative diseases of aging through allocation of scarce micronutrients by triage. *Proc Natl Acad Sci U S A.* 2006;103:17589–17594.
66. Awodele O, Olayemi S, Alimba C, Egbejiogu C, Akintonwa A. Protective effect of vitamin C and or vitamin E on micronuclei induction by rifampicin in mice. *Tanzania J Health Res.* 2010;2:1–7.
67. Jenab M, Salvini S, van Gils CH, et al. Dietary intakes of retinol, β -carotene, vitamin D and vitamin E in the European Prospective Investigation into Cancer and Nutrition cohort. *Eur J Clin Nutr.* 2009;63:S150–S178.
68. Cooke MS, Evand MD, Mistry N, Lunec J. Role of dietary antioxidants in the prevention of *in vivo* oxidative DNA damage. *Nutr Res Rev.* 2002;15:19–41.
69. Beilby J, Ambrosini GL, Rossi E, Klerk NK, Musk AW. Serum levels of folate, lycopene, β -carotene, retinol and vitamin E and prostate cancer. *Eur J Clin Nutr.* 2010;64:1235–1238.
70. Ferguson L, Philpott M. Nutrition and mutagenesis. *Annu Rev Nutr.* 2008;28:313–329.
71. De Barros KV, De Abreu CG, Xavier RA, et al. Effects of a high fat or a balanced omega 3/omega 6 diet on cytokines levels and DNA damage in experimental colitis. *Nutrition.* 2011;27:221–226.
72. Bishop KS, Erdreich S, Karunasinghe N, et al. An investigation into the association between DNA damage and dietary fatty acid in men with prostate cancer. *Nutrients.* 2015;7:405–422.
73. MacLean C, Newberry S, Mojica W, et al. Effects of omega-3 fatty acids on cancer risk: a systematic review. *JAMA.* 2006;295:403–416.
74. Kiefer F, Wiebel FJ. Caffeine potentiates the formation of micronuclei caused by environmental chemical carcinogens in V79 Chinese hamster cells. *Toxicol Lett.* 1998;96–97:131–136.
75. Tsabar M, Eapen VV, Mason JM, et al. Caffeine impairs resection during DNA break repair by reducing the levels of nucleases Sae2 and Dna2. *Nucleic Acids Res.* 2015;43:6889–6901.
76. Kaufmann WK, Heffernan TP, Beaulieu M, et al. Caffeine and human DNA metabolism: the magic and the mystery. *Mutat Res.* 2003;532:85–102.
77. Wasson GR, McKelvey-Martin VJ, Downes CS. The use of the comet assay in the study of human nutrition and cancer. *Mutagenesis.* 2008;23:153–162.