

Molecular dynamics in tilted bilayer smectic phases: A proton nuclear magnetic resonance relaxometry study

A. Carvalho, P. J. Sebastião, A. C. Ribeiro, H. T. Nguyen, and M. Vilfan

Citation: [The Journal of Chemical Physics](#) **115**, 10484 (2001); doi: 10.1063/1.1413744

View online: <https://doi.org/10.1063/1.1413744>

View Table of Contents: <http://aip.scitation.org/toc/jcp/115/22>

Published by the [American Institute of Physics](#)

PHYSICS TODAY

WHITEPAPERS

ADVANCED LIGHT CURE ADHESIVES

Take a closer look at what these environmentally friendly adhesive systems can do

[READ NOW](#)

PRESENTED BY
 **MASTERBOND**
ADHESIVES | SEALANTS | COATINGS

Molecular dynamics in tilted bilayer smectic phases: A proton nuclear magnetic resonance relaxometry study

A. Carvalho

Centro de Física da Matéria Condensada, Av. Prof. Gama Pinto 2, 1649-003 Lisboa, Portugal and Instituto Superior Técnico, Av. Rovisco Pais, 1049-001 Lisboa, Portugal, and Escola Superior de Tecnologia da Saúde de Lisboa, R. José Carlos dos Santos, 7, 1700-256 Lisboa, Portugal

P. J. Sebastião^{a)} and A. C. Ribeiro

Centro de Física da Matéria Condensada, Av. Prof. Gama Pinto 2, 1649-003 Lisboa, Portugal and Instituto Superior Técnico, Av. Rovisco Pais, 1049-001 Lisboa, Portugal

H. T. Nguyen

CNRS, Centre de Recherche Paul Pascal, Av. A. Schweitzer, F-33600 Pessac, France

M. Vilfan

Jozef Stefan Institute, Jamova 39, 1000 Ljubljana, Slovenia

(Received 25 April 2001; accepted 5 September 2001)

A proton nuclear magnetic resonance (NMR) relaxation study of molecular dynamics in the liquid crystal 4-octylphenyl 2-chloro-4-(4-cyanobenzoyloxy)benzoate (DB₈Cl) is presented. DB₈Cl molecules possess a strong polar terminal group and form, in addition to the nematic phase, three different smectic phases: bilayer smectic A, bilayer smectic C, and anticliniclike smectic C phase. The proton spin-lattice relaxation times were measured in all mesophases over a broad frequency range of six decades by applying conventional and fast field-cycling NMR techniques. The parameters obtained in the analysis of the experimental data give quantitative information on molecular motions, particularly for the tilted smectic phases of DB₈Cl. In contrast to former conjectures, we found that the low-frequency relaxation in the bilayer smectic C phases results from director fluctuations about the layer normal, which occur without distortion of the layers, and from layer undulations, similar to those in the smectic A phase. In the low-temperature bilayer smectic C phase, a considerable slowing-down of molecular translational diffusion is observed. It confirms indirectly the anticlinic character of this mesophase. Measurements of angular dependence of the relaxation times at 60 MHz support the conclusions obtained from the frequency dispersion data. © 2001 American Institute of Physics. [DOI: 10.1063/1.1413744]


I. INTRODUCTION

It is well-known that liquid crystals with strong polar terminal groups often present a great variety of mesophases, including the nematic phase and smectic A and C phases, consisting of monolayers, bilayers or partial-bilayers.^{1–6} The structures of these phases and transitions among them have been an object of detailed study by x-ray diffraction,^{7–11} calorimetry,^{12–15} and nuclear magnetic resonance (NMR).^{16–23} Dielectric measurements and NMR relaxometry gave information on molecular dynamics, predominantly in the nematic and bilayer smectic A phases.^{16–22,24} On the other hand, the structural properties and dynamics in tilted smectic phases have been much less investigated and are still not well understood. In this paper we apply proton NMR relaxometry in a broad frequency range to study the peculiar molecular arrangement and mobility of such phases. We focus on the uniformly tilted bilayer smectic C phase (SmC₂), and on the anticliniclike phase (SmC₇), which appears at lower temperature, and which is—as far its proposed structure is concerned—between the alternating and the bilayer tilted phases.^{24,25}

In NMR relaxometry of liquid crystals we are dealing with several mechanisms, i.e., types of molecular motion, responsible for the relaxation of nuclear spins. As different motional processes occur over different time scales they are usually assumed to be statistically independent and the coupling between them is neglected. Previous studies of molecular dynamics in compounds with polar end-groups have revealed that the relaxation mechanisms in the smectic A phase are not considerably different from those observed in the nematic phase.^{18,19,26–29} If protons are used as the molecular probe, the relaxation in the MHz regime of both mesophases is governed by reorientations of the whole molecule, fast conformational changes within the molecule, and translational self-diffusion. The latter affects only intermolecular proton interactions.

The most significant difference between a liquid crystal and isotropic fluids is observed in the kHz frequency range.^{30,31} Collective molecular reorientations, which are known as director fluctuations (DF), produce here a peculiar dependence of the spin-lattice relaxation rate T_1^{-1} on the Larmor frequency ν . In the nematic phase, T_1^{-1} is proportional to $\nu^{-1/2}$ over several frequency decades as theoretically predicted by Pincus and experimentally definitely con-


^{a)}Electronic mail: pedros@lince.cii.fc.ul.pt

FIG. 1. Chemical structure and polymorphism of the DB₈Cl compound.

firmed by Wölfel, Noack, and Stohrer.^{30,32} In the smectic A phase, collective orientational fluctuations are basically layer undulations. They seem to be less effective than nematic fluctuations as the spin-lattice relaxation time in the smectic A phase exceeds that of the nematic phase almost by one order of magnitude in the kHz regime. The relaxation rate associated with pure layer undulations should go as $T_1^{-1} \propto \nu^{-1}$ due to the two-dimensional character of such fluctuations.²⁶

The relaxation in the tilted smectic phases, composed of nonchiral molecules, has been up to now only marginally included in the papers describing the relaxation in liquid crystals.^{27,33,34} No definite conclusion on the frequency dependence of T_1^{-1} in the kHz regime has been given. In some cases it was ascribed to nematiclike director fluctuations and in others to undulations as in the smectic A phase. This should be the reason for the comment of Vold and Vold that director fluctuations should be operative in tilted smectic C phase,³⁵ but adequate theory and experiment are lacking. The situation has not much improved up to now. Recently, Acosta and Pusiol pointed out that both types of dispersion, $T_1^{-1} \propto \nu^{-1/2}$ and $T_1^{-1} \propto \nu^{-1}$, are observed close to the smectic C-nematic phase transition.²¹ Their observation was only qualitative and assigned to the pre-transitional effects, i.e., pseudo nematic domains within the smectic phase. In order to throw additional light on this problem and on the molecular dynamics in tilted smectic phases generally, we decided to perform a comprehensive study of the NMR proton spin-lattice relaxation in the liquid crystal 4-octylphenyl 2-chloro-4-(4-cyanobenzoyloxy)benzoate, DB₈Cl for short.

The DB₈Cl compound shows four mesophases upon decreasing temperature: One nematic and three bilayer smectic phases (Fig. 1).^{9,15} The structures of the SmA₂ and SmC₂ phases are well understood. In the smectic A₂ phase molecules are oriented perpendicular to the plane of the layer which has a thickness of approximately two molecular lengths as illustrated in Fig. 2. In the SmC₂ phase, the molecular pairs are tilted with respect to the layer normal. The tilt angle θ depends on the temperature and amounts up to 30° just above 107°C. At this temperature the transition into another tilted smectic phase, which is denoted by SmC₇,

FIG. 2. Schematic representation of the structures of the SmA₂, SmC₂, and SmC₇ phases of DB₈Cl. **B** represents the external magnetic field.

takes place. On the basis of dielectric measurements, x-rays diffraction and proton NMR studies an anticlinic type of ordering was proposed for this mesophase.^{9,23,24,36} Within this model the anticlinic arrangement is superimposed on the “regular tilt.” Within a bilayer, one layer of molecules are tilted for an angle $\theta + \beta$ and the other for the angle $\theta - \beta$ away from the layer normal (Fig. 2). β was found to be about $13^\circ \pm 2^\circ$.²³

In this work we present a study of relaxation in all mesophases of DB₈Cl in order to correlate the parameters obtained for the description of molecular dynamics in the SmC₂ and SmC₇ phases with those of the N and SmA₂ phases. In Sec. II experimental procedure and results are described. In Sec. III we present the relaxation mechanisms used in the analysis of the measured relaxation times. The values of the parameters obtained in the fitting procedure are discussed in Sec. IV. They show significant differences between different mesophases and provide an insight into molecular dynamics, particularly for the tilted smectic phases. The conclusions are outlined in Sec. V.

II. EXPERIMENTAL TECHNIQUES AND RESULTS

The proton spin-lattice relaxation time T_1 of DB₈Cl was measured as a function of Larmor frequency (ν), sample's orientation angle (Δ) and temperature (T). In order to cover a broad Larmor frequency range from ~ 500 Hz to 300 MHz, three different NMR spectrometers were used: a Bruker MSL 300 operating at 300 MHz; a Bruker SXP-4/100 for frequencies between 5 and 100 MHz, and a home-built fast field-cycling spectrometer in the low-frequency regime.³⁷ Different pulse sequences were applied according to different frequency ranges. The usual inversion-recovery pulse sequence was used to perform the T_1 measurements in the MHz frequency range. On the fast field-cycling spectrometer the sequence ($B_{H \rightarrow L} - \tau_i - B_{L \rightarrow H} - (\pi/2) - FID$) was used, where τ_i is the evolution time, $(\pi/2)$ is the radio frequency (rf) detecting pulse, and $B_{H \rightarrow L}, B_{L \rightarrow H}$ are transitions between the high and low magnetic fields, respectively.³¹ The measurements of the angular dependence of T_1 were performed at a frequency of 60 MHz on the Bruker SXP spectrometer by using a step motor to achieve the angular resolution of 0.6°.

The DB₈Cl sample was sealed in NMR glass tubes of 0.5 mm diameter under moderate vacuum (< 0.133 Pa). Since the observed T_1 might depend somewhat on the procedure by which the mesophases were oriented in the magnetic field, we took care to perform all measurements after cooling the sample, in the presence of the external magnetic field, from the isotropic phase to the desired temperature at the rate 1 °C/min. In this way the nematic and smectic A phases are homogeneously oriented with the director parallel to the magnetic field everywhere in the sample. The normal to the layers in the smectic A₂ phase has the same direction. When the sample is cooled into the smectic C₂ phase, the molecules are still aligned along the magnetic field.³⁸ This fact implies, however, an inclination of the smectic planes and consequently a decomposition of the uniform structure into domains. Each domain is characterized by the orientation of the layers' normal, i.e., by its polar and azimuthal




FIG. 3. Frequency dependence of the proton spin-lattice relaxation time $T_1(\nu)$ in the N, SmA₂, SmC₂, and SmC_γ phases of DB₈Cl. The dashed lines are guide lines to the characteristic slopes of the DF mechanism ($T_1 \sim \nu^{1/2}$ and $T_1 \sim \nu^1$).

angles with respect to the magnetic field. Whereas the polar angle equals θ in all domains, the azimuthal angle assumes all values between 0 and 2π , with equal probability.

In Fig. 3 we present the experimental results of $T_1(\nu)$ at one temperature for each mesophase of DB₈Cl. The random error limits are estimated to be $\pm 10\%$; for the sake of clarity the experimental error bars are omitted. The T_1 frequency dispersions in the N and SmA₂ phases of DB₈Cl are similar to those found in other liquid crystals.^{18,19,39} In both mesophases T_1 increases from the low to high frequencies but at intermediate frequencies there is a plateau in $T_1(\nu)$ in the SmA₂ phase which is not observed in the N phase. In the SmA₂ phase the dispersion $T_1 \propto \nu^1$ is observed in one decade (10^3 – 10^4 Hz), whereas in the nematic phase the director fluctuations ($T_1 \propto \nu^{1/2}$) obviously govern the relaxation from 10^3 to 10^6 Hz. In this frequency range T_1 in the nematic phase is considerably shorter than in the SmA₂ phase.

A first inspection of the T_1 dispersion in the SmC₂ phase (Fig. 3) shows partly the characteristics of the nematic and partly of the SmA₂ phase, namely: (i) at low frequencies (10^3 Hz– 4×10^4 Hz) T_1 increases with frequency as in the N phase ($T_1 \propto \nu^{1/2}$); (ii) at intermediate frequencies (4×10^3 Hz– 10^6 Hz) $T_1(\nu)$ evidences a leveling-off which resembles the plateau in the SmA₂ phase. In the SmC_γ phase the spin-lattice relaxation time is considerably shorter than in other mesophases over the whole frequency range studied. The shape of the dispersion is similar to the one observed for the SmC₂ phase.

In Fig. 4 the angular dependences of the relaxation rate




FIG. 4. Angular dependence of the proton spin-lattice relaxation rate T_1^{-1} in the smectic phases of DB₈Cl at $\nu = 60$ MHz.

($T_1^{-1}(\Delta)$) are presented for one temperature in each smectic mesophase at the frequency 60 MHz; for the sake of clarity the experimental error bars are omitted. The angular dependence is obtained by rotating the sample about an axis perpendicular to the external magnetic field. Δ denotes the angle of rotation of the sample with $\Delta = 0$ denoting the position at which the cooling from the isotropic phase was performed. In the case of the smectic A₂ phase, the molecular director turned away from the magnetic field by sample rotation, does not realign for quite a long time. This enables the measurements of the spin relaxation rate with the director turned away by Δ from the direction of the magnetic field. Figure 4 shows that $T_1^{-1}(\Delta)$ exhibits a characteristic maximum at 90° in the SmA₂ phase. Rotating the tilted smectic phases, SmC₂ and SmC_γ, results in the corresponding rotation of the layer planes. However, the molecules in each domain adjust their orientation closer to the magnetic field while preserving their tilt angle and the orientation of the layers. The director, which represents also in the tilted smectic phases the preferred direction of long axis, has therefore different orientations in different domains for $\Delta \neq 0$.²³ The angle of the director in each m domain with respect to the magnetic field is denoted by δ_m and depends, naturally, on the angle Δ of the rotation of the sample. In both tilted smectic phases, SmC₂ and SmC_γ, the NMR spectra show a well expressed angular dependence.²³ On the other hand, the spin-lattice relaxation rates are almost independent of the sample orientation as shown in Fig. 4.

III. RELAXATION MECHANISMS

The proton spin-lattice relaxation measurements are quantitatively interpreted by a global target nonlinear least-square fitting minimization procedure in which a linear combination of relaxation mechanisms' models is used. The most important relaxation mechanisms are:

- (1) Translational self-diffusion (SD), modulating *inter*-molecular magnetic dipolar proton interactions;
- (2) director fluctuations (DF), expected to be effective at low frequencies;
- (3) molecular rotations/reorientations (R), acting on the *intra*-molecular interactions; i.e., magnetic dipolar interactions among protons in the same molecule.

Assuming that these relaxation mechanisms are statistically independent, the measured relaxation rate can be interpreted as a sum of three contributions:

$$\frac{1}{T_1} = \left(\frac{1}{T_1} \right)_{\text{SD}} + \left(\frac{1}{T_1} \right)_{\text{DF}} + \left(\frac{1}{T_1} \right)_{\text{R}}. \quad (1)$$

The following relaxation models were used in the fitting procedure of the theory to the experimental data:

- (1) **Anisotropic translational self-diffusion** as described by Vilfan and Žumer for nematic and smectic A phases.^{40,41} Its contribution to the relaxation rate induced by the time-modulation of intermolecular interactions is

$$\left(\frac{1}{T_1}\right)_{\text{SD}} = \frac{9}{8} \left(\frac{\mu_0}{4\pi}\right)^2 \gamma^4 \hbar^2 \frac{n \tau_{D\perp}}{d^3} \times Q \left(\nu \tau_{D\perp}, \frac{\langle r_{\perp}^2 \rangle}{d^2}, \frac{D_{\parallel}}{D_{\perp}}, \frac{l}{d}, \delta \right), \quad (2)$$

where γ is the proton gyromagnetic ratio, n is the number of spins per unit volume, d the distance of closest approach of two proton spins on neighboring molecules, $\langle r_{\perp}^2 \rangle$ is the mean square jump distance, and $\tau_{D\perp}$ the average time between two successive jumps in the plane of the layer for the smectic phases and in the direction perpendicular to the director for the nematic phase. Similarly, l denotes layers' spacing in the smectic phases and molecular length in the nematic phase. The relaxation rate $(T_1^{-1})_{\text{SD}}$ depends further on the ratio D_{\parallel}/D_{\perp} of the diffusion coefficients in two perpendicular directions and on the angle δ between the molecular director and the magnetic field. The dimensionless function Q , which determines the frequency dependence of $(T_1^{-1})_{\text{SD}}$, depends on the details of the diffusion process and on the spatial molecular arrangement. It is different for the nematic and for the smectic A phase and was evaluated numerically for both cases in Refs. 40 and 41. It should be also mentioned that the parameters D_{\parallel} and D_{\perp} are related to the ones referred for the perfectly ordered nematic and smectic A structures (D_{\parallel}^0 and D_{\perp}^0) through the orientational order parameter S^{42}

$$D_{\parallel} = \frac{2D_{\perp}^0 + D_{\parallel}^0}{3} (1 - S) + S D_{\parallel}^0, \quad (3a)$$

$$D_{\perp} = \frac{2D_{\perp}^0 + D_{\parallel}^0}{3} (1 - S) + S D_{\perp}^0. \quad (3b)$$

As there is no specific model for the relaxation induced by translational self-diffusion in the tilted smectic phases, the above model for the SmA phase was used also in the analysis of the SmC₂ and SmC₇ phases. The approximation is reasonable in view of a weak dependence of $(T_1^{-1})_{\text{SD}}$ on the angle between the molecular director and the external magnetic field.

In fitting the contribution of translational self-diffusion, $(T_1^{-1})_{\text{SD}}$, to the experimental data, $\tau_{D\perp}$ was the only free parameter. The values of other parameters used in the fit are: $n \approx 3.4 \times 10^{28}$ spins m^{-3} (calculated for DB₈Cl from a specific density of ~ 1 g/cm³ and taking into account the number of proton spins present in each molecule), $\langle r_{\perp}^2 \rangle/d^2 = 1$; $d \approx 5.0 \times 10^{-10}$ m in the nematic and SmA₂ phases, and $d \approx 4.8 \times 10^{-10}$ m in the SmC phases (in agreement with the decreasing of molecular area with temperature);⁴³ $l \approx 28 \times 10^{-10}$ m (estimated from the DB₈Cl molecule in its stretched conformation). As for D_{\parallel}/D_{\perp} , the values obtained in the literature for similar compounds were used, i.e., D_{\parallel}/D_{\perp} is 1.4 in the nematic phase, 1 in the smectic A₂ and C₂ phases and 0.18 in the smectic C₇ phase.^{40,41,44} Anyway, the value of this parameter, taken within reasonable limits does not affect appreciably the fit;

(2) **director fluctuations** modulate in a slow time scale the residual proton-proton interactions left after averaging by

fast molecular motions. The well known square root frequency dependence $(T_1^{-1})_{\text{DF}} \sim \nu^{-1/2}$, obtained for the nematic phase,^{32,45} is modified if the finite dimension of molecules on one side, and the finite size of the nematic domain on the other side, are taken into account. As a result, the square root regime is limited to the frequency range between $\nu_{C_{\max}}^N$ on the upper side and $\nu_{C_{\min}}^N$ at the lower side. For Larmor frequencies ν larger than $\nu_{C_{\max}}^N$ the dispersion becomes quadratic, whereas it levels-off into a plateau for frequencies smaller than $\nu_{C_{\min}}^N$. Such behavior is described by the following equations for the nematic phase:^{35,46–49}

$$\left(\frac{1}{T_1}\right)_{\text{DF}_N} = \frac{A_N}{\sqrt{\nu}} \left[j_N^{(1)}(\nu, \nu_{C_{\max}}^N, \nu_{C_{\min}}^N, \delta) + \frac{1}{\sqrt{2}} j_N^{(2)}(2\nu, \nu_{C_{\max}}^N, \nu_{C_{\min}}^N, \delta) \right], \quad (4)$$

where

$$j_N^{(k)}(\nu, \nu_{C_{\max}}^N, \nu_{C_{\min}}^N, \delta) = f_{k1}(\delta) \left[g_N \left(\frac{\nu_{C_{\max}}^N}{\nu} \right) - g_N \left(\frac{\nu_{C_{\min}}^N}{\nu} \right) \right] \quad (4a)$$

and

$$g_N(a) = \frac{1}{\pi} \left(\arctan(\sqrt{2a+1}) + \arctan(\sqrt{2a-1}) - \operatorname{arctanh} \left(\frac{\sqrt{2a}}{a+1} \right) \right). \quad (4b)$$

The effectiveness of order fluctuations in relaxing the spins is given by the parameter A_N which depends on the viscoelastic properties of the liquid crystal, on the square of the orientational order parameter S and on the strength of the proton-proton interaction. $\nu_{C_{\max}}^N$ and $\nu_{C_{\min}}^N$ are the high and low cut-off frequencies. The angular functions $f_{k1}(\delta)$ determine the $(T_1^{-1})_{\text{DF}}(\delta)$ and can be found elsewhere.^{46,50}

In the smectic A phase, the director fluctuations are associated with layer undulations. If the compressibility of layers is neglected, such fluctuations preserve the layer spacing and the perpendicular orientation of the molecules with respect to the instantaneous layer normal. They are restricted to wave vectors in the layer plane, i.e., $q_z = 0$ and yield a linear dependence $T_1^{-1} \sim \nu^{-1}$ over a broad frequency range, described by equations^{18,19,26}

$$\left(\frac{1}{T_1}\right)_{\text{DF}_{LU}} = \frac{A_{LU}}{\nu} [j_{LU}^{(1)}(\nu, \nu_{C_{\max}}^{LU}, \nu_{C_{\min}}^{LU}, \delta) + \frac{1}{2} j_{LU}^{(2)}(2\nu, \nu_{C_{\max}}^{LU}, \nu_{C_{\min}}^{LU}, \delta)], \quad (5)$$

$$j_{LU}^{(k)}(\nu, \nu_{C_{\max}}^{LU}, \nu_{C_{\min}}^{LU}, \delta) = f_{k1}(\delta) \left[\frac{2}{\pi} \arctan\left(\frac{\nu_{C_{\max}}^{LU}}{\nu}\right) - \frac{2}{\pi} \arctan\left(\frac{\nu_{C_{\min}}^{LU}}{\nu}\right) \right], \quad (5a)$$

where the same notation is used as in Eq. (4) and LU denotes layer undulations.

As far as we know, there is no theory specific for relaxation induced by director fluctuations in the smectic C phase. Light scattering studies clearly show that in this phase the fluctuations are not limited to wave vectors in the plane of the layer ($q_z=0$) but involve also fluctuations of the director, i.e., of the preferred orientation of the long molecular axes, about the layer normal with $q_z \neq 0$.¹ In consequence, the smectic C phase is more turbid in appearance than the smectic A phase. The fluctuations with $q_z \neq 0$, which refer to the rotation of the director $\mathbf{n} \equiv (n_x, n_y, n_z)$ without any distortion of the layers, have the mean square amplitude

$$\langle |n_y(\mathbf{q})|^2 \rangle = \frac{k_B T}{(B_1 q_x^2 + B_2 q_y^2 + B_3 q_z^2 + 2B_{13} q_x q_z) V}, \quad (6)$$

where B_1 , B_2 , B_3 , and B_{13} are the elastic constants of the smectic C phase which are roughly of the same order of magnitude as the nematic elastic constants and V is the sample volume.¹ The component n_y of the director is perpendicular to the plane determined by the layer normal and by the average direction of \mathbf{n} . Assuming that the characteristic damping times of the viscoelastic relaxation modes are given by⁴⁵

$$\tau_y(\mathbf{q}) = \frac{\eta}{B_1 q_x^2 + B_2 q_y^2 + B_3 q_z^2 + 2B_{13} q_x q_z}, \quad (7)$$

where η is an effective viscosity, we sum up the contributions of all modes by integrating over q from 0 to ∞ . A dispersion similar to the nematic phase, i.e., $T_1^{-1} \propto \nu^{-1/2}$, is found. On the other hand, the undulations of the layers in the SmC phase should give a $T_1^{-1} \propto \nu^{-1}$ dependence as in the smectic A phase. Neglecting the coupling term between both types of fluctuations,^{51,52} we fit the experimental T_1 values in the SmC phase by adding up a contribution of nematiclike fluctuations with strength A_N and of smectic fluctuations with strength A_{LU} . It should be mentioned that in the chiral smectic C* phase additional modes related to the distortions of the helix should be taken into account.^{53,54}

In fitting the total relaxation rate [Eq. (1)] to the experimental data, A_N , A_{LU} , and the cut-off frequencies can be obtained. However, the high cut-off frequencies $\nu_{C_{\max}}^N$ and $\nu_{C_{\max}}^{LU}$ do not have much influence on the model fits since in the frequency region where they constrain the relaxation (MHz regime) the DF mechanism is overwhelmed by translational self-diffusion and local molecular reorientations. Therefore, fixed values $\nu_{C_{\max}}^N = 100$ MHz and $\nu_{C_{\max}}^N = 60$ MHz were used in the N and SmC fits and $\nu_{C_{\max}}^{LU} = 100$ MHz in all smectic phases;

(3) **molecular rotations/reorientations.** Their effect upon nuclear relaxation has been intensively worked out to




FIG. 5. Frequency dispersion of the proton spin-lattice relaxation time in the nematic phase of DB₈Cl. The solid line represents the fit of Eq. (1) to the experimental data. The contributions of different relaxation mechanisms are indicated. The values of the fitting parameters are given in Table I.

explain the deuterium spin relaxation rate in the MHz regime in smectic and nematic compounds.^{16,22,35,55,56} Here a rotational diffusion model which assumes that each molecule moves in time in a sequence of small angular steps caused by collisions with surrounding molecules and under the influence of a mean square torque exerted by these molecules is usually used.^{57–59} In the analysis of our data, however, the local molecular rotations and conformational changes give a smaller contribution than in the case of deuterons. They are effective only at the highest frequencies studied. We used, therefore, a simpler model evaluated by Vold *et al.*⁶⁰ The possible anisotropy of the local molecular long-axis-reorientations is neglected,

$$\left(\frac{1}{T_1} \right)_R = \frac{9}{8} \left(\frac{\mu_0}{4\pi} \right)^2 \gamma^4 \hbar^2 (J^{(1)}(\nu, \delta, A^{(0)}, A^{(1)}, A^{(2)}, \tau_{\perp}, \tau_{\parallel}, \tau_{\gamma}, S) + J^{(2)}(2\nu, \delta, A^{(0)}, A^{(1)}, A^{(2)}, \tau_{\perp}, \tau_{\parallel}, \tau_{\gamma}, S)), \quad (8)$$

where the spectral densities $J^{(k)}$ are given by Vold *et al.*⁶⁰ $A^{(m)}$ are geometric factors which depend on the intramolecular interproton distances and orientations. τ_{\perp} , τ_{\parallel} , and τ_{γ} are the correlation times associated to the molecular reorientations in the director frame (τ_{\perp} , τ_{\parallel}) and to the molecular fast rotations around the long molecular axis in the molecular frame (τ_{γ}), respectively. S is the orientational order parameter. The geometric factors $A^{(m)}$ are estimated from the intramolecular interproton distances and orientations with respect to a molecular long axes: $A^{(0)} \approx 4 \times 10^{57} \text{ m}^{-6}$, $A^{(1)} \approx 3 \times 10^{57} \text{ m}^{-6}$, and $A^{(2)} \approx 4 \times 10^{57} \text{ m}^{-6}$. The orientational order parameter S for DB₈Cl is obtained from proton spectra at different temperatures. Its values are $S=0.46$, 0.63 , 0.67 , 0.73 , 0.73 , 0.75 , and 0.75 consequently for the N ($T=162^\circ\text{C}$), SmA₂ ($T=145^\circ\text{C}$ and $T=135^\circ\text{C}$), SmC₂ ($T=115^\circ\text{C}$ and $T=112^\circ\text{C}$), and SmC₇ ($T=105^\circ\text{C}$ and $T=102^\circ\text{C}$) phases. τ_{\perp} , τ_{\parallel} , and τ_{γ} were the fitting parameters of this model.

In the analysis of the angular dependence of T_1^{-1} results in the SmA and SmC phases different treatments are required. In the SmA₂ phase the director is always parallel to layers' normal and, therefore, $\delta \equiv \Delta$. In the case of the SmC₂ and SmC₇ phases it is necessary to take into account a distribution of director orientations, as the sample is rotated in the magnetic field, as described at the end of Sec. II. Besides,




FIG. 6. Frequency dispersion of the proton spin-lattice relaxation time in the SmA_2 phase of DB_8Cl . The solid line represents the fit of Eq. (1) to the experimental data. The contributions of different relaxation mechanisms are indicated. The values of the fitting parameters are given in Table I.

in the SmC_γ phase an additional tilt angle β of the molecules has to be taken into account in order to reflect the particular structure of this phase,²³ similar to the antiferroelectric phase formed by chiral molecules. To account for these details in calculating the relaxation rate, the functions $f_{kj}(\delta)$ which appear in Eqs. 4 and 5 [and which are also implicitly involved as spectral densities prefactors in Eqs. (2) and (8)] are replaced by their values averaged over all domains

$$\overline{f_{kj}(\Delta)} = \sum_{m=1}^N f_{kj}(\delta_m(\Delta)) / N. \quad (9)$$

In Eq. (9) the summation is over all domains with different director orientations. In the case of the SmC_2 phase, the angle $\delta_m(\Delta)$ is obtained by minimizing the free energy in the smectic C_2 phase taking into account the elastic and the magnetic energy of the liquid crystal.²³

IV. ANALYSIS AND DISCUSSION

In the analysis of the DB_8Cl experimental data we reduced the number of fitted parameters as much as possible. As a rule six fitting parameters were considered in the optimization process: $\tau_{D\perp}$, A (A_N , A_{LU}), $\nu_{C\min}$ ($\nu_{C\min}^N$, $\nu_{C\min}^{LU}$), τ_{\perp} , τ_{\parallel} , and τ_{γ} . This might seem to be a large number of parameters, but fortunately different relaxation mechanisms




FIG. 7. Angular dependence of the proton spin-lattice relaxation rate T_1^{-1} in the SmA_2 phase of DB_8Cl at $T=135^\circ\text{C}$ and $\nu=60$ MHz. The solid line represents the fit of Eq. (1) to the experimental data. The contributions of R and SD relaxation mechanisms are indicated (the DF_{LU} contribution is negligible at this frequency). The values of the fitting parameters are identical to those used in the fit presented in Fig. 6.




FIG. 8. Frequency dispersion of the proton spin-lattice relaxation time in the SmC_2 phase of DB_8Cl . The solid line represents the fit of Eq. (1) to the experimental data. The contributions of different relaxation mechanisms are indicated. The values of the fitting parameters are given in Table I.

act in different frequency ranges. Therefore, the total dispersion curve, which extends over six frequency decades, gives quite reliable values for the above parameters. The results of the fit are further supported by the analysis of angular dependences. The values of parameters, which provide the best fit to the experimental data, are summarized in Table I.

The contributions of different relaxation mechanisms calculated with these parameters and the total relaxation rates are plotted, together with experimental data, in Figs. 5–11.

A. Nematic phase

As shown in Fig. 5, $T_1(\nu)$ in the nematic phase reveals a typical distribution of relaxation mechanisms over the studied frequency range. The DF mechanism dominates the relaxation from 1 kHz to ~ 10 MHz exhibiting the $T_1 \sim \nu^{1/2}$ dependence. Its “strength” $A_N = 2600 \text{ s}^{-3/2}$ at 162°C is slightly larger than the values obtained for similar compounds.^{18,19} Translational self-diffusion and rotations/reorientations prevail as relaxation mechanisms in the high frequency range. The self-diffusion coefficient D_{\perp} , estimated from $D_{\perp} = \langle r_{\perp}^2 \rangle / 4\tau_{D\perp}$, amounts to $9 \times 10^{-11} \text{ m}^2/\text{s}$. The measurements of the angular dependence of T_1^{-1} in the nematic phase could not be performed without special equipment or a more elaborated experimental technique, as the director adjusts to the magnetic field in a very short time.^{19,20}




FIG. 9. Angular dependence of the proton spin-lattice relaxation rate T_1^{-1} in the SmC_2 phase of DB_8Cl at $T=112^\circ\text{C}$ and $\nu=60$ MHz. The solid line represents the fit of Eq. (1) to the experimental data. The contributions of different relaxation mechanisms effective at this frequency are indicated (the DF_{LU} contribution is negligible at 60 MHz). The values of the fitting parameters are identical to those used in the fit presented in Fig. 8.




FIG. 10. Frequency dispersion of the proton spin-lattice relaxation time in the SmC_7 phase of DB_8Cl . The solid line represents the fit of Eq. (1) to the experimental data. The contributions of different relaxation mechanisms are indicated. The values of the fitting parameters are given in Table I.

B. Bilayer smectic A phase (SmA_2)

The T_1 versus ν in the SmA_2 phase is presented in Fig. 6 and T_1^{-1} versus Δ in Fig. 7. The experimental data clearly indicate that a relaxation mechanism yielding $T_1 \sim \nu^{-1}$ frequency dependence is effective below 10^4 Hz. It is also indirectly responsible for the “shoulder” in the dispersion curve, which is not observed in the nematic phase. This mechanism can be—in view of its specific dispersion—assigned most probably to layer undulations, though the impact of layers’ compressibility on cut-off frequencies has not yet been experimentally clarified.⁶¹ In our case the low cut-off frequency could not be precisely determined as no true low-frequency plateau was detected in $T_1(\nu)$. The value of $\nu_{C_{\min}}^{LU}$ presented in Table I is merely indicative.

The relaxation in the intermediate frequency range (10 kHz–20 MHz) is dominated by translational self-diffusion, which is slower than in the nematic phase. The estimated diffusion coefficient is $D_{\perp} = 5 \times 10^{-11} \text{ m}^2/\text{s}$. Rotations/reorientations are important only in the high-frequency regime (> 20 MHz).

All parameters obtained in the fit of dispersion curves can be here additionally tested by the angular dependence of the relaxation rate. Parameters listed in Table I were used to plot the solid line in Fig. 7. Obviously it matches well the experimental data. The observed maximum at $\Delta = 90^\circ$ is due mainly to the contribution of the rotation/reorientations mechanism.




FIG. 11. Angular dependence of the proton spin-lattice relaxation rate T_1^{-1} in the SmC_7 phase of DB_8Cl at $T = 105^\circ\text{C}$ and $\nu = 60$ MHz. The solid line represents the fit of Eq. (1) to the experimental data. The contributions of different relaxation mechanisms effective at this frequency are indicated (the DF_{LU} contribution is negligible at 60 MHz). The values of the fitting parameters are identical to those used in the fit presented in Fig. 10.

C. Bilayer tilted smectic phase (SmC_2)

Measurements of the frequency dependence of T_1^{-1} in the bilayer SmC_2 phase at $T = 112^\circ\text{C}$ are presented in Fig. 8. The contributions of rotations/reorientations and particularly of self-diffusion to T_1^{-1} have obviously increased compared to the nematic and smectic A phases. This is due to an increase of parameters τ_{\perp} , τ_{\parallel} , τ_{γ} for rotations and of $\tau_{D\perp}$ for translational diffusion with decreasing temperature, as expected for thermally activated motions. The angular dependence of T_1^{-1} at 60 MHz, presented in Fig. 9, is much less pronounced than in the SmA_2 phase and hardly exceeds the experimental error. This is due to a relatively smaller contribution of local rotations to the total relaxation rate at 60 MHz. The flat $T_1^{-1}(\Delta)$ curve is well explained by the compensating effect of the self-diffusion and rotations/reorientations relaxation mechanisms. Since both the frequency and the angular dependences of T_1^{-1} in the MHz regime can be described by the same values of fitted parameters, we are reasonably sure that the interpretation is correct in spite of a large number of parameters involved.

A particularly interesting point in our study is the proton relaxation in the kHz regime. According to earlier studies of the smectic C phase, director fluctuations should be dominant in this frequency range, but the frequency dependence of their contribution to T_1^{-1} has not been well establi-

TABLE I. Parameters describing molecular dynamics in DB_8Cl as obtained from the fit of theoretical expressions (described in the text) to the measured T_1^{-1} vs ν^{-1} and T_1^{-1} vs Δ dependences.

	Phase T ($^\circ\text{C}$)	N 162	SmA_2		SmC_2		SmC_7	
			145	135	115	112	105	102
SD	$\tau_{D\perp} (10^{-9} \text{ s})$	0.68	1.2	2.1	4.3	6.3	19	27
DF	$\nu_{C_{\min}}^N (10^3 \text{ Hz})$	~ 0.03	1.5	1.5	1.5	1.6
	$A_N (10^3 \text{ s}^{-3/2})$	2.6	1.9	2.0	4.0	6.0
	$\nu_{C_{\min}}^{LU} (10^3 \text{ Hz})$...	0.36	0.32	1.5	1.5	1.5	1.5
	$A_{LU} (10^4 \text{ s}^{-2})$...	2.2	2.6	13	13	15	16
R	$\tau_{\perp} (10^{-10} \text{ s})$	0.73	0.78	1.0	2.3	3.0	3.4	3.4
	$\tau_{\parallel} (10^{-11} \text{ s})$	0.10	0.18	0.28	0.97	1.4	1.7	1.8
	$\tau_{\gamma} (10^{-12} \text{ s})$	0.63	1.1	1.7	8.2	11	17	18

shed.^{21,27,34} We find that T_1^{-1} of DB₈Cl in the kHz range cannot be fitted either with $T_1^{-1} \sim \nu^{-1/2}$ nor with $T_1^{-1} \sim \nu^{-1}$ dispersion alone. Only a superposition of both dispersion “laws” is capable of explaining the experimental data (Fig. 8). This fact is not surprising if we take into account that in the tilted smectic phase the director can reorient about the layer normal without disturbing the layer thickness. This motion, described by Eqs. (6) and (7), leads to a relaxation rate proportional to $\nu^{-1/2}$ in the range between the low and the high cut-off frequencies, as shown in Sec. III-2. The fit, presented in Fig. 8, shows that the contribution of in-layer fluctuations of the director exceeds that of layer undulations in the whole frequency range studied. Its effectiveness is given by the parameter $A_N = 2.0 \times 10^3 \text{ s}^{-3/2}$ (see Table I), which is only slightly smaller than in the nematic phase. This might seem unusual since the director in the smectic C is “free” to rotate only in the plane of the layer, whereas the out-of-plane fluctuations are much more restricted. However, a similar value of the parameters A_N for both mesophases may result from increased viscosity and orientational order parameter in the SmC₂ phase.⁶² The constant describing the effectiveness of layer undulations, A_{LU} , is roughly 5 times larger in the SmC₂ phase than in the SmA phase because of increased positional, i.e., smectic ordering. The values of the cut-off frequencies, $\nu_{C_{\min}}^N$ and $\nu_{C_{\min}}^{LU}$, given in Table I, are only indicative. They should be larger than in the nematic or smectic A phase, respectively, as a low-frequency plateau in the dispersion curve between 10^3 and 10^4 Hz is here clearly observed. But the data do not allow to decide whether this is caused by the onset of the frequency cutoff or by the transition from Zeeman into the dipolar spin “reservoir” which takes place in the same frequency range.

D. Anticliniclike smectic C phase (SmC₇)

Frequency and angular dependences of the proton relaxation in the SmC₇ phase are presented in Figs. 10 and 11, together with the best fit of Eq. (1) to the experimental data. The measurements are explained basically by the same relaxation mechanisms as in the SmC₂ phase, but there are some important differences. Translational self-diffusion is considerably slower in the low-temperature tilted phase. Consequently its importance in the spin relaxation process is increased. Self-diffusion is here the dominating relaxation mechanism in the whole frequency range between 2×10^4 Hz and 2×10^7 Hz, i.e., over three frequency decades. The in-plane correlation time for translational diffusion, $\tau_{D\perp}$, exhibits an abrupt increase by a factor of 3 at the transition from SmC₂ to the SmC₇ phase. This indicates that in the SmC₇ phase the diffusion in the bilayer is hindered by the formation of molecular pairs with anticlinic arrangement. It should be also stressed that the best fit was obtained in the limit $D_{\parallel}^0/D_{\perp}^0 \rightarrow 0$. Obviously a strong slowing-down of molecular exchange between the layers takes place supporting again the conjecture of anticlinic molecular ordering. The contribution of in-plane director fluctuations to T_1^{-1} in DB₈Cl is increased as well, most probably due to the increase in the viscosity coefficient. The determination of the contribution of layer undulations is less reliable as there is no

frequency region where this contribution would dominate. Therefore, the magnitude of $(T_1)_{DFLU}$ was kept here at values close to the ones obtained in the SmC₂ phase. It was not possible to perform a more detailed study of the T_1 dispersion at low frequencies since the T_1 values in the SmC₇ phase are so small that they are close to the fast field-cycling spectrometer’s experimental limit.

V. CONCLUSIONS

In conclusion we may say that a comprehensive study of proton spin-lattice relaxation in the nematic and bilayer smectic phases of DB₈Cl has been performed. Though the same relaxation mechanisms are basically present in all mesophases, their parameters obtained in the fit reveal the specific character of each mesophase. The correlation times associated with local molecular reorientations, which are the most effective relaxation mechanism in the high-frequency regime, increase with decreasing temperature in the usual way, characteristic of thermally activated motions. On the other hand, translational self-diffusion dominating the proton relaxation at intermediate frequencies, shows an abrupt increase in the intra-layer correlation time at the transition from the SmC₂ into SmC₇ phase. Besides, the best fit in this phase is obtained in the limit $D_{\parallel}^0/D_{\perp}^0 \rightarrow 0$ indicating a considerable decrease of out-of-layer molecular diffusion as well. These facts—slowing-down of intra-layer and particularly of out-of-layer molecular diffusion—indirectly support the anticlinic arrangement of molecular pairs in the SmC₇ phase. A similar effect was observed in antiferroelectric liquid crystals, where the anticlinic molecular ordering diminishes the diffusion between layers by two orders of magnitude compared to ferroelectric liquid crystals with uniform short range tilt.⁶³

Director fluctuations dominate the relaxation in the lowest frequency regime under study. In the nematic phase the usual square root dispersion profile is observed. In the SmA₂ phase a linear frequency dependence of $(T_1^{-1})_{DF}$ explains the experimental data. It is ascribed to the undulations of the layers. In contrast to former conjectures, we find that the T_1^{-1} low-frequency dispersion in the SmC₂ and SmC₇ phases can be explained only if—in addition to layer undulations—fluctuations of the molecular director about the layer normal, which occur without distortion of the layers, are taken into account. Both types of fluctuations are well-known from optical studies but have not been identified by NMR so far.⁶² The calculated contribution of the in-plane director fluctuations in the tilted smectic phases yields a dispersion $(T_1^{-1})_{DF} \sim \nu^{-1/2}$, similar to nematic fluctuations. However, the nematiclike T_1 dispersion in the SmC₂ and SmC₇ phases is not a matter of nematic domains in the smectic phase or a pretransitional effect but an intrinsic property of tilted smectic phases.

ACKNOWLEDGMENTS

The authors wish to thank Fundação para a Ciência e Tecnologia (FCT) through projects PBIC/C/CTM/1935/95 and PRAXIS XXI 3/3.1/MMA/1769/95. One of the authors

(A.C.) thanks FCT for Grant No. PRAXIS XXI BD/2900/94 and M.V. acknowledges the Ministry of Science and Technology of Slovenia, Grant No. P1-024. We also thank Dr. Gabriel Feio for the experimental help with the Bruker MSL spectrometer, and Dr. Benoit Heinrich for scientific discussions.

- ¹S. Chandrasekhar, *Liquid Crystals* (Cambridge University Press, Cambridge, 1992).
- ²J. Goodby, in *Handbook of Liquid Crystals*, edited by D. Demus, J. W. Goodby, G. W. Gray, H. W. Spiess, and V. Vill (Wiley-VCH, New York, 1998).
- ³A. Leadbetter, J. L. Durrant, and M. Rugman, *Mol. Cryst. Liq. Cryst.* **34**, 231 (1977).
- ⁴P. Barois, J. Prost, and T. C. Lubensky, *J. Physique* **46**, 391 (1985).
- ⁵J. Prost and P. Barois, *J. de Chim. Phys.* **80**, 66 (1983).
- ⁶H. Nguyen, *J. de Chim. Phys.* **80**, 83 (1983).
- ⁷F. Hardouin and A. M. Levelut, *J. Physique* **41**, 41 (1980).
- ⁸F. Hardouin, A. M. Levelut, and G. Sigaud, *J. Physique* **42**, 71 (1981).
- ⁹F. Hardouin, H. Nguyen, and A. M. Levelut, *J. Phys. (France) Lett.* **43**, L1779 (1982).
- ¹⁰N. Nuu Tinh, F. Hardouin, and C. Destrade, *J. Physique* **43**, 1127 (1982).
- ¹¹F. Hardouin, A. Levelut, G. Sigaud, M. Achard, N. Tinh, and H. Gasparoux, *Symmetries and Broken Symmetries in Condensed Matter Physics*, N. Boccara (IDSET-Paris) (1981).
- ¹²V. Raja, R. Shashidhar, B. Ratna, G. Heppke, and C. Bahr, *Phys. Rev. A* **37**, 303 (1988).
- ¹³G. Sigaud, F. Hardouin, M. F. Achard, and A. M. Levelut, *J. Physique* **42**, 107 (1981).
- ¹⁴A. Levelut, R. Tarento, F. Hardouin, M. Achard, and G. Sigaud, *Phys. Rev. A* **24**, 2180 (1981).
- ¹⁵Y. H. Jeong, K. J. Stine, C. W. Garland, and H. Nguyen, *Phys. Rev. A* **37**, 3465 (1988).
- ¹⁶R. Y. Dong, *Nuclear Magnetic Resonance of Liquid Crystals* (Springer-Verlag, Berlin, 1994).
- ¹⁷A. Ribeiro, P. Sebastião, and M. Vilfan, *Liq. Cryst.* **3**, 937 (1988).
- ¹⁸P. J. Sebastião, A. C. Ribeiro, H. Nguyen, and F. Noack, *Z. Naturforsch., A: Phys. Sci.* **A48**, 851 (1993).
- ¹⁹P. J. Sebastião, A. C. Ribeiro, and F. Noack, *J. Phys. II* **5**, 1707 (1995).
- ²⁰S. Struppe and F. Noack, *Liq. Cryst.* **20**, 595 (1996).
- ²¹R. Acosta and D. Pusiol, *Phys. Rev. E* **60**, 1808 (1999).
- ²²R. Y. Dong, *Phys. Rev. E* **60**, 5631 (1999).
- ²³A. Carvalho, P. Sebastião, A. Ferraz, A. Ribeiro, and H. Nguyen, *Eur. Phys. J. E* **2**, 351 (2000).
- ²⁴L. Benguigui and F. Hardouin, *J. Phys. (France) Lett.* **45**, L179 (1984).
- ²⁵Y. Galerne and L. Liebert, *Phys. Rev. Lett.* **64**, 906 (1990).
- ²⁶R. Blinc, M. Luzar, M. Vilfan, and M. Burgar, *J. Chem. Phys.* **63**, 3445 (1975).
- ²⁷R. Blinc, M. Vilfan, M. Luzar, J. Seliger, and V. Zagar, *J. Chem. Phys.* **68**, 303 (1978).
- ²⁸F. Noack, M. Notter, and W. Weiß, *Liq. Cryst.* **3**, 907 (1988).
- ²⁹F. Noack and K. H. Schweikert, in *The Molecular Dynamics of Liquid Crystals*, edited by G. Luckhurst and C. A. Veracini (Kluwer Academic, Netherlands, 1994).
- ³⁰W. Wölfel, F. Noack, and M. Stohrer, *Z. Naturforsch., A: Phys. Sci.* **30**, 437 (1975).
- ³¹F. Noack, *Prog. Nucl. Magn. Reson. Spectrosc.* **18**, 171 (1986).
- ³²P. Pincus, *Solid State Commun.* **7**, 415 (1969).
- ³³T. Mugele, V. Graf, W. Wölfel, and F. Noack, *Z. Naturforsch., A: Phys. Sci.* **35**, 924 (1980).
- ³⁴C. Cruz, J. L. Figueirinhas, P. J. Sebastião, A. C. Ribeiro, F. Noack, H. T. Nguyen, B. Heinrich, and D. Guillon, *Z. Naturforsch., A: Phys. Sci.* **51**, 155 (1996).
- ³⁵R. Vold and R. Vold, in *The Molecular Dynamics of Liquid Crystals*, edited by G. Luckhurst and C. A. Veracini (Kluwer Academic, Netherlands, 1994), p. 2.
- ³⁶A. Carvalho, P. J. Sebastião, and H. T. Nguyen, *Mol. Cryst. Liq. Cryst.* **331**, 89 (1999).
- ³⁷E. Rommel, K. Mischker, G. Osswald, K. H. Schweikert, and F. Noack, *J. Magn. Reson.* (1969–1992) **70**, 219 (1986).
- ³⁸R. A. Wise, D. H. Smith, and J. W. Doane, *Phys. Rev. A* **7** (4), 1366 (1973).
- ³⁹K. H. Schweikert and F. Noack, *Z. Naturforsch., A: Phys. Sci.* **44**, 597 (1989).
- ⁴⁰S. Žumer and M. Vilfan, *Phys. Rev. A* **17**, 424 (1978).
- ⁴¹M. Vilfan and S. Žumer, *Phys. Rev. A* **21**, 672 (1980).
- ⁴²R. Blinc, M. Burgar, M. Luzar, J. Pirš, I. Zupančič, and S. Žumer, *Phys. Rev. Lett.* **33**, 1192 (1974).
- ⁴³B. Heinrich, *Etude de la structure et propriétés mésomorphes de séries homologues et de mélanges de composés smectogènes*, Ph.D. thesis, Université Louis Pasteur de Strasbourg (1993).
- ⁴⁴G. J. Krüger, *Phys. Rep.* **82**, 229 (1976).
- ⁴⁵P. de Gennes and J. Prost, *The Physics of Liquid Crystals* (Oxford University Press, 2nd ed., 1993), section 5.2.5.
- ⁴⁶J. W. Doane, C. E. Tarr, and M. A. Nickerson, *Phys. Rev. Lett.* **33**, 620 (1974).
- ⁴⁷R. Blinc, M. Vilfan, and V. Rutar, *Solid State Commun.* **17**, 171 (1975).
- ⁴⁸I. Zupančič, V. Žagar, R. Rožmarin, I. Lesvnik, F. Kogovšek, and R. Blinc, *Solid State Commun.* **18**, 1591 (1976).
- ⁴⁹J. H. Freed, *J. Chem. Phys.* **66**, 4183 (1977).
- ⁵⁰P. J. Sebastião, M. H. Godinho, A. C. Ribeiro, D. Guillon, and M. Vilfan, *Liq. Cryst.* **11**, 621 (1992).
- ⁵¹R. Blinc, M. Vilfan, and J. Seliger, *Bull. Magn. Reson.* **5**, 51 (1983).
- ⁵²B. Zalar, O. Lavrentovich, H. Zeng, and D. Finotello, *Phys. Rev. E* **62**, 2252 (2000).
- ⁵³I. Mušević, A. Rastegar, M. Čepič, B. Žekš, M. Čopič, D. Moro, and G. Heppke, *Phys. Rev. Lett.* **77**, 1769 (1996).
- ⁵⁴A. Ferraz, A. C. Ribeiro, and H. T. Nguyen, *Mol. Cryst. Liq. Cryst.* **331**, 67 (1999).
- ⁵⁵R. Y. Dong and M. Cheng, *J. Chem. Phys.* **113**, 3466 (2000).
- ⁵⁶R. Y. Dong, A. Carvalho, P. J. Sebastião, and H. T. Nguyen, *Phys. Rev. E* **62**, 3679 (2000).
- ⁵⁷P. Nordio and P. Busolin, *J. Chem. Phys.* **55**, 5485 (1971).
- ⁵⁸A. Ferrarini, G. J. Moro, and P. L. Nordio, *Liq. Cryst.* **8**, 593 (1990).
- ⁵⁹R. Y. Dong, *Phys. Rev. A* **43**, 4310 (1991).
- ⁶⁰R. R. Vold and R. G. Vold, *J. Chem. Phys.* **88**, 1443 (1988).
- ⁶¹M. Vilfan, M. Kogoj, and R. Blinc, *J. Chem. Phys.* **86**, 1055 (1987).
- ⁶²Y. Galerne, J. Martinand, G. Durand, and M. Veyssie, *Phys. Rev. Lett.* **29**, 562 (1972).
- ⁶³B. Zalar, A. Gregorovič, and R. Blinc, *Phys. Rev. E* **62**, R37 (2000).