

Portuguese Journal of

Public Health

Port J Public Health , DOI: 10.1159/000546311

Received: May 4, 2023

Accepted: April 23, 2025

Published online: May 26, 2025

Social and psychological determinants of COVID-19 vaccination intention in European and North American adults: a systematic review

Almeida-Silva M, Vieira L, Grilo A, Pedro L, Coelho A, Carolino E, Umanets O, Andrade G

ISSN: 2504-3137 (Print), eISSN: 2504-3145 (Online)

<https://www.karger.com/PJP>

Portuguese Journal of Public Health

Disclaimer:

Accepted, unedited article not yet assigned to an issue. The statements, opinions and data contained in this publication are solely those of the individual authors and contributors and not of the publisher and the editor(s). The publisher and the editor(s) disclaim responsibility for any injury to persons or property resulting from any ideas, methods, instructions or products referred to the content.

Copyright:

This article is licensed under the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC) (<http://www.karger.com/Services/OpenAccessLicense>). Usage and distribution for commercial purposes requires written permission.

© 2025 The Author(s). Published by S. Karger AG, Basel on behalf of NOVA National School of Public Health

Social and Psychological Determinants of COVID-19 Vaccination Intention in European and North American Adults – A Systematic Review

ABSTRACT

Since the beginning of the SARS-CoV-19 pandemic, tremendous efforts have been made to address this public health emergency of international concern. The control of the disease has essentially depended on vaccination programs. Many recent studies continue to explore the factors related to the COVID-19 vaccine acceptance. This systematic review was conducted following PRISMA guidelines. PubMed, Scopus, Web of Science, EBSCO, CINAHL, EMBASE and PsycINFO were searched to gather all relevant data on the social and psychological determinants of intention to vaccinate against COVID-19. ROBIS tool was used to evaluate the risk of bias. The main objective of this systematic review was to identify the main social and psychological determinants responsible for the choice to vaccinate against COVID-19 in the adult population of Europe and North America. This systematic review examined literature identified through seven databases yielding 576 PRISMA records, being used 28 articles. The results suggest that older people, higher education level and Caucasian have a stronger intention to be vaccinated. Many studies confirm that beliefs about vaccine costs and benefits are related with vaccine intention. Social concerns, perceived severity of COVID-19 and perceived risk of being infected are confirmed in most studies as determinants of vaccination intention. Conspiracy beliefs seem to promote vaccine hesitance. The results point to some useful conclusions for promoting vaccination in future pandemic situations. In addition to the targeting of the most vaccine-resistant groups, this study suggests the main themes that should be focused on future public communication to promote vaccination.

KEYWORDS: COVID-19; Vaccination; Intention, Hesitancy, Acceptance, Determinants, Psychological

Determinantes Sociais e Psicológicos da Intenção de Vacinação contra a COVID-19 em adultos Europeus e Norte Americanos - Uma Revisão Sistemática

RESUMO

Desde o início da pandemia de SARS-CoV-19, foram feitos esforços enormes para abordar esta emergência de saúde pública de preocupação internacional. O controlo da doença tem dependido, essencialmente, de programas de vacinação. Muitos estudos recentes continuam a explorar os fatores relacionados à aceitação da vacina da COVID-19. Com o objetivo de reunir todos os dados relevantes sobre os determinantes da intenção de se vacinar contra a COVID-19, o objetivo desta revisão da literatura que teve por base a metodologia PRISMA foi identificar os principais determinantes socio-demográficos e psicológicos responsáveis pela escolha da população adulta em se vacinar contra a COVID-19. Os resultados sugerem que, na Europa e América do Norte, pessoas mais velhas, com maior nível de educação e caucasianos têm uma intenção mais forte de serem vacinados. Muitos estudos confirmam que as crenças sobre os custos e benefícios da vacina estão relacionadas com a intenção de vacinação. Preocupações sociais, percepção da gravidade da COVID-19 e percepção do risco de infeção são confirmadas na maioria dos estudos como determinantes da intenção de vacinação. Crenças conspiratórias parecem promover hesitação em relação à vacina. Os resultados apontam para

algumas conclusões úteis para a promoção da vacinação em futuras situações de pandemia. Para além de identificar os grupos mais resistentes à vacinação, o presente estudo aponta os temas principais que devem ser considerados na comunicação com as populações para promover a vacinação.

PALAVRAS-CHAVE: COVID-19; Vacinação; Intenção; Hesitação; Aceitação; Determinantes; Psicológicos

1. INTRODUCTION

The World Health Organization (WHO) declared that the SARS-CoV-2 infection became a pandemic on March 11th, 2020 (1).

Coronavirus Disease (COVID-19) pandemic has substantially influenced daily life for societies and individuals, affecting economics, health, and well-being. From the beginning of the pandemic caused by SARS-CoV-2, tremendous efforts have been made to tackle this public health emergency of international concern (2,3). The scientific community started working together on strategies to combat the virus, being the development of vaccines one of the most relevant.

In the past, the wide use of vaccines has decreased mortality and morbidity of different transmissible diseases (4). Therefore, an effective COVID-19 vaccination program, like other infectious diseases, has played and still plays an essential role in disease control, depending on a broad uptake, to decrease its incidence and prevalence (5,6).

However, universal vaccination is assumed to be difficult (1,7,8). Among other factors, some public safety concerns and scepticism are related with vaccine hesitance (9,10), although most of the approved COVID-19 vaccines have reported an efficacy greater than 90% in the phase 3 trials (11–13). Previous studies of influenza vaccination show that not all citizens took up the vaccines with a series of factors influencing its uptake (14). Also, approximately 5–10% of individuals have solid anti-vaccination beliefs, and a larger proportion could be viewed as being hesitant regarding vaccination in general (15,16). This led the WHO to declare vaccine hesitancy as one of the top challenges to world health (17). Vaccine hesitancy can vary between infectious diseases, and the rationale behind hesitancy for different diseases, such as influenza and COVID-19, can be influenced by a variety of factors, such as perception of disease severity, novelty of the disease, perceived vaccine effectiveness, political and ideological factors and misinformation and social media. Therefore, the SARS-CoV-2 virus has been perceived as more severe and life-threatening, particularly during the early stages of the pandemic. The high mortality rate in specific groups and the potential for long-term health consequences have generally increased vaccine acceptance. Despite the initial clinical trials for COVID-19 vaccines showing high effectiveness in preventing illness, particularly severe cases, COVID-19 is a novel coronavirus, and the vaccines developed to combat it are new, and concerns about vaccine safety and side effects remain. Additionally, the response to COVID-19 has become politicized in some countries, with vaccine hesitancy driven by political affiliations and ideologies. Finally, misinformation and conspiracy theories related to COVID-19 and its vaccines have been widely circulated on social media, leading to significant hesitancy in some communities (18,19).

A large number of studies continue to be published exploring the perception of the COVID-19 vaccination program and other determinants of its acceptance (20–22). The acceptance rate of COVID-19 vaccines has been investigated and reported in previous studies, varying across

countries globally, ranging from 43.4% to 94.3% (23–25). By bringing together relevant data on COVID-19 vaccine intention determinants, the underlying role of these factors could be clarified, which would eventually provide the opportunity for primary interventions targeting the groups of unvaccinated people, not only in COVID-19 but also in any other potential pandemic. Some data had already focused on predicting vaccination intention, using theoretical planned behaviours [26]. Other systematic reviews had highlighted acceptance of COVID-19 vaccination or demographic social and contextual factors that may influence the vaccination intention or hesitancy determinants [27-29]. Nevertheless, this systematic review aims to determine which socio-demographic and psychological determinants could be responsible for the adults' intention to be vaccinated against COVID-19 in European and North American adults. These two regions have political and ideological affinities, determining our choice for its inclusion in this review. In addition, Europe and the United States share trust in government and public health institutions, namely vaccination regulation, multiple approved vaccines, mass vaccination campaigns, and global collaboration (e.g., North America and Europe participated in global initiatives such as COVAX) (30,31).

2. METHODS

2.1 Design

A systematic literature review of quantitative studies was performed to summarize the available evidence on the social and psychological determinants of the intention to be vaccinated against COVID-19, in European and North American adults. The review was conducted taking into consideration the Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) (32), which encompass four phases: Identification, Screening, Eligibility, and Inclusion. The PROSPERO registration number is CRD42023402405. The protocol is also available at the Open Science Framework platform (DOI 10.17605/OSF.IO/QE7CD). A PEO (Population, Exposure, and Outcome) statement was developed by the reviewers' team as an aid to identify search terms and inclusion/exclusion criteria as appreciate for addressing this systematic review (*Table 1*). The definition of adults was provided by the National Institute of Health that affirms that adult is a person with an age between 18 and 65.

Table 1 - PEO criteria for inclusion and exclusion in systematic review.

2.2 Search strategy

Publications which describe original quantitative research were retrieved via electronic database searches of PubMed, Scopus, Web of Science, EBSCO, CINAHL, EMBASE and PsycINFO, in the period from January to February 2023. The research included articles published from January 2020 to December 2022. A combination of several terms was used, such as: COVID-19 vaccines, intention, refusal, hesitation, determinants and social. The keywords used with the Boolean operators AND and OR are presented in Table 2. The last date that the data was consulted was in April 2023.

Table 2 - Detailed description of the keywords used in each electronic database.

No restrictions regarding language or publication type (original articles or reviews) were applied in this phase. This search strategy led to a total of 126 PubMed papers, 313 Scopus papers, 131 Web of Science papers and six EBSCO papers.

A list of inclusion and exclusion criteria was defined, as detailed in *Table 3*.

Table 3 - Detailed description of the inclusion and exclusion criteria.

Firstly, the titles and abstracts of studies identified from the search were independently reviewed by two reviewers. Secondly, the full text of each study deemed possibly relevant was

retrieved and independently reviewed by two reviewers. Each reviewer compiled a list of studies that meet the inclusion criteria, in their view. The lists were compared, and any disagreements resolved by team discussion and consensus.

2.3 Quality Assessment of the Articles

Quality assessment was done using the JBI Critical Appraisal Checklist for Analytical Cross-Sectional Study (33). JBI is an international research organisation based in the Faculty of Health and Medical Sciences at the University of Adelaide, South Australia and accordingly to Ma et al. (2020) (33) this tool is the most used in quality assessment of analytical cross-sectional studies. Two reviewers evaluated independently all articles as "yes", "no", "unclear" and "not applicable" in the eight dimensions proposed on JBI tool. Disagreements were resolved by consensus and a third reviewer was available to arbitrate any issues that remained unresolved. After discussing the ratings and resolving any disagreement, the global rating for each of the selected articles was obtained by dividing the sum of ratings given ("No" =1; "Unclear" =2; "Yes" =3) for the number of dimensions. Those dimensions rated as "not applicable" were not considered, to permit a more adjusted score within studies. All the articles were assessed in the eight dimensions, despite only seven being scored (See Supplementary Material, S1). With the support of a tool called *robvis* (34) it was possible to build the quality assessment results in a creative way.

2.4 Data Extraction

Data was extracted from all articles independently by two reviewers. For each study, information was extracted regarding the authors, date of publication, country, goals, participants recruitment, and study design (including sampling and methodology).

3. RESULTS

3.1 Study selection

As it was previously described, it was used the PRISMA Framework, and the results are represented in the flow diagram PRISMA 2020 (Figure 1).

Figure 1 - Search and study selection PRISMA flow diagram.

Overall, none of the 28 studies were considered to be seriously flawed according to the quality assessment. The 28 included studies scored between 11 and 21 in the JBI Checklist (See Supplementary Material, S1: S1.1 and S1.2). All studies were considered to have a low risk of bias for selection (Figure 2).

Figure 2 - Summary of the Critical Appraisal based on JBI checklist (risk of bias).

3.2 Characteristics of included studies

The 28 selected studies involved a total of 138,487 subjects (See Supplementary Material, S2), and were conducted from February 2020 to December 2022, half of them (n=14) taking place in 2021 (18,35–44). Considering the country of origin, nine (32.1%) studies were carried in the United States of America (35,40,41,43–48), five (17.9%) in the United Kingdom (18,25,42,49,50), even though two of them also included participants from the Republic of Ireland (42,50), three (10.7%) studies were conducted in Germany (38,52,53), two (7.1%) studies in multiple countries (35,49), and the remaining studies (32.2%) took place in Turkey (54), Finland (38), Northern Ireland and Republic of Ireland (36), Netherlands (55), Federation of Bosnia and Herzegovina (56), Switzerland (57), Czech Republic (58), Italy (59) and Belgium (60).

All articles adopted a cross-sectional design. Twenty studies used a non-probabilistic sampling method, namely: quota sampling (25,35,38,40,42–44,49,52,56), convenience sampling

(37,40,47,50,57,60) or snowball sampling (36,54,59). Four studies were conducted using a stratified sample (18,41,53,61), two of them randomly (53,61). For the other five studies, sampling methodology was not described in detail.

Participant recruitment was done essentially online (18,35,36,38,39,43,50–52,56–61), including social media (37,40,54), e-mail (47) or using online panel providers (42–44,49,53). On the Latkin et al. (2021) study, an online panel provider was combined with a telephone contact. For the other two studies, the way participants were recruited was not described in detail (41).

Being intention to be vaccinated against COVID-19 the focus, there are some differences in how this variable was measured in the studies under analysis. Despite most of them measure intention to be vaccinated, in four studies (38,47,53,58) the focus was on acceptability or acceptance of a COVID-19 vaccine and in eight other studies (36,40–43,48,60,61) it was vaccine hesitancy that was measured. For the studies focusing on psychological determinants, 13 of them were theory-based (18,25,35,36,39,41,44,45,50–52,57,60), which was reflected in the questions posed in the questionnaires being used. In the study conducted by Sanders et al. (2021), although it was not theory-based, the beliefs related to vaccination intention, under assessment, are the result of an extensive literature review (55).

3.3 Socio-Demographic Determinants

The full version of the results regarding socio-demographic data and vaccination intention are presented in the Supplementary Material, S3.

Vaccination intention ranged from 45.4% (18) to 89% (58), with an average of 65%.

Regarding socio-demographic determinants, not all were evaluated in every study. Age was addressed in 21 studies (18,38–45,47,48,50,51,53–56,58,61), although six of them (18,38,45,48,54,58) were not considered statistically significant. Apart from the study conducted by Khubchandani et al (2021), vaccination intention was higher in older participants (40).

Only three studies didn't address gender differences (35,52,60). In ten studies (37–39,41,42,44,47,49,51,54), intention to be vaccinated was higher in men and in five studies (43,50,56,61), the opposite. In the remaining studies, gender was found to be a non-statistically significant variable.

Within the studies addressing educational level as a determinant of vaccination intention, only in the studies conducted by Musa et al. (2022) and by Lazarus et al. (2020) the intention to be vaccinated was higher in participants with a lower educational level (namely in Canada, Spain and the United Kingdom) (53,56). In the remaining countries and in all the other studies addressing this determinant (40,41,44,47,49,51,57,59,61) vaccination intention was higher in participants with a higher educational level.

The influence of marital status on the intention to be vaccinated was not addressed in 19 studies. Participants that were married / civil united or living in a cohabitation demonstrated higher vaccination intention in four studies (41,44,47,51), while in two studies (49,54) that was higher for single / separated participants. In the remaining studies, no differences were found concerning marital status.

Finally, in the studies where ethnicity was found to be a determinant for vaccination intention (35,40,41,43,44,47–50,61), white (Caucasian) participants always reflected a higher vaccination intention.

3.4 Psychological Determinants

Of the 28 selected articles, 23 focused on psychological determinants. The identified factors were classified in eleven thematic categories, namely: Trust on Media, Social & Scientific Entities (addressed in 52% of the 23 studies); Disease Severity (57%), Vaccine Costs & Risks (52%), Vaccine Efficacy (35%), Social Concerns (30%), Conspiracy Beliefs (27%), Perceived of Risk to be Infected (30%), Control Perception (26%), Social Norms (30%), Personal Concerns

(22%) and Emotions & Personality (22%). To enable a more substantiated analysis, categories that were addressed only in one study were excluded (e.g., Perceived health status).

- Disease Severity and Perceived Risk to be Infected

Of the 13 studies that addressed Disease Severity (Table 4), 10 studies (76%) fully confirmed that people with a higher perception of COVID-19 severity tended to have a stronger intention to be vaccinated. Only one study found no significant relation with vaccine intention (52).

Sherman et al. (2021) distinguishes between perceived severity of COVID-19 for the individual and for the community, with only the former being a predictor of vaccination intention. .

Regarding Perceived Risk to be Infected (table 4), six studies assessed risk and concern of being infected and one focused the perception of individual health risk factors for COVID-19 (51).

Five studies (71%) confirm that intention to be vaccinated is directly related to a higher perception of risk to be infected (18,39,40,47,50–52). One study found mixed results on different national samples.

Table 4 - List of Beliefs: Disease Severity and Perceived Risk to be Infected.

- Vaccine Costs & Risks, Control Perception and Vaccine Efficacy

Twelve studies focused on beliefs about vaccine costs and risks (Table 5). With exception of Musa et al. (2022) (56), all the studies (92%) confirmed that a strong perception of vaccine costs and risks is related with a low intention to be vaccinated.

Table 5 shows that three of six studies focusing control perception (i.e., to be able to obtain the vaccine, when/if available) found no significant relation with vaccination intention (18,45,52). Bleakley et al. (2022) (35) presented heterogeneous results concerning this relation, varying according to age and ethnicity. Only two studies (33%) found that intention to vaccinate is related with the perception of easy access (36,56).

Five of eight studies (62%) presented in Table 5 support that people who perceive high vaccine effectiveness/efficacy (e.g.,: the perception/belief in the positive effect of the vaccine in preventing COVID-19 or reducing its symptoms) tend to have a stronger intention to be vaccinated (18,36,47,49,50). Two articles found this relation but only in specific ethnic and age groups (36,38). The remaining two articles do not confirm this association.

Table 5 - List of Beliefs: Vaccine Costs & Risks, Cost perception and Vaccine efficacy.

- Personal Concerns, Social Concerns, Social Norms and Emotions and Personality

Three (18,38,50) of the five studies (60%) confirm the relevance of personal concerns (e.g., protection of oneself; avoid future regret for not having been vaccinated) in the development of vaccination intention (Table 6).

With respect to social concerns (Table 6), seven articles (39,42,45,49,50,55,59) confirm that vaccination intention is related with social concerns (86%): altruistic motivations (42,59), collective importance of COVID-19 vaccine (49), civic responsibility (50), the perception of community benefits (39,45) and moral duty to be vaccinated (50,55). Only one study does not confirm this conclusion (39).

As shown in Table 6, influence of social pressure on vaccine intention was confirmed by four (66,7%) studies (18,36,50,52). Bleakley et al. (2022) (35) found a significant relation between normative pressure and vaccine intention but only for people over 50 years old.

Table 6 - List of Beliefs: Personal Concerns, Social Concerns and Social Norms

- Trust on media, social & scientific Entities; Conspiracy beliefs; Emotions & personality

Table 7 shows the conclusions of the nineteen studies that addressed the relation between vaccine intention and a) trust on media, social and scientific entities, b) conspiracy beliefs and c) emotions & personality (18,37–39,48,50–52,56,59,60). From the eleven studies that assess specifically confidence in political institutions and health authorities seven confirm the relation of this variable with vaccine intention (18,37,43,50,51,56,60). Two do not confirm this relation (25,52) and the remaining two present heterogeneous results depending on age (38) and country (39). The impact of trust in health professionals on willingness to vaccinate is specifically assessed in three studies with heterogeneous results (38,39,52). Two studies do not confirm that trust in science is relevant in the decision to take the vaccine (48,52).

Concerning media information, El-Far Cardo et al. (2021) concluded that those who use media as a source of information have a stronger intention to vaccinate which was not confirmed in Stoler et al. (2022) study (48). Musa et al. (2022) found that people who think that COVID-19 information is exaggerated by the media tend to have a low vaccination intention (56).

In four studies the confidence is assessed combining two or more entities (science, health and political institutions, health professionals) (42,55,56,59) with three of them confirming a direct relation with the intention to be vaccinated (39,52,56). Overall, the relation with trust on media, social & scientific entities is fully confirmed in 10 studies (62,5%).

The results concerning adherence to conspiracy beliefs (Table 7) show heterogeneous conclusions, with four studies (57%) concluding that they are predictors of the unwillingness to be vaccinated (44,49,57,60), while two studies do not confirm this idea (48,52) and another presents mixed results (42)

Emotions as a predictor of vaccine intention are analyzed in four of six studies (Table 7) (51,54,56,57). Though Musa et al. (2022) (56) conclude that negative affective states are predictors of vaccine intention, Roberts et al. (2022) (43) do not confirm this relation for depression and stress. Embitterment and life satisfaction seems to be positively related with the intention to be vaccinated (51,57). The results of the one study that focus on personality variables do not confirm their influence on vaccine intention (48).

From a more global analysis, the most widely confirmed determinants categories of vaccine intention are Vaccine costs & risks (92%), Social concerns (86%) Disease severity (76%) and Perceived risk to be infected (71%). The remaining determinants are confirmed in around 50% of the studies, with exception for Control perception (33%) and Emotions & personality.

Table 7 - List of Beliefs: Trust on Media, Social & Scientific Entities; Conspiracy Beliefs; Emotion & Personality

4. DISCUSSION

Vaccines are essential to accelerate the immunization process (62). However, the COVID-19 vaccines acceptance is influenced by psychosocial variables. The present study aims to systematize the socio-demographic and psychological determinants of the adult intention to be vaccinated against COVID-19 in European and North American adults.

Considering the general characteristics of the 28 selected studies, some methodological and theoretical shortcomings can be identified; first, most studies use non-probabilistic sampling, and second, only half of the studies focusing on psychological determinants are theory-based. The selection of psychological determinants under study without a theoretical framework is even more serious given a substantive number of consolidated theories explaining health behaviors (e.g., Health Beliefs Model, Theory of Planned Behavior; Transtheoretical Model), which have already been widely used in studies related to vaccination adherence.

Cognitive factors related to vaccine intention can be affected over time. For instance, risk perception and the perception of vaccine efficacy can be influenced by the evolution of the number of deaths and knowledge about the disease/vaccine, among other aspects that change over time. Cognitive factors can be also affected by the periods of lockdown and access to vaccination. However, this time variables were not analysed in the present

study due to the absence of international data on dates of lockdown and vaccine access as well as the rate of vaccination for all European and North American countries.

Figure 3 summarizes the results described below, dividing the sociodemographic and psychological determinants along 2 axes: the number of articles found for each determinant and, among these, the percentage of articles fully confirming the relationship between the determinant and the intention to vaccinate.

Figure 3 - Determinants of COVID-19 Vaccine intention by number of articles and percentage of confirmation

4.1 SOCIO-DEMOGRAPHIC VARIABLES

In the present systematic review study, it was observed in 35% of the studies a positive relation between older adults and their intention to vaccinate (42–44,47,49), which may be attributed to the fact that the mortality rate is higher in the elderly population, and consequently there is a perception of greater fragility of this group when facing this disease, leading to a greater need to be vaccinated. These results are corroborated by other authors (6,22,63–66).

Regarding gender, a high prevalence of vaccination intention was found in men (9 of the assessed studies) (37,38,41,42,44,47,49,54). These results are consistent with many studies that found greater acceptance in men, as well as in individuals with higher educational levels (22,63–67).

Concerning marital status, those who are married or in a union were the ones who showed the highest vaccination intention, as found in other studies (22,63–66).

As for race/ethnicity, there was a vast diversity of classification, varying according to the country or region where the study was conducted. Therefore, it is challenging to extract relevant conclusions, except for the higher intention to be vaccinated in white (Caucasian) people. In Cascini's, Fisk's, and Wang's studies (64,68,69), minorities of ethnic groups have lower levels of acceptance (64,68–70). Nevertheless, the study developed by Szilagyi et al. (2021) (67) only showed lower levels of acceptance for Black race than White, Hispanic and Asian as (67). The results of these studies indicate, as pointed out by Fisk (2021) (68) that awareness of vaccination should be strengthened among ethnic minority leaders.

4.2 PSYCHOLOGICAL VARIABLES

One of the aims of this study was to determine which psychological variables could be responsible for the intention to be vaccinated against COVID-19. With exception of emotion and personality, focused only in five studies, the main psychological variables were beliefs that were grouped into seven categories: Disease Severity, Perceived Risk of being Infected, Vaccine Costs and Risks, Vaccine efficacy, Social concerns, Conspiracy Beliefs, and Trust in Social Scientific Entities. A relevant number of studies found a significant relationship between vaccination intention and each category, with which underlies their importance in accepting the COVID-19 vaccine.

All studies confirm an adverse effect of conspiracy beliefs on vaccination intention. These findings are corroborated by Wang's systematic review that, despite just including United States studies, emphasized the association between COVID-19-related conspiracy theories to vaccine acceptance (69,70). Trust in science and health institutions/professionals also emerges as a relevant factor since most studies found a positive relationship between those beliefs and vaccination intention (25,42,71,72). This confirms Nindrea's results (2021) (66) that highlight the relevance of trust in the health system to vaccine acceptance (66), and Joshi's study (2021) that identifies valuing doctor's recommendations and government satisfaction as associated factors of vaccine acceptance (65). These results point out that individual perceptions of social phenomena influence vaccination intention and are not directly related to personal health.

Regarding the perceptions about the vaccine, beliefs about adverse effects are essential determinants of vaccination hesitation in all the studies addressing this relation. The side-effects was one of the vaccine risks identified as predictors of vaccine hesitancy (25,49,55). In addition, results confirmed that the perception of vaccine efficacy is related to the intention to be vaccinated (47,49), being corroborated by Lin's study (73). This outcome confirms Joshi's scoping review finding that vaccine efficacy is related to a high perception of benefits and a low perception of barriers, which were associated with higher vaccination acceptance (Joshi et al., 2021). Cassini's systematic review (64) also highlighted vaccine safety and efficacy perceptions, as well as the study developed by Fernandes et al., (2021) (74). Finally, Dimasa's systematic review (63) identified individual susceptibility to COVID-19 as one of the main determinants of vaccine acceptance. That confirms the assertion in

many psychological models on health behaviour (75–77) that behaviour adopting results from balancing its benefits and disadvantages.

In our systematic review, the studies addressing the risk perception of COVID-19 (severity of the disease; risk to be infected) present the most heterogeneous results since around one-third of the studies do not confirm that intention to be vaccinated is related to the risk perceived of COVID-19. However, regarding previous studies, Nindrea's systematic review found that fear of COVID-19 and perceived risks were positively correlated with vaccine acceptance (66). Similarly, Joshi's scoping review observed that a higher perceived risk of COVID-19 infection was associated with vaccine acceptance (65). The results interpretation must consider that half of the included studies were held in 2020 when many people worldwide had no access to clear information about the risks of the disease.

4.3 LIMITATIONS

Several limitations were identified during the development of this systematic review and should be taking into consideration: 1) different criteria were used to classify some socio-demographic variables, such as race or ethnicity; 2) several studies were developed in multiple contexts, social, cultural, political, economic, etc., differences may have an impact on the results, which increases the relevance of complementary studies at national or continental level. However, the most critical limitation identified by the authors was that 4) several studies were carried out in different periods of the lockdown and phases of the vaccine studies/clinical trials. This is one of the biggest issues regarding the COVID-19' vaccination studies were the timing of the surveys, as highlighted in the review of 209 studies carried out by Cascini and colleagues (64).

To overcome the limitations identified in future research on vaccination hesitancy, particularly given its status as a major challenge to global health, several approaches could be implemented, namely: Standardization of Socio-Demographic Criteria; Context-Specific Research; Longitudinal Studies; Meta-Analyses Across Timeframes; Use of Advanced Analytical Tools; Global Collaboration and Data Sharing; and Survey Timing Considerations.

One limitation in current studies is the use of different criteria to classify socio-demographic variables like race, ethnicity, and other factors. To address this, future research can establish standardized frameworks for classifying and reporting these variables. International collaborations among researchers, healthcare organizations, and governments can help create consensus-based guidelines for consistent data collection. This will make it easier to compare and synthesize findings across different studies and contexts.

Moreover, given the influence of social, cultural, political, and economic factors on vaccination hesitancy, future studies should focus on tailoring research to specific regions or populations. Conducting more localized or national-level studies can better reflect the unique challenges within different populations. This approach will also help identify context-specific drivers of hesitancy, leading to more effective, targeted interventions.

Although deeply expensive, longitudinal studies that track vaccine hesitancy and uptake over time—before, during, and after critical events like lockdowns or vaccine rollouts—would provide more comprehensive data. These studies would allow researchers to better understand how attitudes shift based on changing conditions, including evolving pandemic dynamics or the introduction of new vaccines. To address the issue of varying timelines in existing research, future systematic reviews could conduct meta-analyses that account for the period of the studies. By organizing results according to different stages of the pandemic or vaccine development, researchers can better identify temporal patterns in hesitancy.

Several research points that the future is in the use of advanced analytical tools, such as machine learning models, can help future researchers analyse complex, multi-contextual datasets. These tools can identify patterns across socio-demographic variables and contextual factors (social, political, etc.) in a way that might be missed in more traditional analyses.

Plus, enhancing collaboration between countries and institutions can facilitate the sharing of data, which may help overcome some of the variability caused by different contexts. Creating global databases that allow for cross-contextual research will help harmonize findings and better identify universal trends and challenges in vaccine hesitancy.

Finally, when planning future studies, researchers should design surveys and data collection methods that can account for the timing of significant events, such as lockdowns or new vaccine introductions. Timing surveys more strategically, or conducting them continuously throughout vaccine rollouts, will provide more accurate snapshots of public sentiment.

5. CONCLUSION

This systematic review showed the socio-demographic and psychological determinants that contributed to the intention of adults being vaccinated against COVID-19 in Europe and North America.

There is, in general, a good acceptance of COVID-19 vaccines in these two regions, with wide discrepancies across the identified determinants, such as age; gender; educational level; marital status; race/ethnicity are found. Regarding the psychological determinants identified in this systematic review, the perception of COVID-19 severity for the population and personnel and the perceived risk of being infected should be highlighted. Several included studies also underlined the significance of several beliefs regarding the COVID-19 vaccines, such as lack of confidence, concerns about potential harm, safety, and side effects. Other social concerns and conspiracy beliefs were also linked to being vaccinated.

The findings of this study may help European and North American health institutions and governments to identify the best approach to implement COVID-19 vaccination programs. Moreover, this could contribute to the fight against vaccination hesitancy to promote herd immunity around these countries.

6. ACKNOWLEDGEMENTS

H&TRC authors gratefully acknowledge the FCT/MCTES national support through the UIDB/05608/2020 and UIDP/05608/2020. This work was also supported by the project “Youth Breakdown in the post-COVID era and their Vaccination Intention” (IPL/2021/Vaccin2You(th)_ESTeSL), funded by Instituto Politécnico de Lisboa, Portugal.

7. STATEMENT OF ETHICS

Ethical approval and participant consent are not applicable in this case, as it is a systematic review.

8. CONFLICT OF INTEREST STATEMENT

The authors have no conflicts of interest to declare.

9. FUNDING SOURCES

Nothing to declare.

10. DATA AVAILABLE STATEMENT

All data generated or analysed during this study are included in this. Further enquiries can be directed to the corresponding author.

11. REFERENCES

1. World Health Organization. WHO Coronavirus (COVID-19) Dashboard. 2022.
2. Hanney SR, Wooding S, Sussex J, Grant J. From COVID-19 research to vaccine application: why might it take 17 months not 17 years and what are the wider lessons? *Health Res Policy Syst.* 2020 Dec 8;18(1):61.
3. Lurie N, Saville M, Hatchett R, Halton J. Developing Covid-19 Vaccines at Pandemic Speed. *New England Journal of Medicine.* 2020 May 21;382(21):1969–73.
4. Greenwood B. The contribution of vaccination to global health: past, present and future. *Philosophical Transactions of the Royal Society B: Biological Sciences.* 2014 Jun 19;369(1645):20130433.
5. Orenstein, WA, Bernier, RH, Dondero TJ, Hinman, AR, Marks JS, Bart, KJ, et al. Field evaluation of vaccine efficacy*. Vol. 63, *Bulletin of the World Health Organization.* 1985.
6. Weinberg GA, Szilagyi PG. Vaccine Epidemiology: Efficacy, Effectiveness, and the Translational Research Roadmap. *J Infect Dis.* 2010 Jun;201(11):1607–10.
7. Hardwick RJ, Truscott JE, Oswald WE, Werkman M, Halliday KE, Pullan RL, et al. Dynamics of individual adherence to mass drug administration in a conditional probability model. Available from: <https://doi.org/10.1101/2020.04.17.20069476>
8. Levine OS. What drivers will influence global immunizations in the era of grand convergence in global health? *Vaccine.* 2017 Jan;35:A6–9.
9. Betsch C, Renkewitz F, Haase N. Effect of Narrative Reports about Vaccine Adverse Events and Bias-Awareness Disclaimers on Vaccine Decisions. *Medical Decision Making.* 2013 Jan 8;33(1):14–25.
10. Bhattacharyya S, Bauch CT. “Wait and see” vaccinating behaviour during a pandemic: A game theoretic analysis. *Vaccine.* 2011 Jul;29(33):5519–25.

11. Chagla Z. The BNT162b2 (BioNTech/Pfizer) vaccine had 95% efficacy against COVID-19 ≥ 7 days after the 2nd dose. *Ann Intern Med.* 2021 Feb;174(2):JC15.
12. Polack FP, Thomas SJ, Kitchin N, Absalon J, Gurtman A, Lockhart S, et al. Safety and Efficacy of the BNT162b2 mRNA Covid-19 Vaccine. *New England Journal of Medicine.* 2020;383(27):2603–15.
13. Voysey M, Clemens SAC, Madhi SA, Weckx LY, Folegatti PM, Aley PK, et al. Safety and efficacy of the ChAdOx1 nCoV-19 vaccine (AZD1222) against SARS-CoV-2: an interim analysis of four randomised controlled trials in Brazil, South Africa, and the UK. *The Lancet.* 2021;397(10269):99–111.
14. Bish A, Yardley L, Nicoll A, Michie S. Factors associated with uptake of vaccination against pandemic influenza: A systematic review. *Vaccine.* 2011 Sep;29(38):6472–84.
15. Lane S, MacDonald NE, Marti M, Dumolard L. Vaccine hesitancy around the globe: Analysis of three years of WHO/UNICEF Joint Reporting Form data-2015–2017. *Vaccine.* 2018 Jun;36(26):3861–7.
16. Dror AA, Eisenbach N, Taiber S, Morozov NG, Mizrahi M, Zigron A, et al. Vaccine hesitancy: the next challenge in the fight against COVID-19. *Eur J Epidemiol.* 2020 Aug 12;35(8):775–9.
17. World Health Organization. Ten threats to global health in 2019 [Internet]. Accessed at January 2022. Available from: <https://www.who.int/news-room/spotlight/ten-threats-to-global-health-in-2019>.
18. Goffe L, Antonopoulou V, Meyer CJ, Graham F, Tang MY, Lecouturier J, et al. Factors associated with vaccine intention in adults living in England who either did not want or had not yet decided to be vaccinated against COVID-19. *Hum Vaccin Immunother.* 2021;17(12):5242–54.
19. Dombrádi V, Joó T, Palla G, Pollner P, Belicza É. Comparison of hesitancy between COVID-19 and seasonal influenza vaccinations within the general Hungarian population: a cross-sectional study. *BMC Public Health* [Internet]. 2021;21(1):2317. Available from: <https://doi.org/10.1186/s12889-021-12386-0>
20. Daly M, Robinson E. Willingness to vaccinate against COVID-19 in the US: Longitudinal evidence from a nationally representative sample of adults from April-October 2020. *medRxiv* [Internet]. 2020 Nov 30; Available from: <http://www.ncbi.nlm.nih.gov/pubmed/33269354>
21. Kreps S, Prasad S, Brownstein JS, Hswen Y, Garibaldi BT, Zhang B, et al. Factors Associated With US Adults' Likelihood of Accepting COVID-19 Vaccination. *JAMA Netw Open.* 2020 Oct 20;3(10):e2025594.
22. Malik AA, McFadden SM, Elharake J, Omer SB. Determinants of COVID-19 vaccine acceptance in the US. *EClinicalMedicine.* 2020 Sep;26:100495.
23. Sallam M, Dababseh D, Eid H, Hasan H, Taim D, Al-Mahzoum K, et al. Low covid-19 vaccine acceptance is correlated with conspiracy beliefs among university students in Jordan. *Int J Environ Res Public Health.* 2021 Mar 2;18(5):1–14.
24. Sallam M, Dababseh D, Eid H, Al-Mahzoum K, Al-Haidar A, Taim D, et al. High rates of covid-19 vaccine hesitancy and its association with conspiracy beliefs: A study in Jordan and Kuwait among other Arab countries. *Vaccines (Basel).* 2021 Jan 1;9(1):1–16.
25. Sherman SM, Smith LE, Sim J, Amlôt R, Cutts M, Dasch H, et al. COVID-19 vaccination intention in the UK: results from the COVID-19 vaccination acceptability study (CoVAccS), a nationally representative cross-sectional survey. *Hum Vaccin Immunother.* 2021;17(6):1612–21.
26. Limbu YB, Gautam RK, Zhou W. Predicting Vaccination Intention against COVID-19 Using Theory of Planned Behavior: A Systematic Review and Meta-Analysis. Vol. 10, *Vaccines.* MDPI; 2022.
27. Alarcón-Braga EA, Hernandez-Bustamante EA, Salazar-Valdivia FE, Valdez-Cornejo VA, Mosquera-Rojas MD, Ulloque-Badaracco JR, et al. Acceptance towards COVID-19 vaccination in Latin America and the Caribbean: a systematic review and meta-analysis. *Travel Med Infect Dis.* (2022) 49:102369. doi: 10.1016/j.tmaid.2022.102369
28. AlShurman BA, Khan AF, Mac C, Majeed M, Butt ZA. What demographic, social, and contextual factors influence the intention to use COVID-19 vaccines: a scoping review. *Int J Environ Res Public Health.* (2021) 18:9342. doi: 10.3390/ijerph18179342.
29. Biswas MdR, Alzubaidi MS, Shah U, Abd-Alrazaq AA, Shah Z. A scoping review to find out worldwide COVID-19 vaccine hesitancy and its underlying determinants. *Vaccines.* (2021) 9:1243. doi: 10.3390/vaccines9111243)
30. Sachs JD, Karim SSA, Akinin L, Allen J, Brosbøl K, Colombo F, et al. The Lancet Commission on lessons for the future from the COVID-19 pandemic. Vol. 400, *The Lancet.* Elsevier B.V.; 2022. p. 1224–80.
31. Hutton B, Salanti G, Caldwell DM, Chaimani A, Schmid CH, Cameron C, et al. The PRISMA Extension Statement for Reporting of Systematic Reviews Incorporating Network Meta-analyses of Health Care Interventions: Checklist and Explanations. *Ann Intern Med.* 2015 Jun 2;162(11):777–84.

32. Moola S, Munn Z, Tufanaru C, Aromataris E, Sears K, Sfec R, et al. Chapter 7: Systematic Reviews of Etiology and Risk. In: JBI Manual for Evidence Synthesis. JBI; 2020.
33. Ma LL, Wang YY, Yang ZH, Huang D, Weng H, Zeng XT. Methodological quality (risk of bias) assessment tools for primary and secondary medical studies: What are they and which is better? Vol. 7, Military Medical Research. BioMed Central Ltd.; 2020.
34. McGuinness LA, Higgins JPT. Risk-of-bias VISualization (robvis): An R package and Shiny web app for visualizing risk-of-bias assessments. *Res Synth Methods*. 2021 Jan 6;12(1):55–61.
35. Bleakley A, Hennessy M, Maloney E, Young DG, Crowley J, Silk K, et al. Psychosocial Determinants of COVID-19 Vaccination Intention Among White, Black, and Hispanic Adults in the US. *Annals of Behavioral Medicine*. 2022 Apr 1;56(4):347–56.
36. Breslin G, Dempster M, Berry E, Cavanagh M, Armstrong NC. COVID-19 vaccine uptake and hesitancy survey in Northern Ireland and Republic of Ireland: Applying the theory of planned behaviour. *PLoS One*. 2021 Nov 1;16.
37. El-Far Cardo A, Kraus T, Kaifie A. Factors that shape people's attitudes towards the covid-19 pandemic in germany—the influence of media, politics and personal characteristics. *Int J Environ Res Public Health*. 2021 Aug 1;18(15).
38. Hammer CC, Cristea V, Dub T, Sivelä J. High but slightly declining COVID-19 vaccine acceptance and reasons for vaccine acceptance, Finland April to December 2020. *Epidemiol Infect*. 2021.
39. Kerr JR, Schneider CR, Recchia G, Dryhurst S, Sahlin U, Dufouil C, et al. Correlates of intended COVID-19 vaccine acceptance across time and countries: Results from a series of cross-sectional surveys. *BMJ Open*. 2021 Aug 2;11(8).
40. Khubchandani J, Sharma S, Price JH, Wiblishauser MJ, Sharma M, Webb FJ. COVID-19 Vaccination Hesitancy in the United States: A Rapid National Assessment. *J Community Health*. 2021 Apr 1;46(2):270–7.
41. Latkin CA, Dayton L, Yi G, Colon B, Kong X. Mask usage, social distancing, racial, and gender correlates of COVID-19 vaccine intentions among adults in the US. *PLoS One*. 2021 Feb 1;16.
42. Murphy J, Vallières F, Bentall RP, Shevlin M, McBride O, Hartman TK, et al. Psychological characteristics associated with COVID-19 vaccine hesitancy and resistance in Ireland and the United Kingdom. *Nat Commun*. 2021 Dec 1;12(1).
43. Roberts HA, Clark DA, Kalina C, Sherman C, Brislin S, Heitzeg MM, et al. To vax or not to vax: Predictors of anti-vax attitudes and COVID-19 vaccine hesitancy prior to widespread vaccine availability. *PLoS One*. 2022 Feb 1;17.
44. Ruiz JB, Bell RA. Predictors of intention to vaccinate against COVID-19: Results of a nationwide survey. *Vaccine*. 2021 Feb 12;39(7):1080–6.
45. Hayashi Y, Romanowich P, Hantula DA. Predicting Intention to Take a COVID-19 Vaccine in the United States: Application and Extension of Theory of Planned Behavior. *American Journal of Health Promotion*. 2022 May 1;36(4):710–3.
46. Piltch-Loeb R, Silver DR, Kim Y, Norris H, McNeill E, Abramson DM. Determinants of the COVID-19 vaccine hesitancy spectrum. *PLoS One*. 2022 Jun 1;17.
47. Reiter PL, Pennell ML, Katz ML. Acceptability of a COVID-19 vaccine among adults in the United States: How many people would get vaccinated? *Vaccine*. 2020 Sep 29;38(42):6500–7.
48. Stoler J, Klofstad CA, Enders AM, Uscinski JE. Sociopolitical and psychological correlates of COVID-19 vaccine hesitancy in the United States during summer 2021. *Soc Sci Med*. 2022 Aug 1;306.
49. Freeman D, Loe BS, Chadwick A, Vaccari C, Waite F, Rosebrock L, et al. COVID-19 vaccine hesitancy in the UK: The Oxford coronavirus explanations, attitudes, and narratives survey (Oceans) II. *Psychol Med*. 2021.
50. Walsh JC, Comar M, Folan J, Williams S, Kola-Palmer S. The psychological and behavioural correlates of COVID-19 vaccine hesitancy and resistance in Ireland and the UK. *Acta Psychol (Amst)*. 2022 May 1;225.
51. Fobiwe JP, Martus P, Poole BD, Jensen JL, Joos S. Influences on Attitudes Regarding COVID-19 Vaccination in Germany. *Vaccines (Basel)*. 2022 May 1;10(5).
52. Seddig D, Maskileyson D, Davidov E, Ajzen I, Schmidt P. Correlates of COVID-19 vaccination intentions: Attitudes, institutional trust, fear, conspiracy beliefs, and vaccine skepticism. *Soc Sci Med*. 2022 Jun 1;302.
53. Lazarus J V., Wyka K, Rauh L, Rabin K, Ratzan S, Gostin LO, et al. Hesitant or Not? The Association of Age, Gender, and Education with Potential Acceptance of a COVID-19 Vaccine: A Country-level Analysis. *J Health Commun*. 2020;25(10):799–807.

54. Akarsu B, Canbay Özdemir D, Ayhan Baser D, Aksoy H, Fidancı İ, Cankurtaran M. While studies on COVID-19 vaccine is ongoing, the public's thoughts and attitudes to the future COVID-19 vaccine. *Int J Clin Pract*. 2021 Apr 1;75(4).
55. Sanders JG, Dijk M van, Elberse J, Lambooi MS, Kroese FM, Bruin M de. Understanding a national increase in COVID-19 vaccination intention, the Netherlands, November 2020–March 2021. *Vol. 26, Euro Surveill*. 2021.
56. Musa S, Cilovic-Lagarija S, Kavazovic A, Bosankic-Cmajcanin N, Stefanelli A, Scott NA, et al. COVID-19 Risk Perception, Trust in Institutions and Negative Affect Drive Positive COVID-19 Vaccine Intentions. *Int J Public Health*. 2022 Apr 11;67.
57. Koroma D, Pestalozzi MI, Znoj H. How Social Exclusion, Embitterment, and Conspiracy Beliefs Mediate Individual's Intention to Vaccination against COVID-19: Results from a Moderated Serial Mediation Analysis. *Psychopathology*. 2022 Mar 1;55(2):93–103.
58. Blanka K, Vladimir P, Jan MH, Ladislav S. Vaccination against COVID-19 as prevention of occupational disease in University of Defence members in the Czech Republic - motivation for vaccination and reasons for hesitancy. *Bratislava Medical Journal*. 2022;123(10):745–51.
59. Caserotti M, Girardi P, Tasso A, Rubaltelli E, Lotto L, Gavaruzzi T. Joint analysis of the intention to vaccinate and to use contact tracing app during the COVID-19 pandemic. *Sci Rep*. 2022 Dec 1;12(1).
60. Van Oost P, Yzerbyt V, Schmitz M, Vansteenkiste M, Luminet O, Morbée S, et al. The relation between conspiracism, government trust, and COVID-19 vaccination intentions: The key role of motivation. *Soc Sci Med*. 2022 May 1;301.
61. Piltch-Loeb R, Su M, Bonetti M, Testa M, Stanton E, Toffolutti V, et al. Cross-National Vaccine Concerns and Predictors of Vaccine Hesitancy in Not-Fully Vaccinated Individuals: Findings from USA, Canada, Sweden, and Italy. *Vaccines (Basel)*. 2022 Oct 1;10(10).
62. Hall V, Foulkes S, Insalata F, Kirwan P, Saei A, Atti A, et al. Protection against SARS-CoV-2 after Covid-19 Vaccination and Previous Infection. *New England Journal of Medicine*. 2022;386(13):1207–20.
63. Akem Dimala C, Kadia BM, Nguyen H, Donato A. Community and Provider Acceptability of the COVID-19 Vaccine: A Systematic Review and Meta-analysis. *Advances in Clinical Medical Research and Healthcare Delivery*. 2021 Nov 12;1(3).
64. Cascini F, Pantovic A, Al-Ajlouni Y, Failla G, Ricciardi W. Attitudes, acceptance and hesitancy among the general population worldwide to receive the COVID-19 vaccines and their contributing factors: A systematic review. *EClinicalMedicine*. 2021 Oct 1;40.
65. Joshi A, Kaur M, Kaur R, Grover A, Nash D, El-Mohandes A. Predictors of COVID-19 Vaccine Acceptance, Intention, and Hesitancy: A Scoping Review. *Vol. 9, Frontiers in Public Health*. Frontiers Media S.A.; 2021.
66. Nindrea RD, Usman E, Katar Y, Sari NP. Acceptance of COVID-19 vaccination and correlated variables among global populations: A systematic review and meta-analysis. *Clin Epidemiol Glob Health*. 2021 Oct 1;12.
67. Szilagyi PG, Thomas K, Shah MD, Vizueta N, Cui Y, Vangala S, et al. National Trends in the US Public's Likelihood of Getting a COVID-19 Vaccine - April 1 to December 8, 2020. *Vol. 325, JAMA - Journal of the American Medical Association*. American Medical Association; 2021. p. 396–8.
68. Fisk RJ. Barriers to vaccination for coronavirus disease 2019 (COVID-19) control: experience from the United States. *Vol. 5, Global Health Journal*. KeAi Communications Co.; 2021. p. 51–5.
69. Wang J, Lu X, Lai X, Lyu Y, Zhang H, Fenghuang Y, et al. The changing acceptance of COVID-19 vaccination in different epidemic phases in China: A longitudinal study. *Vaccines (Basel)*. 2021 Mar 1;9(3):1–17.
70. Wang K, Wong ELY, Ho KF, Cheung AWL, Yau PSY, Dong D, et al. Change of willingness to accept covid-19 vaccine and reasons of vaccine hesitancy of working people at different waves of local epidemic in hong kong, china: Repeated cross-sectional surveys. *Vaccines (Basel)*. 2021 Jan 1;9(1):1–15.
71. Al-Mohaithef M, Padhi BK. Determinants of covid-19 vaccine acceptance in saudi arabia: A web-based national survey. *J Multidiscip Healthc*. 2020;13:1657–63.
72. da Silva DT, Biello K, Lin WY, Valente PK, Mayer KH, Hightow-Weidman L, et al. Covid-19 vaccine acceptance among an online sample of sexual and gender minority men and transgender women. *Vaccines (Basel)*. 2021;9(3):1–10.
73. Lin Y, Hu Z, Zhao Q, Alias H, Danaee M, Wong LP. Understanding COVID-19 vaccine demand and hesitancy: A nationwide online survey in China. *PLoS Negl Trop Dis*. 2020 Dec 1;14(12):e0008961.
74. Fernandes N, Costa D, Costa D, Keating J, Arantes J. Predicting covid-19 vaccination intention: The determinants of vaccine hesitancy. *Vaccines (Basel)*. 2021 Oct 1;9(10).

75. Ajzen I. From intentions to action: a theory of planned behavior. In: J. Kuhl & J. Beckmann, editor. Action-control: From cognition to behavior. Heidelberg: Springer; 1985. p. 11–39.
76. Prochaska, J.O., Redding, C.A., & Evers KE. The transtheoretical model and stages of change. In: K. Glanz BKR& KV, editor. Paper Knowledge Toward a Media History of Documents. San Francisco: Jossey-Bass; 2014. p. 125–48.
77. Rosenstock IM. The Health Belief Model and preventive health behavior. Health Educ Monogr. 1974;2(4):354–84.

STATEMENT OF EACH AUTHOR'S CONTRIBUTION:

Marina Almeida-Silva (MAS): Conceptualization, Methodology, Quality Assessment and Writing- Reviewing and Editing.

Lina Vieira (LV): Writing- Original draft preparation, Quality Assessment and data treatment.

Ana Grilo (AG): Methodology and Writing- Original draft preparation.

Luísa Pedro (LP): Writing- Original draft preparation.

André Coelho (AC): Methodology.

Elisabete Carolino (EC): Writing- Reviewing and Editing.

Oleksandra Umamets (OU): Writing- Reviewing and Editing.

Maria de Graça Andrade (MGA): Conceptualization, Methodology, Quality Assessment and Writing- Reviewing and Editing.

SUPPLEMENTARY MATERIAL

S1 - Overall Quality Assessment

Table S1.1 shows the overall Quality Assessment based on JBI Critical Appraisal Checklist for Analytical Cross-Sectional Studies. Twenty-nine studies were considered eligible to be assessed and its quality is presented as supplementary material (**Error! Reference source not found.**).

Figure S1.2 presents the graphical scheme of the Critical Appraisal using the robvis tool.

S2 - General characterization of the Results

Error! Reference source not found. summarizes the data provided from each eligible article, such as: Year, Country, Theory based, Goals, Participants' recruitment and Sampling Methodology.

S3 – Socio-demographic characteristics of the Results

Error! Reference source not found. presents the socio-demographic characteristics of the total 28 eligible articles assessed under this systematic review




Figure 3 - Search and study selection PRISMA flow diagram.




Figure 4 - Summary of the Critical Appraisal based on JBI checklist (risk of bias).

Accepted Manuscript




Figure 3 - Determinants of COVID-19 Vaccine intention by number of articles and percentage of confirmation (The quadrant in which each determinant is found depends on: 1) the number of articles that analyse each determinant (top quadrant >10 articles, bottom quadrant <10 articles); b) the percentage of articles that fully confirm the determinant's link with the intention to vaccinate (left quadrant < 60%; right quadrant \geq 60%).

Table 1 - PEO criteria for inclusion and exclusion in systematic review.

P	European and North American Adults
E	Social and/or Psychological and Determinants
O	COVID-19 vaccine intention

Accepted Manuscript

Table 2 - Detailed description of the keywords used in each electronic database.

Databases	Keywords
Pubmed	("COVID-19 Vaccines"[MeSH]) AND ("Intention"[MeSH] OR "Vaccination Refusal"[MeSH] OR "Hesitation") AND ("Social Determinants of Health"[MeSH] OR "Sociology, Medical"[MeSH] OR "Decision Theory"[MeSH] OR "Psychology Determinants")
Web of Science	ALL=("COVID-19" OR "SARS COV 2") AND ALL=("Vaccin*") AND ALL=("Intention" OR "Hesitation" OR "Refus*") AND ALL=("determinants") AND ALL=("Social" OR "Sociolog*" OR "Psychologic*")
Scopus	ALL ("COVID-19" OR "SARS COV 2") AND ALL ("Vaccin*") AND ALL ("Intention" OR "Hesitation" OR "Refus*") AND ALL ("determinants") AND ALL ("Social" OR "Sociolog*" OR "Psychologic*")
EBSCO; CINAHL; EMBASE; PsycINFO	COVID-19 or sars-cov-2 AND Vaccine or Vaccination AND Intention OR Hesitation OR Refusal

Table 3 - Detailed description of the inclusion and exclusion criteria.

Inclusion criteria	Exclusion criteria
- European and North America studies	- Health professionals
	- Children (<18 years old)
- Quantitative studies	- Parents of children < 18 years old
	- Parents representing children < 18 years old
- Adults	- Review and meta-analysis studies
- Articles in English, Portuguese	

Accepted Manuscript

	Disease Severity	Perceived Risk to be Infected
(47)	Perceived severity of COVID-19 infection: > Intention	Perceived likelihood of getting COVID-19 infection in the future: > Intention
(40)	Perceived threat & Concern of getting infected with COVID-19 in the next year: >Intention	Perceived threat & Concern of getting infected with COVID-19 in the next year: > Intention
(25)	Perceived risk of COVID-19 for population: > Intention	Probability of being infected: NSSD
(44)	Personal risk of COVID-19: NSSD	NE
(41)	Health threat appraisal: > Intention	Worried about COVID-19 infecting family or self: NSSD
(37)	COVID- 19 health threat: > Intention	
(38)	Worried about severe disease: > Intention	NE
(18)	Disease severity: > Intention	
(39) ¹	Worry about covid: > Intention (Germany, Italy, Sweden)	Perceived infection risk: > Intention (USA, Germany, Italy)
(56)	COVID-19 risk: > Intention	
(58)	Fear of COVID-19: > Intention	NE
(59)	COVID-19 Perceived risks: > Intention	
(52)	Fear of COVID-19: NSSD	NE
	Risk of COVID-19: NSSD	
(50) ²	Fear of COVID-19: > Intention	Fear to be infected: > Intention
	Perceived Severity: > Intention	
	Perceived susceptibility: > Intention	
(51)	NE	Having risk factors for COVID-19: > Intention

Table 4 - List of Beliefs: Disease Severity and Perceived Risk to be Infected.

1 - Only stronger predictors are mentioned, due to the high number of determinants and countries under study. Countries with more than one evaluation, only the last was considered. 2 - Only results for combined samples (Ireland & UK) are presented; NE - Not Evaluated; NSSD - Non-Statistically Significant Determinant

Table 5 - List of Beliefs: Vaccine Costs & Risks, Cost perception and Vaccine efficacy.

	Vaccine Costs & Risks	Control perception	Vaccine efficacy
(47)	Perceived potential harm of a COVID-19 vaccine: > Intention	NE	Perceived effectiveness of a COVID-19 vaccine: > Intention
(25)	Vaccine side-effects: < Intention	NE	NE
(49)	Vaccine side-effects: < Intention	NE	Perceived effectiveness of a COVID-19 vaccine: > Intention
(38)	Worries about potential side effects: < Intention	NE	Vaccine efficacy: NSSD (<50 Years Old) Vaccine efficacy: > Intention (>50 Years Old)
(36)	Vaccine Negative attitude ¹ : < Intention Mistrust of vaccine benefits: < Intention Worries about unforeseen future events: < Intention Preference for natural immunity: < Intention	Control ¹ : > Intention	Confidence in the COVID-19 vaccine: > Intention
(35)	NE	Control to get the vaccine when it becomes available ¹ : Autonomy: NSSD for 18-49 years old; < Intention for > 50 years old (only White) Self-efficacy: NSSD for 18-49 years old; > Intention for 50 years old+ (only Black)	Attitudes (favourability toward being vaccinated) ¹ : > Intention for 18-49 years old (White, Black, but not in Hispanic) >Intention for > 50 years old (for the three ethnic groups)
(55)	Short or long-term side effects: < Intention	NE	NE
(18)	NE	Control to get vaccinated: NSSD	Perceived vaccine benefits: > Intention
(56)	Side effects of Vaccine: NSSD	Vaccine being easy to get: > Intention	NE
(43)	Anti-vaccine beliefs: < Intention	NE	NE
(45)	Vaccine negative attitude ¹ : < Intention	Control against COVID-19: NSSD	NE
(59)	Doubts about vaccine: < Intention	NE	NE
(52)	Vaccine Negative attitude ¹ : < Intention	Perceived control ¹ : NSSD	Scepticism toward vaccines: NSSD
(50) ²	Vaccine risk: < Intention	NE	Perceived benefit: > Intention

1 - Theory of Planned Behavior; 2 - Only results for combined samples (Ireland & UK) are presented; NE - Not Evaluated; NSSD - Non-Statistically Significant Determinant.

Table 6 - List of Beliefs: Personal Concerns, Social Concerns and Social Norms

	Personal concern	Social concerns	Social Norms
(42)	NE	Ireland and UK sample: Altruistic motivations: > Intention	NE
(36)	NE	NE	Subjective norms: > Intention
(35)	NE	NE	Normative pressure ¹ : NSSD in 18-49 years old (for the three ethnic groups); > Intention in >50 years old (for the three ethnic groups)
(49)	NE	Collective Importance of COVID-19 vaccine: > Intention	NE
(38)	Assumed protection of oneself was a reason to get vaccine: > Intention	NE	NE
(55)	Will protect one-self: NSSD	Vaccination protects others (moral duty): > Intention	NE
		Vaccination is key to reopening society: NSSD	NE
(18)	Anticipated regret of not having a vaccine: > Intention	NE	Subjective norm ² : > Intention
(39) ³	NE	Pro-sociality (to do things for the benefit of others and society): NSSD	NE
(45)	Anticipated regret (not to vaccinate): NSSD	Perceived community benefits: > Intention	Subjective norm ² : NSSD
(59)	NE	Protection of others (Altruistic motivations): > Intention	NE
(50) ⁴	Vaccine attitudes (need to be vaccinated): >Intention	Civic responsibility:> Intention Moral duty (VS Choice)> Intention	Peer influence: > Intention GP influence: > Intention Govern. influence:>Intention
(52)	NE	NE	Subjective norm ² :< Intention

1 - Normative pressure: perceptions about what others think and do about performing the behavior; 2 - Theory of Planned Behavior; 3 - Only more stronger predictors are mentioned, due to the due to high number of determinants and countries under study. Countries with more than one evaluation, only the last was considered; 4 - Only results for combined samples (Ireland & UK) are presented; NE - Not Evaluated; NSSD - Non-Statistically Significant Determinant.

Table 7 - List of Beliefs: Trust on Media, Social & Scientific Entities; Conspiracy Beliefs; Emotion & Personality

	Trust on Media, Social & Scientific Entities	Conspiracy Beliefs	Emotions & Personality
(55)	Trust in Science or Institutions: > Intention	NE	
(54)	NE	NE	Anxiety: > Intention
(42)	Ireland and UK, respectively Trust in scientists, health professionals and state: > Intention and NSSD	Ireland and UK, respectively Conspiracy beliefs: < Intention and NSSD Paranoia: NSSD and < Intention	NE
(25)	Perceived trust in the virus management: NSSD	NE	NE
(49)	NE	Conspiracy beliefs: < Intention	NE
(38)	Vaccine given are safe: > Intention (>50Y) Recommendation from a health professionals > Intention (<50Y)	NE	NE
(37)	Trust in health authorities: > Intention Use of media as inform source: > Intention	NE	NE
(39) ¹	Trust in National sci/med advisors > Intention (Spain and EUA) Trust in experts: > Intention (Spain) Trust in WHO: > Intention (Germany)	NE	NE
(44)	NE	Conspiracy beliefs: < Intention	NE
(59)	Trust in politics and science: > Intention	NE	NE

1- Only stronger predictors are mentioned, due to the due to high number of determinants and countries under study. Countries with more than one evaluation, only the last was considered; 2 - Embitterment defined as a feeling of disappointment and anger in response to perceived injustices by others toward the self and to negative life events, and feelings of social exclusion, a lack of social support and the perception of unrightfully been let down; 3 - Only results for combined samples (Ireland & UK) are presented; NE - Not Evaluated; NSSD - Non-Statistically Significant Determinant.