


CQB-day 2016

June 28th



Book of Abstracts



Ciências
ULisboa

CQB
Centro
de Química
e Bioquímica

FCT

Fundação para a Ciência e a Tecnologia
MINISTÉRIO DA CIÊNCIA, TECNOLOGIA E ENSINO SUPERIOR

CQB-Day 2016

June 28th

Faculdade de Ciências – Universidade de Lisboa



Ciências
ULisboa

Faculdade
de Ciências
da Universidade
de Lisboa



FCT
Fundação para a Ciência e a Tecnologia
MINISTÉRIO DA CIÊNCIA, TECNOLOGIA E ENSINO SUPERIOR

ORGANISING COMMITTEE

ANA PAULA CARVALHO

OLINDA C. MONTEIRO

ANA MOURATO

WELCOME TO CQB-DAY 2016

Centro de Química e Bioquímica (CQB) celebrates 15 years since its foundation in 2001, with the aim of creating an environment oriented toward fruitful collaborations between chemistry and biochemistry groups.

The Strategic Project for 2015-2020 (PEst 2015-2020), defined in 2014, and approved by FCT, should strengthen the links between groups around two thematic lines, reflecting the interest and expertise of CQB:

1. *Chemistry and biochemistry for a clean environment.*
2. *Healthy life: molecular interventions and regulation mechanisms.*

During the last CQB-Day, in September 2015, all the groups had the opportunity to present their contributions towards these topics, either orally (selected contributions highlighting collaborative work within the thematic line) or as a poster. Half-way through 2016 and with the triennium evaluation planned by FCT at national level for the second half of 2017, we propose to carry out an internal mid-term evaluation to check our progress in achieving the project goals.

The program of CQB Day 2016 will open with a brief historical account of CQB, by its first coordinator, José Artur Martinho Simões. Lectures will follow, addressing the work of the two thematic lines and the contribution of the science manager, as well as new projects funded and opportunities for the future. A large number (around 80) of poster presentations will contribute to give a general idea of the wealth of ongoing projects and to increase the degree of cross-fertilization between the twelve groups, their members and collaborators. Two members of the External Advisory Committee (Sir William Wakeham and Hans Peter Wessel) will be present and will help us in the self-evaluation exercise. A round table will be dedicated to discuss CQB Activity and Future.

We count on all of you to make this CQB Day very successful and use this unique opportunity to discuss the implementation and progress of our Strategic Project, to strengthen collaborations between groups supported by novel scientific synergies, to find new common grounds for research and grasp the most advantageous funding opportunities.

Maria José Calhorda
CQB coordinator

SCIENTIFIC PROGRAM

9.30h CQB: 15 years and beyond

Martinho Simões

Maria José Calhorda

10.00h Science at CQB - Oral Communications

O1. *Chemistry and Biochemistry for a Clean Environment*

Carla D. Nunes

O2. *“Healthy Life”: Interaction of Polyphenols with Lipid Bilayers and their Effects in Human Cells*

Hugo A. L. Filipe, Catarina Peneda, Joaquim T. Marquês, Miguel Machuqueiro, João C. Ramos, Maria da Soledade Santos, H. Susana Marinho, Ana S. Viana, Helena Soares, Rodrigo F. M. de Almeida

O3. *S&T Management at CQB*

Ana Mourato

10.45h Science at CQB – Poster Communications & Coffee Break

11.30h Science at CQB - Oral Communications

O4. *Exploring the Catalytic Behaviour of Hierarchical Mcm-22 Zeolite in Low Temperature Friedel-Crafts Acylation of Heteroaromatics*

Nelson Nunes, Rodrigo Aleixo, Ruben Elvas-Leitão, Filomena Martins, Ana P. Carvalho, Amadeu Brigas, Angela Martins

O5. *New Insights on the Immobilization Mechanism of Escherichia Coli Onto Activated Carbons*

Susana Marques, Marta Pacheco, Jossano Marcuzzo, Ana S. Mestre, Ricardo Dias, Ana P. Carvalho

O6. *CoFe₂O₄ Nanoparticles Synthesized with Natural Templates for Magnetic Hyperthermia*

Liliana P. Ferreira, Maria Margarida Cruz, Maria L. Oliveira, Sofia G. Mendo, André F. Alves, Maria Helena Mendonça, Margarida Godinho, Maria Deus Carvalho

O7. *Herbal Infusions in Age-Related Diseases*

Pedro Luis Falé, Rita Pacheco, Maria Helena Florêncio, Maria Luisa Serralheiro

12.30h – 14.30h Lunch & Science at CQB – Poster Communications

14.30h CQB Scientific Activity and Future

Chairperson: Martinho Simões

Miguel Castanho, Fundação para a Ciência e Tecnologia

Hans Peter Wessel, Universidade de Aveiro

Sir William Wakeham, University of Southampton

15.30h FCT Funded Projects Launched in 2016

Flexible Biomimetic/Nanoconjugated Platforms for Sensitive Immunosensing

Ana S. Viana

CpHMD-L Simulations of Phlip Peptides: Design of New Tumor-Targeted Drug Delivery Systems

Diogo Vila-Viçosa, Pedro Reis, Vitor H. Teixeira, Maria J. Calhorda, António M. Baptista, Miguel Machuqueiro

Multifunctional Luminescent Spin Labile Hybrid Materials

Paulo N. Martinho, Liliana P. Ferreira, Maria de Deus Carvalho, Sara Realista, Nuno A. G. Bandeira, d, Janaína C. Almeida, Frederico F. Martins, Maria José Calhorda

*Sphingolipid Organization in the Plasma Membrane of *Saccharomyces Cerevisiae**

Rodrigo F. M. de Almeida

16.10h Science at CQB - Oral Communications

O8. *New Generation of Microextraction Devices for Trace Analysis Based on Floating Sampling Technology*

Alessandra H. Ide, Alexandra M. Fernandes, André M. Segurado, Samir M. Ahmad, Nuno R. Neng, José M. F. Nogueira

O9. *Unraveling The Role Of Tbccd1 Protein On Cell Size Control: The Regulation Of Cytoskeleton Dynamics And Cell Junctions*

Carolina Camelo, Catarina Peneda, Étienne Coyaude, Brian Raught, Ana I. Câmara, Bruno Carmona, Francisco Pinto, H. Susana Marinho, Helena Soares

O10. *Exploratory Chemistry of Butyrylcholinesterase Nucleoside-Based Inhibitors*

Vasco Cachatra, Ignazio Schino, Nicola Colabufo, Amélia P. Rauter

17.00h Closing Session

Maria José Calhorda

17.20h Happy Hour

INDEX

ORAL COMMUNICATIONS

O1	CHEMISTRY AND BIOCHEMISTRY FOR A CLEAN ENVIRONMENT <u>Carla D. Nunes</u>	3
O2	“HEALTHY LIFE”: INTERACTION OF POLYPHENOLS WITH LIPID BILAYERS AND THEIR EFFECTS IN HUMAN CELLS <u>Hugo A. L. Filipe</u> , Catarina Peneda, Joaquim T. Marquês, Miguel Machuqueiro, João C. Ramos, Maria da Soledade Santos, H. Susana Marinho, Ana S. Viana, Helena Soares, Rodrigo F. M. de Almeida.....	4
O3	S&T MANAGEMENT AT CQB <u>Ana Mourato</u>	5
O4	EXPLORING THE CATALYTIC BEHAVIOUR OF HIERARCHICAL MCM-22 ZEOLITE IN LOW TEMPERATURE FRIEDEL-CRAFTS ACYLATION OF HETEROAROMATICS <u>Nelson Nunes</u> , Rodrigo Aleixo, Ruben Elvas-Leitão, Filomena Martins, Ana P. Carvalho, Amadeu Brigas, Angela Martins.....	6
O5	NEW INSIGHTS ON THE IMMOBILIZATION MECHANISM OF <i>ESCHERICHIA COLI</i> ONTO ACTIVATED CARBONS <u>Susana Marques</u> , Marta Pacheco, Jossano Marcuzzo, Ana S. Mestre, Ricardo Dias, Ana P. Carvalho.....	7
O6	CoFe ₂ O ₄ NANOPARTICLES SYNTHESIZED WITH NATURAL TEMPLATES FOR MAGNETIC HYPERTHERMIA Liliana P. Ferreira, Maria Margarida Cruz, Maria L. Oliveira, Sofia G. Mendo, André F. Alves, Maria Helena Mendonça, Margarida Godinho, <u>Maria Deus Carvalho</u>	8
O7	HERBAL INFUSIONS IN AGE-RELATED DISEASES Pedro Luis Falé, Rita Pacheco, Maria Helena Florêncio, <u>Maria Luisa Serralheiro</u>	9
O8	NEW GENERATION OF MICROEXTRACTION DEVICES FOR TRACE ANALYSIS BASED ON FLOATING SAMPLING TECHNOLOGY Alessandra H. Ide, Alexandra M. Fernandes, André M. Segurado, Samir M. Ahmad, Nuno R. Neng, <u>José M. F. Nogueira</u>	10
O9	UNRAVELING THE ROLE OF TBCCD1 PROTEIN ON CELL SIZE CONTROL: THE REGULATION OF CYTOSKELETON DYNAMICS AND CELL JUNCTIONS Carolina Camelo, <u>Catarina Peneda</u> , Étienne Coyaude, Brian Raught, Ana I. Câmara, Bruno Carmona, Francisco Pinto, H. Susana Marinho, Helena Soares.....	11
O10	EXPLORATORY CHEMISTRY OF BUTYRYLCHOLINESTERASE NUCLEOSIDE-BASED INHIBITORS <u>Vasco Cachatra</u> , Ignazio Schino, Nicola Colabufo, Amélia P. Rauter.....	12

FCT FUNDED PROJECTS LAUNCHED IN 2016

FLEXIBLE BIOMIMETIC/NANOCONJUGATED PLATFORMS FOR SENSITIVE IMMUNOSENSING

Ana S. Viana.....15

CpHMD-L SIMULATIONS OF pHLIP PEPTIDES: DESIGN OF NEW TUMOR-TARGETED DRUG DELIVERY SYSTEMS

Diogo Vila-Viçosa, Pedro Reis, Vitor H. Teixeira, Maria J. Calhorda, António M. Baptista, Miguel Machuqueiro.....16

MULTIFUNCTIONAL LUMINESCENT SPIN LABILE HYBRID MATERIALS

Paulo N. Martinho, Liliana P. Ferreira, Maria de Deus Carvalho, Sara Realista, Nuno A. G. Bandeira, Janaina C. Almeida, Frederico F. Martins, Maria José Calhorda.....17

SPHINGOLIPID ORGANIZATION IN THE PLASMA MEMBRANE OF SACCHAROMYCES CEREVISIAE

Rodrigo F. M. de Almeida.....18

POSTER COMMUNICATIONS

P1 SURFACES MODIFIED BY BIO-INSPIRED MOLECULES FOR MOLECULAR OXYGEN REDUCTION CATALYSIS

Rui S. Sampaio, Jorge P. Correia.....21

P2 MACROPHAGE TYPE LECTIN -GALACTOSE LIGANDS: CHEMICAL SYNTHESIS

João Manuel de Barros, Filipa Marcelo, Amélia P. Rauter.....22

P3 EXTRACTION AND ANALYSIS OF SPHINGOLIPIDS FROM SACCHAROMYCES CEREVISIAE WILD TYPE CELLS AND IN A SPHINGOLIPID BIOSYNTHETIC MUTANT

Filipe M. Moreira, Catarina A. Antunes, H. Susana Marinho, Rodrigo F.M. de Almeida.....23

P4 POLYDOPAMINE FILMS MODIFIED WITH LACCASE FOR ELECTROCHEMICAL BIOSENSING

Luís C. Almeida, Jorge P. Correia, Ana S. Viana.....24

P5 SYNTHESIS OF NEW ANTIBIOTIC GLYCOSIDES AND COMPUTATIONAL STUDIES ON THEIR INTERACTION WITH MODEL LIPID BILAYERS

Rafael Nunes, João Pais, Diogo Vila-Viçosa, Miguel Machuqueiro, Amélia P. Rauter.....25

P6 COMPARATIVE STUDY BETWEEN GRUNWALD-WINSTEIN AND TAKA PARAMETERS USED FOR THE STUDY OF SOLVENT EFFECTS

Ruben Elvas-Leitão, Ricardo J. Nunes, Filomena Martins.....26

P7 SYNTHESIS AND HEPATOTOXICITY OF PSYCHOACTIVE CATHINONES

Soraia Bronze, Rita Pacheco, Susana Santos, Helena Gaspar.....27

P8 THERMODYNAMIC STUDY OF AQUEOUS MIXTURES OF 3-BUTOXYPROPAN-1-AMINE

Ângela F. S. Santos, Maria Luisa C. J. Moita, Luis C. S. Nobre, Isabel M. S. Lampreia.....28

P9 THE SPIN CROSSOVER PROFILE IN IRON(III) SCHIFF-BASE COMPOUNDS: HALOGEN INFLUENCE

F. F. Martins, A. I. Vicente, L. P. Ferreira, M. D. Carvalho, V. H. N. Rodrigues, M. J. Calhorda, P. N. Martinho.....29

P10	THE ABILITY OF STEROLS TO FORM LIQUID ORDERED DOMAINS LINKS MEMBRANE BIOPHYSICAL PROPERTIES TO BIOLOGICAL PROCESSES <u>Catarina A. C. Antunes</u> , Alena Khmelinskaia, André E. P. Bastos, Gerson M. da S. Lobo, Rui Malhó, M. L. Cyrne, H. Susana Marinho, Rodrigo F. M. de Almeida.....	30
P11	VCL ₃ { η^3 -HC(Pz) ₃ }]@CNT AS CATALYST FOR THE MICROWAVE-ASSISTED OXIDATION OF XYLENES <u>Jiawei Wang</u> , Luísa M. D. R. S. Martins, Ana P. C. Ribeiro, Marta Saraiva, Sónia A. C. Carabineiro, José L. Figueiredo, Armando J. L. Pombeiro.....	31
P12	FORCE FIELDS FOR LIPID BILAYERS: HOW DO THEY WORK? <u>Ricardo O. Dias</u> , Diogo Vila-Viçosa, Paulo J. Costa, Miguel Machuqueiro.....	32
P13	ONE-STEP CHEMICAL IMMOBILIZATION OF LACCASE AND Fe ₃ O ₄ NANOPARTICLES ON GOLD FOR BIOSENSING APPLICATIONS <u>I. Almeida</u> , F. Henriques, M. D. Carvalho, A. S. Viana.....	33
P14	BIOACTIVE COMPOUNDS FROM AGRICULTURE AND FOOD INDUSTRY RESIDUES <u>Joana Henriques</u> , Ana Assunção, Pedro L. Falé, Maria Helena Florêncio, Rita Pacheco, Maria Luísa Serralheiro.....	34
P15	ANTICANCER AND ANTIOXIDANT ACTIVITY FROM CORK WASTE WATER <u>Luís Gomes</u> , Maria Luísa Serralheiro, Miguel Minhalma, Rita Pacheco.....	35
P16	SYNTHESIS OF NOVEL NUCLEOS(T)IDE AND GLYCOSYL PHOSPHATE ANALOGS OR MIMETICS POSSESSING THERAPEUTIC POTENTIAL <u>Nuno M. Xavier</u> , Daniela Batista, Alexandre Porcheron, Vítor J. Martins, Andreia Fortuna, Paulo J. Costa, Stefan Schwarz, Anne Loesche, René Csuk, Radek Jorda.....	36
P17	SUSTAINABLE CARBON MATERIALS AS SUPPORT FOR OXIDATION CATALYSTS Cécile Petit, Miguel V. Silva, Ana S. Mestre, Ana P. Carvalho, <u>Carla D. Nunes</u>	37
P18	MOLYBDENUM(II) COMPLEXES AS CATALYSTS IN REDUCTION REACTIONS <u>Marta S. Saraiva</u> , Ana Condinho, Maria José Calhorda.....	38
P19	RECOVERY OF PHENOLIC COMPOUNDS FROM INFUSIONS: ANTIACETYLCHOLINESTERASE AND ANTIOXIDANT ACTIVITY <u>João Carvalho</u> , Inês Vasco, Andreia Maia, Tiago Santos, Rita Pacheco, Maria Luísa Serralheiro.....	39
P20	MODELLING HALOGEN BONDS IN MOLECULAR DYNAMICS SIMULATIONS OF BACTERIOPHAGE T4 LYSOZYME <u>Ricardo J. M. Rosa</u> , Diogo Vila-Viçosa, Miguel Machuqueiro, Paulo J. Costa.....	40
P21	SPONTANEOUS INCORPORATION OF SILVER PARTICLES IN THIOPHENE-BASED CONDUCTING POLYMERS Ricardo M. Assunção, <u>Jorge P. Correia</u>	41
P22	MOB1, HIPPO PATHWAY MEMBER, IS CRITICAL FOR <i>TOXOPLASMA GONDII</i> REPLICATION Alexandra Tavares, Inês Delgado, Samuel Francisco, João Coelho, Alexandre Leitão, <u>Helena Soares</u> , Sofia Nolasco.....	42


P23	RESOURCE ALLOCATION TO MAINTENANCE AND STRESS RESPONSE: SET UP OF A QUANTITATIVE EXPERIMENTAL MODEL <u>Miguel A. Rendas</u> , Rafael Bento, Carlos Bernardes, Manuel E.M. da Piedade; Fernando Antunes.....	43
P24	REORGANIZATION OF PLASMA MEMBRANE LIPID DOMAINS IN <i>NEUROSPORA CRASSA</i> DURING CONIDIAL GERMINATION <u>Filipa C. Santos</u> , Andreia S. Fernandes, Catarina A. C. Antunes, Filipe P. Moreira, Arnaldo Videira, H. Susana Marinho, Rodrigo F. M. de Almeida.....	44
P25	INSECTICIDAL ACTIVITY OF TROPICAL AND SUBTROPICAL PLANT ESSENTIAL OILS AGAINST <i>Aedes Aegypti</i> LARVAE <u>Diara Rocha</u> , Maria Novo, Olívia Matos, Ana C. Figueiredo, Manuel Delgado, Cristina Moiteiro.....	45
P26	PREPARATION AND PHYSICOCHEMICAL AND STRUCTURAL CHARACTERIZATION OF NATURAL DEEP EUTECTIC SOLVENTS <u>Frederico Ferreira</u> , <u>Lucas Ribeiro</u> , Ruben Elvas-Leitão, Filomena Martins.....	46
P27	PERSONALISED ICT SUPPORTED SERVICE FOR INDEPENDENT LIVING AND ACTIVE AGEING: THE CONTRIBUTION OF FUNCTIONAL INGREDIENTS AND THE WEBSITE NUTRIAGEING <u>Alice Martins</u> , Marta Sousa Silva, António Ferreira, Luís Sousa, Catarina Dias, Ana Marta Matos, Vasco Cachatra, Daniela Batista, Antónia Turkman, Feridun Turkman, Marília Antunes, Helena S. Costa, Tânia Albuquerque, Ana Sanches, Amélia P. Rauter.....	47
P28	SYNTHETIC COBALT CLAYS FOR THE STORAGE AND SLOW RELEASE OF THERAPEUTIC NITRIC OXIDE <u>Ana C. Fernandes</u> , Moisés L. Pinto, Fernando Antunes, João Pires.....	48
P29	METAL-ORGANIC FRAMEWORKS AS POTENTIAL NITRIC OXIDE STORAGE AND DELIVERY VEHICLES FOR THERAPEUTIC APPLICATIONS <u>Moisés L. Pinto</u> , Paula Brandão, Rosana V. Pinto, Fernando Antunes, João Pires.....	49
P30	CAPTURE AND ACTIVATION OF CO ₂ BY METAL-ORGANIC STRUCTURES <u>Janaína C. Almeida</u> , Sara Realista, Sónia Barroso, Maria José Calhorda, Paulo N. Martinho.....	50
P31	PHENOLIC QUANTIFICATION IN DIFFERENT CAMELLIA SINENSIS INFUSIONS AND RELATION WITH THEIR ANTIOXIDANT ACTIVITY Rute Martins, Daniela Jones Jones, Ana Marques, Maria Helena Florêncio, Rita Pacheco, <u>Maria Luisa Serralheiro</u>	51
...		
P32	DEVELOPMENT OF LYSOSOMAL-MIMICKING VESICLES TO STUDY THE EFFECT OF SPHINGOSINE ABNORMAL ACCUMULATION ON MEMBRANE BIOPHYSICAL PROPERTIES <u>Ana C. Carreira</u> , Rodrigo F. M. de Almeida, Liana C. Silva.....	52
P33	INTERACTION OF D-GLUCOSE END-CAPPED POLYLACTIDE RUTHENIUM CYCLOPENTADIENYL COMPLEXES WITH GLUT: INSIGHTS FROM MOLECULAR DOCKING <u>Carla Antunes</u> , Andreia Valente, Paulo J. Costa.....	53

P34	USING SOLVATOCHROMIC PROBES FOR THE STUDY OF INTERMOLECULAR INTERACTIONS IN 1,4-DIOXANE/METHANOL/ACETONITRILE SOLVENT MIXTURES <u>Ricardo J. Nunes</u> , Nelson Nunes, Ruben Elvas-Leitão, Filomena Martins.....	54
P35	GLYCOSIDE ANTIBIOTICS AND MECHANISM OF ACTION <u>João P. Pais</u> , Ana S. Viana, Amélia P. Rauter, Luís. Sobral, Ricardo Dias, Rogério Tenreiro.....	55
P36	BIOCIDAL MICROCAPSULES: A NEW STRATEGY FOR BIOFOULING CONTROL <u>Elisabete R. Silva</u> , Olga Ferreira, Raquel B. Feio, João C. M. Bordado, Patrícia Rijo, Ana C. Marques.....	56
P37	BIOMEMBRANE ORGANIZATION: BIOPHYSICAL CHARACTERIZATION OF ORDERED DOMAINS <u>Gerson M. Lobo</u> , Telmo O. Paiva, Ana S. Viana, Rodrigo F. M. de Almeida.....	57
P38	STUDY OF THE PERMEABILITY OF PHENOLIC COMPOUNDS PRESENT IN CLADODES OF <i>OPUNTIA FICUS INDICA</i> AND THEIR EFFECT ON THE CHOLESTEROL PERMEABILITY THROUGH CACO-2 CELL LINES <u>Asma Ressaissi</u> , Pedro L. Falé, Nebil Attia, Rita Pacheco, Maria Luísa M. Serralheiro.....	58
P39	MULTIFUNCTIONAL LUMINESCENT SPIN LABILE HYBRID MATERIALS <u>Paulo N. Martinho</u> , Liliana P. Ferreira, Maria de Deus Carvalho, Sara Realista, Nuno A. G. Bandeira, Janaína C. Almeida, Frederico F. Martins, Maria José Calhorda.....	59
P40	CHARACTERIZATION AND IDENTIFICATION OF ESSENTIAL OILS BY GC-MS <u>J. Aço</u> , M. H. Florêncio, C. Borges, A. P. Marques.....	60
P41	[FeCl ₂ (HCPz ₃)] HETEROGENIZED AT HIERARCHICAL FAU FOR ECO-FRIENDLY ALKANE OXIDATIONS Vanmira Van-Dúnem, Ana P. Carvalho, Armando J. L. Pombeiro, Luísa M. D. R. S. Martins, <u>Angela Martins</u>	61
P42	FUNCTIONAL PROTEIN-BASED NANOPARTICLES FOR REDUCTION OF CARDIOVASCULAR DISEASES RISK <u>André F. Silva</u> , Joaquim T. Marquês, Ana S. Viana, Rita Pacheco, Maria Luísa M. Serralheiro.....	62
P43	CO ₂ ELECTROREDUCTION BY BINUCLEAR Ni(II) AND Cu(II) COMPLEXES <u>Paulo N. Martinho</u> , Sara Realista, Paulo J. Costa, Maria José Calhorda.....	63
P44	HETERODINUCLEAR Ni(II) AND Cu(II) SCHIFF-BASE COMPLEXES AND THEIR OXYGEN REDUCTION MATERIALS <u>Priscila C. Ramgi</u> , Sara Realista, Bernardo de P. Cardoso, Ana I. Melato, Ana S. Viana, Maria José Calhorda, Paulo N. Martinho.....	64
P45	ENERGY-FREE REDUCTION OF SILVER FROM DILUTE SOLUTIONS BY ELECTROLESS PRECIPITATION (RE-)USING POLYANILINE FILMS <u>Inês J. Pereira</u> , Jorge P. Correia.....	65
P46	LUMINESCENT HYDROPHOBIC SURFACES FOR SMART WINDOWS <u>J. B. Aldeias</u> , S. Realista, M. J. Calhorda, P. N. Martinho.....	66

P47	SYNTHESIS AND PHOTOCATALYTIC ACTIVITY EVALUATION OF NEW TITANATE NANOTUBES MODIFIED BY CO-DOPING WITH COBALT AND RHENIUM <u>B. Barrocas</u> , O. C. Monteiro	67
P48	SYNTHESIS OF NEW CNS-TARGETED DRUG-LIKE LEADS AGAINST ALZHEIMER'S DISEASE: THE ROLE OF THE SUGAR MOIETY IN C-GLUCOSYL FLAVONOID ANALOGUES <u>Ana M. Matos</u> , Joana Cristóvão, Cláudio Gomes, Amélia P. Rauter	68
P49	REVERSE SELECTIVITY OF ZEOLITES AND MOFs IN THE ETHANE/ETHYLENE SEPARATION BY ADSORPTION <u>João Pires</u> , Joana Fernandes, Ana C. Fernandes, Moisés L. Pinto	69
P50	OPI1P TRANSLOCATION TO THE NUCLEUS IS REGULATED THROUGH OXIDATION BY HYDROGEN PEROXIDE IN <i>SACCHAROMYCES CEREVISIAE</i> <u>C. Camelo</u> , F. Vilas-Boas, A. Cepeda, C. Real, F. Pinto, G. da Costa, H. Soares, H. S. Marinho, L. Cyrne	70
P51	NOVEL BIOCL-BI ₂ S ₃ NANOSTRUCTURES SYNTHESIZED FROM DEEP EUTECTIC SOLVENTS AND ITS PHOTOCATALYTIC PROPERTIES <u>Virgínia C. Ferreira</u> , Olinda C. Monteiro	71
P52	DEVELOPMENT OF A LC-MS/MS METHOD FOR QUANTITATIVE ANALYSIS OF NINE PHARMACEUTICAL COMPOUNDS IN SURFACE WATER SAMPLES <u>Rodrigo A. Osawa</u> , Maria Helena Florêncio, Maria Rosário Bronze	72
P53	ACHe INHIBITION AND ANTIOXIDANT ACTIVITY OF <i>CENTAURIUM ERYTHRAE</i> (FEL DA TERRA) INFUSIONS <u>Laura Guedes</u> , Samuel Barata, Maria Helena Florêncio, Rita Pacheco, Miguel Machuqueiro, Maria Luísa Serralheiro	73
P54	NOVEL TITANATE NANOTUBULAR MATERIALS MODIFIED WITH TRANSITION METALS FOR THE PHOTOCATALYTIC DEGRADATION OF PPCPs <u>J. C. Cardoso</u> , O. C. Monteiro	74
P55	A NEW SEMI-EMPIRICAL EQUATION TO DESCRIBE THE SURFACE TENSION OF AQUEOUS ORGANIC LIQUID MIXTURES <u>M. Soledade C.S. Santos</u> , João Carlos R. Reis	75
P56	EXPLOITING 2-DEOXY GLYCOSIDES TOWARDS NEW ANTIMICROBIAL AND NEUROPROTECTIVE COMPOUNDS <u>Catarina Dias</u> , A. P. Rauter	76
P57	ACETYLCHOLINESTERASE INHIBITION, ANTIOXIDANT ACTIVITY, <i>IN VITRO</i> GASTROINTESTINAL DIGESTION AND TOXICITY OF CLADODES FROM <i>OPUNTIA FICUS INDICA</i> EXTRACTS ON HEPG2, MCF-7 AND CACO-2 CELL LINES <u>Asma Ressaissi</u> , Rita Pacheco, Nebil Attia, Maria Luisa Serralheiro	77
P58	ARE pH-SENSITIVE AMINO ACIDS IN THEIR MOST COMMON PROTONATION STATES AT THE WATER/MEMBRANE INTERFACE? <u>Pedro B. P. S. Reis</u> , Vitor H. Teixeira, Diogo Vila-Viçosa, Miguel Machuqueiro	78

P59	MN(III) SINGLE ION MAGNET WITH A TRIDENTATE SCHIFF-BASE LIGAND <u>S. Realista</u> , A. J. Fitzpatrick, G. Santos, L. P. Ferreira, S. Barroso, L. C. J. Pereira, N. A. G. Bandeira, P. Neugebauer, G. G. Morgan, J. van Slageren, M. J. Calhorda, P. N. Martinho.....	79
P60	POLYMORPHISM OR THERMAL DISORDER IN 4-HYDROXYHEPTANOPHENONE? <u>Cátia S. D. Lopes</u> , M. Fátima M. Piedade, Ricardo G. Simões, Herminio Diogo, Carlos E. S. Bernardes, Manuel E. Minas da Piedade.....	80
P61	PHOTOCATALYTIC DEGRADATION OF CARBAMAZEPINE AND IBUPROPHEN IN WATER USING NOVEL TiO ₂ -MOO ₂ MATERIALS <u>J. Diniz</u> , C. D. Nunes, O. C. Monteiro.....	81
P62	AMINO ACID INTERCALATED CLAYS AS GREEN MATERIALS FOR CARBON DIOXIDE ADSORPTION <u>Joanna Jużkówa</u> , Ana C. Fernandes, Moisés L. Pinto, João Pires.....	82
P63	pH-DEPENDENT INSERTION OF pHLIP PEPTIDE INTO LIPID BILAYERS: pK _a VALUES OF KEY RESIDUES <u>Diogo Vila-Viçosa</u> , Vitor H. Teixeira, Miguel Machuqueiro.....	83
P64	[VO ₂ {HB(3,5-ME ₂ PZ) ₃ }] SUPPORTED AT HIERARCHICAL MOR FOR ECO-FRIENDLY CICLOHEXANE OXIDATION <u>Davide Ottaviani</u> , Ana P. Carvalho, Luísa M. D. R. S. Martins, Armando J. L. Pombeiro, Angela Martins.	84
P65	NEW INTERMETALLIC ELECTROCATALYSTS FOR HYDROGEN PRODUCTION <u>D. Siopa</u> , A. P. Gonçalves, J. Branco, A. Gomes.....	85
P66	BIOACTIVITY AND PHYTOCHEMICAL STUDIES ON TUNISIAN GLOBE ARTICHOKE (CYNARA SCOLYMUS) WASTES <u>Faiza Mejria</u> , Alice Martins, Karim Hosni, Hervé Casabianca, Amélia P. Rauter.....	86
P67	THE EFFECT OF P. BARBATUS AND P. ZULUENSIS INFUSIONS ON PROTEIN SECONDARY STRUCTURE INVESTIGATED USING FTIR SPECTROSCOPY <u>Elsa Brito</u> , André Silva, Pedro L. Falé, Lia Ascensão, Parvez I. Haris, António J. Serralheiro, Maria L. M. Serralheiro, Rita Pacheco	87
P68	WHY IS INH-C ₁₀ MORE ACTIVE THAN INH AGAINST MYCOBACTERIUM TUBERCULOSIS? DECODING THE PUZZLE... <u>Filomena Martins</u> , Miguel Machuqueiro, Peter C. Loewen.....	88
P69	IDENTIFICATION OF PSYCHOACTIVE SUBSTANCES IN SEIZED PRODUCTS IN PORTUGAL <u>Christophe Gonçalves</u> , Cláudio R. Queirós, Sara Ciriaco, M. João Caldeira, Helena Gaspar.	89
P70	CONDUCTING POLYMERS EXHIBITING PSEUDO CATIONIC DOPING FOR CATHODES IN LI-ION BATTERIES Ariana R. Silva, <u>Jorge P. Correia</u>	90
P71	Mo NP'S AS SELECTIVE CATALYST IN THE OXIDATION OF BENZYL ALCOHOL <u>Filipe Gaspar</u> , Carla Nunes.....	91

P72	CATALYTIC ESTERIFICATION USING HYDROTHERMALLY SYNTHESIZED CARBON SPHERES <u>Tiago A.G. Duarte</u> , Luísa M. D. R. S. Martins, Ana P. Carvalho, Armando J. L. Pombeiro.....	92
P73	PHOTOCATALYTIC DEGRADATION OF WATER ORGANIC POLLUTANTS <u>Pedro Mota</u> , Joana B. Aldeias, Sara Realista, Olinda C. Monteiro, Maria José Calhorda, Paulo N. Martinho.....	93
P74	TUNING THE PORE SIZE DISTRIBUTION OF MESOPOROUS CARBONS: THE KEY ROLE OF ALKALINE CHLORIDES IN MIXTURES WITH ZINC CHLORIDE <u>Ana S. Mestre</u> , Raquel Gonçalves, Valentine Vigneron, Cristina Freire, Ana P. Carvalho.....	94
P75	INFLUENCE OF ABIOTIC EFFECTS ON ANTIOXIDANT AND POLYPHENOL CONTENT OF SOME ECONOMICALLY VALUABLE MEDITERRANEAN CROPS <u>Basma Mabrouk</u> , Bárbara Oliveira, Leila Bettaie, Ben Kaab , Maria Eduarda Araújo.....	95
P76	STEP BY STEP DEVELOPMENT OF A LIPID-BASED IMMUNOSENSOR PLATFORM WITH HIGH CAPABILITY TO INHIBIT NONSPECIFIC ADSORPTION <u>Telmo O. Paiva</u> , Inês Almeida, Joaquim T. Marquês, Yan-Yan Chen, Niu Yuc, Gang Jin, Rodrigo F. M. de Almeida, Ana S. Viana.....	96
P77	THERMOCHEMISTRY OF PEROXY RADICALS: A COMBINED PHOTOACOUSTIC CALORIMETRY, QUANTUM CHEMISTRY AND MOLECULAR DYNAMICS STUDY <u>Rui M. Borges dos Santos</u> , Elsa M. Gonçalves, Filipe Agapito, Nuno Galamba, José A. Martinho Simões.....	97
P78	THE IMPACT OF LIGAND DERIVATISATION ON Mn(III) THERMAL SPIN TRANSITIONS <u>Patrícia S. Arranja</u> , Liliana P. Ferreira, Sónia Barroso, Maria José Calhorda, Paulo N. Martinho....	98
P79	RECOVERY OF PALLADIUM FROM A SPENT INDUSTRIAL CATALYST: A HYDROMETALLURGICAL APPROACH <u>A.P. Paiva</u> , P. P. Abreu, C. A. Morgado, G. I. Carvalho, O. Orteta, C.A.Nogueira.....	99
P80	“HEALTHY LIFE”: INTERACTION OF POLYPHENOLS WITH LIPID BILAYERS AND THEIR EFFECTS IN HUMAN CELLS <u>Hugo A. L. Filipe</u> , Catarina Peneda, Joaquim T. Marquês, Miguel Machuqueiro, João C. Ramos, Maria da Soledade Santos, H. Susana Marinho, Ana S. Viana, Helena Soares, Rodrigo F. M. de Almeida.....	100
P81	DESIGNING” POROUS NETWORKS FOR WATER TREATMENT: REMOVAL OF ATENOLOL AND ANTIBIOTICS BY ACTIVATED CARBONS <u>Susana Marques</u> , Stephane Gonçalves, Ana S. Mestre, Andreja Z. Gotvajn, Ana P. Carvalho.....	101



1

Chemistry and biochemistry for a clean environment

Carla D. Nunes

Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal
cmnunes@fc.ul.pt

This thematic line was defined taking into account the expertise of the groups and the added value obtained by a closer collaboration between them, aiming at warranting a clean environment. The synthetic skills of several groups provide the conditions to develop new molecules and materials, ranging from organic to inorganic. Heterogeneous catalysts have been designed and built from bulk oxides, nanoparticles and carbon based materials, to be applied as supports (passive) or catalysts (active). Particular attention was paid to low-cost raw materials. Such combined approaches led to applications in asymmetric and photo-catalysis; capture and reduction of CO₂, photodegradation of new generation pollutants, namely carbamazepine and ibuprofen (drugs), and clofibric acid (pesticide), which have been selected as representative examples of their classes. They represent a major concern owing to the continuously increasing consumption of medicinal drugs and crop protection agents worldwide.




Figure 1: Examples of materials used in the described applications

New analytical methodologies are being developed for the detection of emerging contaminants in water resources as well as for their reduction/removal. Selection of this matrix is of utmost relevance, since water is a valuable commodity for any population and ecosystem on Earth. The developed methodologies comprised separation methods for the analysis of priority organic species, including the design, validation and application of innovative and alternative analytical approaches.

Here they will be present the achievements, contributions the last developments in this thematic line from the CQB research groups.

Acknowledgements

The authors thank FCT for financial support (UID/MULTI/00612/2013).

- A.S. Mestre, A. Nabiço, P.L. Figueiredo, M.L. Pinto, M.S.C.S. Santos, I.M. Fonseca, *Chem. Eng. J.* 2016, 286, 538–554.
G. Naudin, T. Entradas, B. Barrocas, O.C. Monteiro, *J. Mat. Sci. Tech.*, 2016 in press.
M.K.S. Batista, A. S. Mestre, I. Matos, I. M. Fonseca, A. P. Carvalho *RSC Advances*, 2016, 6, 45419-45427.
Y. Tokudome, T. Morimoto T., N. Tarutani, P.D. Vaz, C.D. Nunes, V. Prevot, G. Stenning, M. Takahashia, *ACS Nano*, 2016, 10, 5550–5559.
B. Barrocas, S. Serio, A. Rovisco, Y. Nunes, M.E.M. Jorge, *Appl. Surf. Sci.*, 2016, 360, 798-806.
O. Ortet, M.S.C.S. Santos, A.P. Paiva *Sep. Sci. Technol.*, 2016 (in press).
F.N. Andrade, A.H. Ide, N.R. Neng, F.M. Lanças, J.M. Nogueira, *J. Sep. Sci.*, 2016, 39(4), 756-761.

“Healthy Life”: Interaction of polyphenols with lipid bilayers and their effects in human cells

Hugo A. L. Filipe^{a*}, Catarina Peneda^a, Joaquim T. Marquês^a, Miguel Machuqueiro^a, João C. Ramos^a, Maria da Soledade Santos^a, H. Susana Marinho^a, Ana S. Viana^a, Helena Soares^{a,b}, Rodrigo F. M. de Almeida^a

^aCentro de Química e Bioquímica, Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Ed. C8, Campo Grande, 1749-016 Lisboa, Portugal

^bEscola Superior de Tecnologia da Saúde de Lisboa, IPL, 1990-096 Lisboa, Portugal

*hafilipe@fc.ul.pt

This work concerns the transversal project of the CQB thematic line: “Healthy Life: Molecular Interventions and Regulation Mechanisms”.

Biologically active plant phytochemicals have a broad range of pharmacological effects including anticarcinogenic, antimicrobial, antioxidant, and anti-inflammatory activity. [1] Notwithstanding the possibility of having a specific target, phytochemicals must interact and permeate through cell membranes in the body. Indeed, it was suggested that those molecules insert into the membranes and thereby may have a promiscuous activity by changing structural properties of lipid bilayers. [2]

Some well-known phenolic acids such as caffeic (CA), rosmarinic (RA) and chlorogenic (CGA) acids, whose identification in plant extracts has been achieved by CQB research groups, were selected to be addressed in first place (Figure 1).

All the phenolic acids studied have low lipophilicity and among them, RA was the only one with a partition to biological membrane models measurable by fluorescence spectroscopy, as opposed to CA and CGA. Cyclic voltammetry measurements using an electrode modified with a supported lipid bilayer, also indicated a higher affinity of RA to lipid membranes. In addition, oxidation/reduction of the phenolic acids displayed higher reversibility in the lipid milieu than in the aqueous bulk. Indeed, the reduced form of phenolic acids was unstable in aqueous solution. In particular, in DMEM/F-12 cell culture media, a colour change observed after incubation with each compound could be reverted by the addition of a reducing agent. The higher reversibility of phenolic acids oxidation/reduction, once they were inserted in the lipid membrane, may contribute to the stability of the compounds and prevent the formation of degradation products. Molecular dynamics (MD) simulations are being performed to probe the location and orientation of these and other selected compounds in lipid bilayers.

The influence of the phenolic acids in the cytoskeleton organization, both actin filaments and microtubules, of a human retinal pigment epithelial cell line (RPE1) was also investigated. All compounds induced concentration and time dependent effects, translated in structural alterations mainly at the cell periphery, and also in the perturbation of cell division. Moreover, it was not evident that these compounds induce apoptosis under the conditions tested. RA seemed to induce evident effects at earlier times and at lower concentrations, as compared to CA and CGA. This higher sensibility of RPE1 cells to RA correlates with the higher affinity of this compound to the lipid bilayer.





Figure 1. Effects in the actin filaments (red) and microtubules (green) of RPE1 cells, after 8 h incubation with 100 μ M of phenolic acid.

Acknowledgements

Support for this work was provided by F.C.T. through IF2012/2013 initiatives (P.O.P.H., F.S.E.) and Project UID/MULTI/00612/2013. H.A.L.F. acknowledges post-doc fellowship under the same Project. M.L. Serralheiro is acknowledged for the phenolic acids.

[1] A. Stefanson and M. Bakovic, *Nutrients* 6 (2014) 3777-3801.

[2] H. I. Ingólfsson, P. Thakur, K. F. Herold, E. A. Hobart, N. B. Ramsey, X. Periole, D. H. de Jong, M. Zwama, D. Yilmaz, K. Hall, T. Maretzky, H. C. Hemmings, C. Blobel, S. J. Marrink, A. Koçer, J. T. Sack and O. S. Andersen, *ACS Chem. Biol.* 9 (2014) 1788-1798.



S&T Management at CQB

Ana Mourato

Centro de Química e Bioquímica, DQB, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

**acmourato@fc.ul.pt*

A manager in Science and Technology of a research unit is usually working simultaneously at multiple interfaces and needs to establish the necessary bridges between researchers & unit coordination, researchers & society and researchers & institutions. In the Research Units, all researcher have their own ideas of what should be the functions of a manager in Science and Technology. The most common are gathering information on funding opportunities, write projects, communicate science, write press releases, manage and organize data, support the unit coordination, promote and organize events, etc. But, which of them are the most important, relevant and decisive for a Research Unit? Many are realistic, others less and, therefore, to help clarify and understand the expectations of scientists regarding this theme, a multiple-choice survey was created and distributed to all CQB members. This survey aimed to enhance and manage more effectively the interaction between researchers and Manager of Science and Technology. The information collected was also used to compare the chosen tasks with those already performed by the science manager of CQB. With this knowledge, we were able to evaluate how researchers expectations align or diverge from the work plan developed by the Science manager.

Acknowledgements

Support for this work was provided by Fundação para a Ciência e Tecnologia (FCT), Portugal, through projects UID/MULTI/00612.

Exploring the catalytic behaviour of hierarchical MCM-22 zeolite in low temperature Friedel-Crafts acylation of heteroaromatics

Nelson Nunes^{a,b}, Rodrigo Aleixo^b, Ruben Elvas-Leitão^{a,b}, Filomena Martins^a, Ana P. Carvalho^a, Amadeu Brigas^c, Angela Martins^{a,b}

^aCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Ed. C8, Campo Grande 1749-016 Lisboa, Portugal

^aÁrea Departamental de Engenharia Química, Instituto Superior de Engenharia de Lisboa, IPL, R. Cons. Emídio Navarro, 1959-007 Lisbon, Portugal

^aDepartamento de Química e Farmácia, Universidade do Algarve, Campus da Penha, Estrada da Penha, 8005-139 Faro, Portugal
*nnunes@deq.isel.ipl.pt

MCM-22 is a synthetic zeolite (Mobil 1990) with a peculiar structure, presenting three independent pore systems, two of them internal and the last one located at the external surface of the crystals. Even though this structure presents internal supercages, with a high concentration of acidic active sites, they are only accessed through narrow windows. Thus, the modification of the textural properties of this zeolite, through the creation of a micro + mesopore hierarchical structure, allows the improvement of molecular diffusion and access to the active sites, especially when larger molecules are involved. In this study hierarchical MCM-22 was prepared by performing post-synthesis basic or a combination of basic + acid treatments. The experimental procedures as well as the discussion of structural, textural and acidic characterization are reported elsewhere [1]. Friedel-Crafts acylation of heteroaromatics is an important industrial reaction that traditionally uses homogenous catalysts, such as AlCl_3 and FeCl_3 that are harmful to the environment. The use of commercial zeolites as environmentally friendly catalysts has been reported [2], however, the use of hierarchical zeolites, and especially hierarchical MCM-22, has never been explored. In this study the catalytic behaviour was investigated in the acylation of simple heteroaromatics such as furan, anisole or pyrrole, by acetic anhydride (Figure 1) using a molar ratio of 1:5 and 150 mg of zeolite, at 60 °C. Samples of the reaction mixture were analysed periodically by GC to follow conversion and yields as a function of time. Langmuir-Hinshelwood model was used to calculate kinetic parameters as well as turnover frequencies (TOF). The analysis of the results shows an improved catalytic behaviour for hierarchical MCM-22 modified through alkaline + acid treatment when compared to parent MCM-22, presenting enhanced mass transport and access to the active sites, making hierarchical MCM-22 a promising catalyst for Friedel–Crafts acylation of larger heteroaromatic molecules.




Figure 1. General scheme for the acylation of heteroaromatic compounds.

Acknowledgements

Support for this work was provided by FCT through UID/MULTI/00612/2013 and PEst-OE/QUI/UI0612/2013.

[1] V. Machado, J. Rocha, A. P. Carvalho, A. Martins, App. Catal. A: Gen. 329 (2012) 445-446.

[2] V.F.D. Álvaro, A.F. Brigas, E.G. Derouane, J.P. Lourenço, B.S. Santos, J. Molecular. Catalysis A: Chemical 305 (2009) 100-103.

New insights on the immobilization mechanism of *Escherichia coli* onto activated carbons

Susana Marques^{a,b,*}, Marta Pacheco^b, Jossano Marcuzzo^c, Ana S. Mestre^a, Ricardo Dias^b, Ana P. Carvalho^a

^aCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^bInstituto de Biosistemas & Ciências Integrativas, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^cFaculdade de Tecnologia do Estado de São Paulo FATEC - Campus São José dos Campos, 12247-014 – São José dos Campos – SP -Brasil

*scrmarques@fc.ul.pt

Adsorption on activated carbons is a well-established method used in water treatment to remove numerous pollutants. Microorganisms present in water can become immobilized in the adsorbent, possibly affecting the adsorption process. As new forms of adsorbents are developed, it is important to understand the impact of these materials' properties in the mechanism of bacteria immobilization, to conclude about the possible advantages of using these new materials in water treatment purposes.

In this work, the influence of the adsorbent morphology (granular, powder and cloth), texture and superficial chemistry in the immobilization of *E. coli* cells in aqueous media was investigated.

The results indicate that favorable electrostatic interactions carbon surface-cells is a key factor in bacteria immobilization by activated carbons that must be complemented by its morphology and textural characteristics. Thus using basic carbons, electron microscopy results revealed that powder morphology allows the formation of agglomerates carbon-cell-cell-carbon. In the case of granular and cloth samples the data already obtained suggest that, in an initial stage, individual cells adhere to the external surface of the carbon particles (more efficiently in cloth morphology). Further assays, considering different cellular densities and the simultaneous presence of pharmaceutical compounds in the aqueous medium are being performed.

Acknowledgements

The authors thank FCT, Portugal, for financial support to CQB center (Project UID/MULTI/00612/2013) and fellowships to SM (SFRH/BD/91767/2012) and ASM (SFRH/BPD/86693/2012). Salmon & Cia (Norit) is also acknowledged by kindly supplying of the commercial carbon sample.

REFERENCES

- [1] T.J. Bandoz in: T.J. Bandoz (Eds.), Activated carbon surfaces in environmental remediation, New York, 2006, pp. 159-230.
- [2] A.L. Cukierman, ISRN Chem Eng (2013) (DOI: 10.1155/2013/261523)
- [3] A.M. Paruch, Water Sci Technol. 63 (2011) 558-564.
- [4] F. Veglio, F. Beolchini, *Hydrometallurgy* 44 (1997) 301-316.
- [5] X. Bai, X. Ma, F. Xu, J. Li, H. Zhang, X. Xiao, Sci Total Environ 533 (2015) 24-31.
- [6] J. Rivera-Utrilla, I. Toledo, M.A. Ferro-García, C. Moreno-Castilla, J Chem Technol Biotechnol 76 (2001) 1209-1215.
- [7] S.C. Edberg, E.W. Rice, R.J. Karlin, M.J. Allen, J Appl Microbiol. 88(S1) (2000) 106-116.
- [8] H.P. Boehm, Carbon 32(5) (1994) 759-769.
- [9] A.S. Mestre, J. Pires, J.M.F. Nogueira, A.P. Carvalho, Carbon 45 (2007) 1979-1988.
- [10] K.S.W. Sing, Pro. Int Symp., 1970.

CoFe₂O₄ nanoparticles synthesized with natural templates for magnetic hyperthermia

Liliana P. Ferreira^{a,b}, Maria Margarida Cruz^a, Maria L. Oliveira^c, Sofia G. Mendo^c, André F. Alves^c, Maria Helena Mendonça^c, Margarida Godinho^a, Maria Deus Carvalho^{c*}

^aBioISI, Biosystems and Integrative Sciences Institute, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^bDepartment of Physics, University of Coimbra, 3004-516 Coimbra, Portugal

^c CQB, Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

*mdcarvalho@ciencias.ulisboa.pt

Magnetic nanostructured materials have a wide range of applications in nanodevices magnetic information storage media, and biomedical applications like drug delivery, resonance imaging (MRI) and magnetic hyperthermia. This latter technique uses magnetic nanoparticles under an alternating magnetic field as local heat dissipating agents for cancer therapy [1]. Besides magnetite (Fe_3O_4), other ferrites such as CoFe_2O_4 have been also investigated for such applications. Cobalt ferrite is ferrimagnetic with a high saturation magnetization and uniaxial magnetic anisotropy. These properties make CoFe_2O_4 nanoparticles (NP) good candidates for magnetic hyperthermia since anisotropic magnetic NP are expected to be associated with high heating efficiency.

Aiming at controlling the nanoparticles size and morphology, new synthesis methods have been explored in order to enhance their magnetic properties, namely the magnetic anisotropy. In this work, the synthesis of CoFe_2O_4 was performed in the presence of natural fibers, following an established methodology [2], resulting in two batches of samples: the first one are nanoparticles obtained directly from the solution, and the second one are the nanoparticles resulting from the template calcination. The structural, morphological and magnetic characterization of these particles was performed using several techniques. Their potential for magnetic hyperthermia was assessed through induction heating measurements under an alternating magnetic field and evaluated through the specific loss power (SLP) parameter using the methodology and the experimental set-up previously described in detail [3].

The NP obtained after the templates calcination show morphologies that are determined by the threaded templates, leading to magnetic coercivity values strongly enhanced when compared with other NP obtained by conventional methods. The coercive fields reach values of 1.3 T at 10 K, decreasing to 70 mT at room temperature, values comparable to the ones obtained in CoFe₂O₄ nanofibers or nanorods (Figure 1). Concerning the magnetic hyperthermia results, it was shown that for nanoparticles obtained from the solution, SLP values of ~ 7 W g⁻¹ were obtained for dispersions in water and were mainly attributed to Brown relaxation due to a high surface contribution to the magnetic anisotropy constant. These values decrease for the nanoparticles obtained after template calcination, showing that higher magnetic anisotropy does not necessarily imply higher heating efficiency and that the optimization of the NP heating performance by tailoring the NP morphology requires a careful balance between size and shape.




Figure 1. SEM image of gauze (left); TEM image of CoFe₂O₄ nanoparticles obtained after gauze calcination (middle); magnetization results of all samples obtained after template calcination (right).

Acknowledgements

Work funded by the Portuguese Foundation for Science and Technology (FCT) under the projects PTDC/CTM-BIO/2102/2012, UID/MULTI/00612/2013 and UID/MULTI/04046/2013.

- [1] A. Jordan, P. Wust, R. Scholz, H. Faehling, J. Krause, R. Felix, Magnetic fluid hyperthermia, In *Scientific and Clinical Applications of Magnetic Carriers*; ed. U. Häfeli, W. Schütt, J. Teller and M. Zborowski, Plenum Press, New York, 1997.
- [2] S. G. Mendo, A.F. Alves, L. P. Ferreira, M. M. Cruz, M.H. Mendonça, M. Godinho, M. D. Carvalho, New J. Chem., 39, (2015) 7182-7193.
- [3] A. F. Alves, S. G. Mendo, L. P. Ferreira, M. H. Mendonça, P. Ferreira, M. Godinho, M. M. Cruz, M. D. Carvalho, J. Nanopart Res. 18 (2016) 27 (13 pag).

Herbal infusions in age-related diseases

Pedro Luis Falé^{a,b}, Rita Pacheco^{a,c}, Maria Helena Florêncio^{a,b}, Maria Luisa Serralheiro^{a,b,*}

^aCentro de Química e Bioquímica. Faculdade de Ciências. Universidade de Lisboa. Campo Grande. 1749-016 Lisboa.

^bDepartamento de Química e Bioquímica. Faculdade de Ciências. Universidade de Lisboa. Campo Grande. 1749-016 Lisboa.

^cÁrea Departamental de Engenharia Química. Instituto Superior de Engenharia de Lisboa. Av. Conselheiro Emídio Navarro, 1959-007 Lisboa, Portugal.

*mlserralheiro@fc.ul.pt

The aim of the group is to study the effect of plant infusions and their components (Fig.1) in the prevention and treatment of age-related diseases such as Alzheimer disease (AD), atherosclerosis and cancer. Plant aqueous extracts have been used throughout history for their medicinal¹ and health-promoting properties. Scientific studies have related

these therapeutic properties with several molecular mechanisms on which the phenolic components of plant extracts have shown some activity. Age-related diseases are multifactorial-caused and also require treatments using multiple approaches. The main approaches selected are the inhibition of enzymatic activities involved in AD (acetylcholinesterase)², hypercholesterolemia (HMG-co reductase³ and cholesterol transporter proteins⁴). The




Figure 1: Example of herbs, LC-MS/MS, FTIR of proteins and protein-based nanoparticles

effect of bioactive molecules for cancer treatment is also an objective of the group, focusing on the expression of proteins and on metabolomics. A relationship between bioactivity and chemical constitution of the plant extracts has been established for several plant extracts. The chemical constituents have been identified by means of mass spectrometry and mass spectrometry- hyphenated techniques. Additionally, the group has also been involved in the production of protein-based nanoparticles to encapsulate bioactive phenolic compounds and to study the permeability through Caco-2 monolayers as models of the intestinal barrier. This work allowed the group to present 8 posters all having the same objective, that is, to study the bioactivity of phenolic compounds either from infusions or from leaves of plants or even recovering them from peels of fruits. These studies allowed to establish collaborations between CQB groups, such as the Interfacial Electrochemistry Group for AFM studies of nanoparticles and cell image, and the Inorganic and Theoretical Chemistry Group for docking studies of phenolic compounds with AChE. The group also studies the effect of phenolics on protein secondary structure (FT-IR) in collaboration with DMU, Leicester, UK, and the effect of drugs on living cells using FT-IR (ATR), in collaboration with King's College London, UK.

In the near future the group will focus on the effect of active compounds, drugs and natural products, in cancer mechanisms using mass spectrometry and mass spectrometry hyphenated techniques as well as FT-IR applied to living cells; on mechanisms of hypercholesterolemia and atherosclerosis treatments; on protein aggregation, associated with several age-related diseases; on new models to test new treatments, closer to the living organism (such as 3D cell culture); and also on the development of new methodologies for studying antioxidant and anti-inflammatory activities using cells *in situ*.

Acknowledgements

FCT for the programmatic financing reference number: UID/MULTI/00612/2013.

- [1] A. Proença-da-Cunha, A.P. Silva, O.R. Roque. Plantas medicinais em fitoterapia. Fundação Calouste Gulbenkian, Portugal, Lisboa, 2003.
- [2] M. Heinrich, H.L. Teoh, J. Ethnopharmacol. 92 (2004) 147-162.
- [3] D. Iqbal, M.S. Khan, S. Ahmad, A.K. Srivastava, Phytoter. Res. 28 (2014) 899-908.
- [4] D. Feng, L. Ohlsson, R.D. Duan, Lipids Health Dis. 9 (2010) 40

New generation of microextraction devices for trace analysis based on floating sampling technology

Alessandra H. Ide^a, Alexandra M. Fernandes^a, André M. Segurado^a, Samir M. Ahmad^a, Nuno R. Neng^a, José M. F. Nogueira^{a,*}

^a Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

*nogueira@fc.ul.pt

In the analysis of complex matrices, sample preparation is the most time consuming step. For this reason, the modern sample preparation approaches aim the miniaturization of the analytical devices and easy manipulation, as well as reduced sample volumes and absence (*i.e.* solventless) of toxic organic solvents in compliance with the green analytical chemistry principles.

“Bar adsorptive microextraction” (BA μ E) is a novel technique introduced by our group, which uses an approach based on the static mode and operates under the floating sampling technology. This novel technique uses an analytical device, light in weight in comparison to water density, simultaneously with a conventional Teflon magnetic stirring bar at the bottom of the sampling flask. When the sample matrix is rapidly spinning around due to centripetal force promoted by the magnetic bar, the analytical device stays under free-floating motion just below the centre of the formed vortex. Several coating phases have also been applied by this technique, including many types of activated carbons, polymers, ionic liquids, etc., in particular for trace analysis of the more polar compounds in aqueous media.

During a static process, the analytes migrate by diffusion from the sample bulk and are retained in a convenient sorbent phase, where the microextraction takes place. Then, the devices are removed from the samples, transferred into vials containing inserts with the stripping solvent, where the back-extraction stage takes place under sonication. Subsequently, the devices are removed from the inserts and, after evaporation and solvent-switch, the microextracts become ready for instrumental analysis.

Recently, our group have been involved on reducing the number of analytical steps required for the back-extraction stage. For this purpose, new generation BA μ E devices, smaller and more flexible than the original ones have been developed. The extraction and back-extraction stages still remain similar to the methodology described before, although the latter being performed in only-one liquid desorption step.

Another type of analytical devices proposed uses porous hollow fibres or polyurethane polymers, both having cylindrical geometry, in which are dipped with appropriate organic solvent for the microextraction of many classes of non-polar compounds in aqueous media.

By using these analytical approaches, trace analysis can be performed in matrices from several scientific areas, such as environment (pharmaceutical and personal care products, pesticides, endocrine disruptors, etc.), forensic (drugs of abuse, legal highs, steroid hormones, etc.), food (disinfection by-products of water, flavonoids, polyphenols, etc.) and many others.

Acknowledgements

The authors wish to thank “Fundação para a Ciência e Tecnologia” (Portugal) for the Post-Doc (SFRH/BPD/86071/2012) and PhD (SFRH/BD/107892/2015) grants and funding (UID/MULTI/00612/2013) and “Coordenação de Aperfeiçoamento de Pessoal de Nível Superior” (Brazil) for the PhD grant (CAPES BEX 0394-14-9).

Unraveling the role of TBCCD1 protein on cell size control: the regulation of cytoskeleton dynamics and cell junctions

Carolina Camelo^a, Catarina Peneda^a, Étienne Coyaud^b, Brian Raught^{b,c}, Ana I. Câmara^a, Bruno Carmona^{a,d}, Francisco Pinto^e, H. Susana Marinho^a, Helena Soares^{a,d,*}

^aCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^bPrincess Margaret Cancer Centre, University Health Network, Toronto, Canada

^cDepartment of Medical Biophysics, University of Toronto, Toronto, Canada

^dEscola Superior de Tecnologia da Saúde de Lisboa, IPL, 1990-096 Lisboa, Portugal

^e Instituto de Biosistemas e Ciências Integrativas, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

*mhsoares@fc.ul.pt

During their lifetime most cells maintain their size. There is increasing evidence showing that this process may be dynamic and that cells can adapt their size in response to external signals and changes in the environment [1], which strongly suggests that cell size is regulated. Both Hippo and IGF/PI3K/AKT/mTORC1 pathways have been described as being involved in cell size/growth control [1]. Interestingly, these pathways are in a cross-talk with others involved and/or dependent on cellular polarity [2]. Our group characterized a centrosomal protein, TBCCD1 (TBCC domain-containing human protein 1) which, when depleted in human retinal epithelial (RPE-1) cells, leads to an abnormal localization of the centrosome at the cell periphery accompanied by the fragmentation of the Golgi apparatus, resulting in the disruption of the intrinsic cell polarity axis “Nucleus-Centrosome-Golgi Apparatus”. Moreover, TBCCD1-depleted cells are larger, slower and have a lower efficiency in primary cilia assembly than control cells [3]. We identified the TBCCD1 interactome that showed that most of its partners are involved in cell polarity. Furthermore, most of them participate in the formation/maintenance of cell junctions, which are main regulators of cell polarity in epithelia and are upstream of pathways, like Hippo pathway. We also observed that TBCCD1 overexpression affects tubulin acetylation, which supports our results showing that some of the partners are involved in the regulation of the cytoskeleton dynamics, which may affect cell size. Therefore, it is tempting to hypothesize that the mechanisms involved in the establishment of intrinsic cell polarity may also directly/indirectly participate in the regulation of cell size.

Acknowledgements

Support for this work was provided by CQB's PEST UID/MULTI/00612/2013.

[1] Lloyd AC. The regulation of cell size. *Cell*. 2013 Sep 12; 154(6):1194-205.

[2] Soares H, Marinho HS, Real C, Antunes F. Cellular polarity in aging: role of redox regulation and nutrition. *Genes Nutr*. 2014 Jan;9(1):371.

[3] Gonçalves J, Nolasco S, Nascimento R, Lopez Fanarraga M, Zabala JC, Soares H. TBCCD1, a new centrosomal protein, is required for centrosome and Golgi apparatus positioning. *EMBO Rep*. 2010 Mar;11(3):194-200.

Exploratory chemistry of butyrylcholinesterase nucleoside-based inhibitors

Vasco Cachatra^{a*}, Ignazio Schino^{a,b}, Nicola Colabufio^b, Amélia P. Rauter^a


^a Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Ed C8, Piso 5, Campo Grande, 1749-016 Lisboa, Portugal

^b Biofordrug, Via Orabona, 4, Bari, Italy

*vmcachatra@fc.ul.pt

The increase in average life expectancy raises a number of challenges resulting in the development of new strategies pointing to a healthy ageing. As our organism begins to age, dementia is one of the many hurdles to overcome affecting 46.8 million people worldwide, with recent trends showing an increase in this number for the coming years.¹ Alzheimer's disease (AD) is the most common type of dementia but the complete disease ethiology is not known. While current treatments are based on the inhibition of acetylcholinesterase (AChE) or dual inhibition of AChE and butyrylcholinesterase (BChE) to restore acetylcholine levels on the brain, this effect is only significant in the early disease stages. BChE takes the role of AChE in hydrolysing acetylcholine in later stages,² so the search for selective inhibitors of BChE has the possibility to provide better treatment options and/or to have new insights onto AD ethiology.


Our research group has developed a new family of nucleosides that demonstrated a potent and selective inhibition of BChE.³ They have key structural features, namely a 2-acetamido-6-chloropurine N-linked to an unusual bicyclic sugar moiety (type I), or linked to glycosyl residues as depicted in structure-types II and III. Optimization of synthetic procedures for regio- and stereoselectivity toward potent activity and selectivity, synthesis of analogues and computational studies have been carried out to understand the binding mode and identify the key structural features required for the activity. These results will be disclosed and discussed.



Acknowledgements

The authors thank the European Commission for the support of the project entitled “Diagnostic and Drug Discovery Initiative for Alzheimer’s Disease”, Industry-Academia Partnerships and Pathways (IAPP), FP7-PEOPLE-2013-IAPP, GA 612347, 2014-2018. Fundação para a Ciência e a Tecnologia is also gratefully acknowledged for Vasco Cachatra PhD grant (SFRH/BD/90359/2012) and for the support of the project UID/MULTI/00612/2013.

- [1] Alzheimer's Disease International *World Alzheimer's report* **2015**, 1;
[2] I. R. Macdonald, K. Rockwood, E. Martin, S. Darvesh *J. Alzheimers Dis.* **2014**, *42*, 379;
[3] a) F. Marcelo, F. M. V. Silva, M. Goulart, J. Justino, P. Sinaÿ, Y. Blériot, A. P. Rauter, *Bioorg. Med. Chem.* **2009**, *17*(14) 5106; b) S. Schwarz, R. Csuk, A. P. Rauter, *Org. Biomol. Chem.* **2014**, *12*, 2446.



CQB-Day 2016

Flexible biomimetic/nanoconjugated platforms for sensitive immunosensing

Ana S. Viana

Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

**anaviana@fc.ul.pt*

The major scientific goal of this work is to develop simple and disruptive surface chemistry to push forward the field of biosensing. The preparation of biosensors, as the sandwich-type immunosensors, with improved sensitivity regarding expensive enzyme immunoassays (Enzyme linked immunosorbent assay, ELISA) [1], can follow two main strategies. One is to increase the capture of antigen by the primary antibody adsorbed on the surface (recognition layer), and the other is to optimize the signal amplification methodology to detect the target antigen by a secondary antibody. Herewith, we will tackle both aspects and present innovative yet simple approaches to enhance the detection (electrochemical and optical) of the biological event. It is worth noting that the sandwich-type immunosensors to be developed can be easily adapted and tailored to a trial of interest. One crucial demand is the biocompatibility of the recognition layer and in this context supported lipid bilayers (SLB) may provide an ideal matrix for antibody immobilization. However, the formation of flat, uniform and air stable SLB on conductive surfaces remains a challenge and most works fail to use biologically relevant lipid mixtures. Recently, the team was able to prepare robust and raft forming SLB on gold surfaces [2-4], enabling direct electrochemical monitoring and control. The expertise of the team on the synthesis of metallic and semiconducting nanoparticles and their functionalization [5], enable to use suitable surface chemistry for the direct bio-functionalization of nanoparticles, with antibodies. The biorecognition reactions will be transduced electrochemically and optically in a partnership with a Chinese Group (Institute of Mechanics, Beijing). In addition, a startup company, Lumisense Lda., will design and construct a microfluidic miniaturized immunosensor based on the developed platforms, directed to the detection of toxins in beverages.

Acknowledgements

Fundação para a Ciência e a Tecnologia (FCT) is acknowledged for financial support through projects: PEst 2015-2020 (UID/Multi/00612/2013), IF/00808/2013/CP1159/CT0003, IF2013 initiative (POPH, FSE) and PTDC/CTMNAN/0994/2014.

- [1] J Rusling, The Chemical Record 12 (2012) 164–176.
- [2] J. T. Marquês, R. F. M. de Almeida, A. S. Viana Soft Matter 8 (2012) 2007-2016.
- [3] J. T. Marquês, A. S. Viana, R. M. F. de Almeida, Langmuir 30 (2014) 12627-12637.
- [4] J. T. Marquês, I. Almeida, W. Liu, Y. Niu, R.F.M. de Almeida, G. Jin, A. S. Viana, Colloids and Surfaces B: Biointerfaces 136 (2015) 997-1003.
- [5] I. Almeida, S.G. Mendo, M.D. Carvalho, J.P. Correia, A.S. Viana, Electrochimica Acta 188 (2016) 1–12.

CpHMD-L simulations of pHLIP peptides: design of new tumor-targeted drug delivery systems

Diogo Vila-Viçosa^a, Pedro Reis^a, Vitor H. Teixeira^a, Maria J. Calhorda^a, António M. Baptista^b, Miguel Machuqueiro^a

^a Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, PT

^b Instituto de Tecnologia Química e Biológica António Xavier, Universidade Nova de Lisboa, Av. da República, 2780-157 Oeiras, PT

machuque@ciencias.ulisboa.pt

The pH (Low) Insertion Peptide (pHLIP) is a 36 amino acid peptide derived from bacteriorhodopsin [1] that targets tissues with acidic pH [2]. It simultaneously targets tumors, carries the cargo, and translocates it across the plasma membrane at low pH values [3]. At neutral pH, pHLIP is soluble as a monomer in water and associate with lipid bilayer surfaces largely as an unstructured peptide. Under acidic conditions, pHLIP inserts across a lipid bilayer with an apparent pK of 6, forming a transmembrane helix (Figure 1) [1]. The pH-dependent insertion process is coupled to the protonation of one or both of the Asp residues located in the transmembrane region of the peptide [1, 2].

Despite the extensive experimental studies on the mechanism and thermodynamics of pHLIP-membrane interactions [4], there is not enough information at the molecular level for a good understanding of the insertion phenomenon and its pre-requisites. These processes are usually only followed indirectly in typical biophysical experiments and cannot be modelled using conventional methodologies, like molecular dynamics or continuum electrostatics due to the inherent complexity associated with pH. Therefore, in this project, (PTDC/QEQCOM/5904/2014) we will employ computational methodologies (like our recently developed CpHMD-L method) to understand the molecular details of pH dependent peptide/membrane interaction and insertion.




Figure 1. The pHLIP peptide inserted in a lipid bilayer.

Acknowledgements

We acknowledge financial support from FCT projects PTDC/OE/COM/5904/2014 and UID/MULTI/00612/2013.

- [1] J. F. Hunt, P. Rath, K. J. Rothschild, D. M. Engelman, *Biochemistry*, 36:15177 (1997)
- [2] Y. K. Reshetnyak, O. A. Andreev, U. Lehnert, D. M. Engelman, *PNAS*, 103, 6460 (2006)
- [3] Yao, L., Danniels, J, Moshnikova, A, Kuznetsov, S, Ahmed, A, Engelman, DM, Reshetnyak, YK, Andreev, OA (2012) *PNAS*, 110, 465
- [4] D. Weerakkody, A. Moshinoka, M. S. Thakur, V. Moshinoka, J. Daniels, D. M. Engelman, O. A. Andreev, Y. R. Reshetnyak, *PNAS*, 9:5834 (2013).

Multifunctional luminescent spin labile hybrid materials

Paulo N. Martinho^{a*}, Liliana P. Ferreira^{b,c}, Maria de Deus Carvalho^a, Sara Realista^a, Nuno A. G. Bandeira^{a,d}, Janaína C. Almeida^a, Frederico F. Martins^a, Maria José Calhorda^a

^aCentro de Química e Bioquímica, DQB, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^bBioISI, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

^cDepartamento de Física, Faculdade de Ciências e Tecnologia, Universidade de Coimbra, 3004-516 Coimbra, Portugal

^dInstitute of Chemical Research of Catalonia (ICIQ), Avda. Països Catalans, 16- 43007 Tarragona, Spain


*pnmartinho@ciencias.ulisboa.pt

Data storage in binary systems operates by switching between two stable states under ambient conditions and information is written by application of a stimulus. Fabrication of typical data storage devices is made using a top-down approach where the size of magnetic domains has been constantly decreasing.[1] Reduction of size of magnetic domains is only possible until the super paramagnetic limit is reached, producing highly unstable devices.[2,3] To overcome the instability conferred by the superparamagnetic limit, scientists have been developing strategies using diverse and imaginative solutions. However, a new model of spin-based electronics, based on the orientation of individual electron spins to store binary information, offers the tantalizing possibility of non-volatility, increased data processing speed, decreased electric power consumption and increased integration densities.

Magnetic compounds with good potential for incorporation into spintronic materials include spin crossover transition metal compounds.[4] Research has been developed where bistable magnetic states in molecules are addressed mainly by application of temperature or light.

Engineering of multifunctional materials by combining both SCO and luminescence yields hybrid molecules and opens opportunities to develop a range of materials with applications in molecular electronics, nanomedicine and sensors technology.

This research proposal wishes to exploit the fact that is possible to combine more than one function in the same molecule forming hybrid molecular systems. This will be achieved by coordination of luminescent centres based on coinage metals to SCO molecules with functional acetylide groups. The research project here proposed consists of five major parts. The first part of the project is dedicated to the synthesis of both amphiphilic and unfunctionalised SCO Fe(III) molecules.



Acknowledgements

Support for this work was provided by Fundação para a Ciência e Tecnologia (FCT), Portugal, through projects UID/MULTI/00612/2013, PTDC/QEQ-QIN/3414/2014 and fellowship SFRH/BPD/73345/2010 (PNM).

- [1] Dietzel, A., Hard Disk Drives in Nanoelectronics and Information Technology Wiley-VCH: Berlin, 2005.
- [2] Charap, S. H., Pu-Ling, L., Yanjun, H., IEEE Trans. Magn., 1997, 33, 978-983.
- [3] Martín, J. I., Nogués, J., Liu, K., Vicent, J. L., Schuller, I. K., J. Magn. Magn. Mater., 2003, 256, 449-501.
- [4] Kahn, O., Martinez, C. J., Science, 1998, 279, 44-48.

Sphingolipid organization in the plasma membrane of *Saccharomyces cerevisiae*

Rodrigo F. M. de Almeida^{a,*}

^a Centro de Química e Bioquímica, Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

*e-mail corresponding author: rfalmeida@fc.ul.pt


Sphingolipids [SLs] are major lipid components of the plasma membrane [PM] of eukaryotes and their organization in the PM is critical for many cellular vital functions [1]. There is evidence implicating SLs in the mechanism of action and resistance to antifungal agents in clinical use [2, 3]. Moreover, fungal resistance to antifungals is an emerging public health problem [4], many important antifungal agents act on the PM and SLs are potential therapeutic targets in fungal infections [3, 5]. Despite its recognized importance, SL organization in the PM of fungi remains poorly understood, even in *Saccharomyces cerevisiae*, and biophysical studies with yeast SLs are almost inexistent. Thus, the goal of this project is to define the structural features and molecular interactions of SLs which are crucial for the organization of the PM in *S. cerevisiae*, establishing the biophysical principles governing SLs role in the mode of action of antifungals and fungal resistance.

Acknowledgements

Fundação para a Ciência e a Tecnologia (FCT), Portugal is acknowledged for PTDC/BBB-BQB/6071/2014, UID/Multi/00612/2013, and IF2012 initiative (P.O.P.H., F.S.E.).


References

1. J. C. Holthuis, T. Pomorski, R. J. Raggars, H. Sprong, G. Van Meer, *Physiol Rev.* 81 (2001) 1689-1723.
2. I.E. François, A. Bink, J. Vandercappellen, K.R. Ayscough, A. Toulmay, R. Schneiter, E. van Gyseghem, G. Van den Mooter, M. Borgers, D. Vandenbosch, T. Coenye, B.P.A. Cammue, K. Thevissen, *J. Biol. Chem.* 284 (2009) 32680-32685.
3. K. Mukhopadhyay, T. Prasad, P. Saini, T.J. Pucadyil, A. Chattopadhyay, R. Prasad, *Antimicrob. Agents Chemother.* 48 (2004) 1778-1787.
4. A. Chakrabarti., *Regional Health Forum* 15 (2011) 97-103.
5. G.B. Wells, R.C. Dickson, R.L. Lester, *J. Bacteriol.* 178 (1996) 6223-6226.



Science at CQB

Poster Communications



Surfaces modified by bio-inspired molecules for molecular oxygen reduction catalysis

Rui S Sampaio*, Jorge P Correia

Centro de Química e Bioquímica, DQB, Faculdade de Ciências, Universidade de Lisboa, Portugal

**kruder@hotmail.com*

The oxygen reduction reaction (ORR) is a sluggish reaction over any metal substrate, limiting the kinetics of the electrochemical devices using the molecular oxygen as oxidising agent [1]. The pursuit of new reliable substitutes of the conventional catalysts will lead to the preparation of materials with increased electrocatalytic activity and economically interesting in order to push forward the market of such electrochemical devices.

The purpose of this work is to develop a high performance cathodic catalyst for the ORR based on bio-inspired molecules (synthetic metalloporphyrins and Vitamin B12), that display a high intrinsic catalytic potential, as widely recognized by the scientific community [2-3]. To achieve this goal, porphyrin moieties were electrochemically immobilized on electrode surfaces. Cyclic voltammetry was employed to understand the electrochemical behaviour of the metal complexes in solutions over different substrates as well as that of the matrices assembled on the electrode surfaces. The catalyst was electrochemically deposited using two different approaches; a) by continuous potential cycling of the substrate in solutions containing the coordination compound and b) incorporation of the catalyst into a conducting polymer matrix during its synthesis using potentiodynamic and potentiostatic techniques. The electrochemical behaviour of the modified electrodes was characterized by cyclic voltammetry and their morphology by atomic force microscopy (AFM) and x-ray photoelectron spectroscopy (XPS). The dielectric constants and thickness of the thin films were assessed by multiangle *ex-situ* ellipsometry. The electrocatalytic performance of such materials towards the ORR was evaluated in neutral and acid media. To investigate the mechanism and kinetics of ORR at the different modified electrodes with the porphyrins it was used the rotating ring disk electrode technique.

Acknowledgements

This work is funded by FCT (Fundação para a Ciência e a Tecnologia) through the project UID/MULTI/00612/2013

[1] Perla B Balbuena and Venkat Subramanian, *Theory and Experiment in Electrocatalysis*, Springer, 2010

[2] R Fraga and JP Correia, *Electrochimica Acta*, 2005, 50, 1653-1659

[3] Karl M Kadish, *Handbook of Porphyrin Science*, Elsevier Science, 2003

Macrophage type lectin -galactose ligands: chemical synthesis

João Manuel de Barros^{a*}, Filipa Marcelo^b, Amélia P. Rauter^a

^a*Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Ed C8, Piso 5, Campo Grande, 1749-016 Lisboa, Portugal*

^b*Unidade UCIBIO, Departamento de Química, Faculdade de Ciências e Tecnologias, Universidade Nova de Lisboa, Quinta da Torre, 2829 -516 Caparica, Portugal*

*jmvbarros@fc.ul.pt

Filoviruses, like Marburg and Ebola viruses, cause hemorrhagic diseases in humans and nonhuman primates with high rate of mortality. As other C-type lectins like DC-SIGN (Dendritic Cell-Specific Intercellular adhesion molecule-3-Grabbing Non-integrin), the macrophage galactose C-type lectin (MGL) binds to Ebola or Marburg virus glycoprotein (GP). The interaction is formed via the GalNAc residue directly linked to Ser/Thr side chains on the GP of the virus and promotes viral infectivity *in vitro*, by enhancing viral attachment to cellular entry receptors [1]. Furthermore, MGL is present on cells known to be major targets of filoviruses (i.e., macrophages and dendritic cells), suggesting a role for this receptor in viral replication *in vivo*. In this context, it was recently unravelled the interactions of GalNAc, Gal and tumor-associated MUC1 glycopeptides and MGL[2].

In this project we are synthesizing mimetics of GalNAc, for example, phenylselenenyl galactosides bearing an imide functionality at position 2, that could better bind to MGL receptor than GalNAc itself, therefore competing to the interaction between MGL and the GP of Ebola or Marburg filovirus.

NMR spectroscopy has demonstrated its suitability to provide structural and dynamics information, at an atomic level, of carbohydrate-protein complexes [3]. The interaction of GalNAc mimetics and MGL lectin receptor will be monitored by NMR binding studies in combination with molecular modelling simulations, to clarify the key interacting features of the new molecules. NMR data of the new glycomimetics, including affinity data, will also be compared to those of the natural GalNAc ligand.

Acknowledgements

The authors thank Fundação para a Ciência e a Tecnologia for the support of the project UID/MULTI/00612/2013.

References

- [1] Takada, A., Fujioka, K., Tsuiji, M., Morikawa, A., Higashi, N., Ebihara, H., Kobasa, D., Feldmann, H., Irimura, T., Kawaoka, Y., (2004) Human Macrophage C-Type Lectin Specific for Galactose and N-Acetylgalactosamine Promotes Filovirus Entry, *J. Virology*, 78, 2943-2947;
- [2] Marcelo F., Garcia-Martin F., Matsushita T., Sardinha J., Coelho H., Oude-Vrielink A., Koller C., André S., Cabrita E. J., Gabius H.-J., Nishimura S.-I., Jiménez-Barbero J., Cañada F. J., (2014) Delineating binding modes of Gal/GalNAc and structural elements of the molecular recognition of tumor-associated mucin glycopeptides by the human macrophage galactose-type lectin, *Chem. Eur. J.* 20, 16147-16155. VIP Paper.
- [3] Fernández-Alonso M. C., Díaz D., Berbis M. A., Marcelo F., Jiménez-Barbero J., (2012) Protein-carbohydrate interactions. From molecular recognition to drug design. *Curr. Protein Pept. Sci.*, 13, 816-8

Extraction and analysis of sphingolipids from *Saccharomyces cerevisiae* wild type cells and in a sphingolipid biosynthetic mutant

Filipe M. Moreira*, Catarina A. Antunes, H. Susana Marinho, Rodrigo F.M. de Almeida

Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

**fc42633@alunos.fc.ul.pt*

The yeast *Saccharomyces cerevisiae* exhibits a composition in ergosterol and sphingolipids in the plasma membrane similar to pathogenic fungi, being ergosterol the main target of chemical agents against these fungi¹.

On another hand, *S. cerevisiae* cells, such as *ipt1Δ* deletion mutant, which are unable to synthesize the sphingolipid mannosyl-diinositolphosphorylceramide, M(IP)₂C, show greater resistance to antifungal agents, such as miconazole² and nystatin³. This strain has ergosterol content similar to the wild type strain. Therefore, the mechanisms of this resistance may involve sphingolipid domains, which in yeast may be ergosterol depleted⁴.

For the reasons above, it is important to study the sphingolipid influence in the plasma membrane organization of *S. cerevisiae*. To this end, we intend to isolate and perform a thorough biophysical characterization of each complex sphingolipid class present in the plasma membrane of that organism separately. Before that, it is however, necessary that the sphingolipid extraction from yeast cells is optimized.

After obtaining the total lipid extracts, the lipids have to undergo mild alkaline hydrolysis to hydrolyze the glycerophospholipids, but not the sphingolipids. The sphingolipid extracts thus obtained are then analyzed by thin layer chromatography. If good separation is reached, a purified sample of each sphingolipid can be extracted from the respective band.

Several optimization steps using model lipid mixtures were needed to reach the most suitable time of hydrolysis, since longer times can hydrolyze not only glycerophospholipids but also sphingolipids and shorter times may hydrolyze incompletely the glycerophospholipids.

Regarding the yeast lipid extracts, obtained using the Fölch method, it was only possible to identify unequivocally the sphingolipid mannosyl-inositolphosphorylceramide (MIPC), in both wt and *ipt1Δ* extracts. The impossibility to observe M(IP)₂C may be due to low extraction efficiency of this highly polar lipid and consequently new methods are required to overcome this obstacle, which are now being attempted.

It was possible to observe no differences when it comes to the contents in glycerophospholipids and ergosterol between the two strains. However, an increase in the amount of MIPC in *ipt1Δ* extracts was perceptible, as expected.

Acknowledgments

Fundação para a Ciência e a Tecnologia (FCT), Portugal is acknowledged for PTDC/BBB-BQB/6071/2014, UID/Multi/00612/2013, and IF2012 initiative (POPH, Fundo Social Europeu).

1. Castelli, M. V. *et al.* Novel antifungal agents: a patent review (2011 - present). *Expert Opin. Ther. Pat.* **24**, 323–338 (2014).
2. François, I. E. J. A. *et al.* Membrane rafts are involved in intracellular miconazole accumulation in yeast cells. *J. Biol. Chem.* **284**, 32680–32685 (2009).
3. Leber, A., Fischer, P., Schneiter, R., Kohlwein, S. D. & Daum, G. The yeast mic2 mutant is defective in the formation of mannosyl-diinositolphosphorylceramide. *FEBS Lett.* **411**, 211–214 (1997).
4. Aresta-Branco, F. *et al.* Gel domains in the plasma membrane of *Saccharomyces cerevisiae*: highly ordered, ergosterol-free, and sphingolipid-enriched lipid rafts. *J. Biol. Chem.* **286**, 5043–5054 (2011).

Polydopamine films modified with Laccase for electrochemical biosensing

Luís C. Almeida*, Jorge P. Correia, Ana S. Viana

^aCiências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

**fc42424@alunos.fc.ul.pt*

Laccases (Lac) are versatile oxidoreductases known by their wide applicability in wastewater treatments, textile and paper industries, delignification processes, biofuel cells and biosensors^[1]. Despite their excellent catalytic properties towards several substrates, more reliable immobilization strategies are needed to stabilize the enzyme in a solid surface and to allow their reusability^[2]. Common electrode modification approaches, such as self-assembled monolayers, are not applicable to all surfaces and may involve multi-step procedures^[3]. In contrast, the bio-inspired polydopamine (PDA) films can spontaneously grow on a variety of surfaces, creating a biocompatible interface with interesting adhesive properties^[4]. This highly functional films enriched with quinone groups are able to covalently bind target biomolecules, such as Laccases, through a Schiff base reaction or Michael type additions^[5]. In this work we have modified glassy carbon and graphite electrodes with spontaneously formed or electrochemically synthesized polydopamine films for the covalent immobilization of two types of commercial Laccases: purified powder and cocktail booster form (Figure 1). Several PDA films with different thicknesses were grown on glassy carbon and their electrochemical properties were evaluated by cyclic voltammetry. Ellipsometric measurements allow to correlate the film thicknesses with time deposition or polymerization potential cycles. Contact angle goniometry disclose the hydrophilic nature of PDA, whereas morphological information was provided by AFM onto highly oriented pyrolytic graphite. This multitude of techniques revealed the properties of both polydopamine types and allow to maximize the catalytic performances of carbon/PDA/Lac biosensors, that were evaluated afterwards by cyclic voltammetry and chronoamperometry towards different substrates.





Figure 1. Two-step methodology used for the carbon/PDA/enzyme preparation.

Acknowledgements

The authors acknowledge Professors A. Morana and F. La Cara for providing the Laccase Booster. This work was financially supported by FCT (Fundação para a Ciência e a Tecnologia): through projects: PEst 2015-2020-UID/Multi/00612/2013, IF/00808/201.

- [1] S.R. Couto, J.L.T. Herrena, *Biotechnology Advances*, 24 (2006) 500-513.
- [2] C. Mateo, J.M. Palomo, G. Fernandez-Lorente, J.M. Guisan, R. Fernandez-Lafuente, *Enzyme and Microbial Technology* 40 (2007) 1451–1463.
- [3] I. Almeida, V.C. Ferreira, M.F. Montemor, L.M. Abrantes, A.S. Viana, *Electrochimica Acta*, 30 (2012) 311-320.
- [4] Y. Lui, K. Ai, L. Lu, *Chemical Reviews*, 114 (2014) 5077-5115.
- [5] E. Faurea, C. Falentin-Daudréa, C. Jérômea, J. Lyskawa, D. Fournier, P. Woisel, C. Detrembleur, *Progress in Polymer Science*, 38 (2013) 236-270.



Synthesis of new antibiotic glycosides and computational studies on their interaction with model lipid bilayers

Rafael Nunes*, João Pais, Diogo Vila-Viçosa, Miguel Machuqueiro, Amélia P. Rauter

Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

*rsnunes@fc.ul.pt

The spread of bacterial resistance to the available therapeutics is considered a major public health threat to contemporary societies. In the search for innovative sugar-based antibiotics with new mechanisms of action, our research group has successfully generated a family of surfactant glycosides exhibiting potent antibacterial activity against several pathogens including *Bacillus anthracis*, which is considered a bioterrorism agent [1]. In particular, a series of alkyl 2-deoxy- and 2,6-dideoxy-*arabino*-hexopyranosides were accessed by reaction of fatty alcohols with protected glycals in the presence of triphenylphosphane hydrobromide, an efficient procedure that stereoselectively delivers the bioactive α -glycosides in high yields. Their antibacterial activity is modulated by the deoxygenation pattern of the sugar moiety, and preliminary mechanistic studies indicate that these molecules act through the destabilization of bacterial cell membranes.

With the intent to further rationalize the structural key features for glycoside bioactivity, we designed and synthesized a small library of analogues related to the most active compound, including 2-deoxyglycosides derived from pentoses, 2-fluorinated analogues and glycosides deoxygenated at 6-position of the sugar, which were screened for their potential antibacterial activity. In addition, we applied molecular dynamics simulations to study the interactions of these molecules, as well as the aggregates they form in solution, with model phospholipid membranes. Particularly, we modelled the partitioning of glycosides into the lipid bilayer at the interface and analyzed the effect of these events on membrane thermotropic properties. This computational study provided important insights into their mechanism of action which will also be disclosed.

Acknowledgements

Support for this work was provided by Fundação para a Ciência e a Tecnologia (project UID/MULTI/00612/2013), QREN-COMPETE (project n°. 21547) and COST (action CMST/COST-Action/CM1102). João Pais acknowledges financial support from Fundação para a Ciência e a Tecnologia (fellowship SFRH/BDE/51957/2012).

- [1] A. Martins, M. S. Santos, C. Dias, P. Serra, V. Cachatra, J. Pais, J. Caio, V. H. Teixeira, M. Machuqueiro, M. S. Silva, A. Pelerito, J. Justino, M. Goulart, F. V. Silva, A. P. Rauter, *Eur. J. Org. Chem.* 8 (2013) 1448-1459.

Comparative study between Grunwald-Winstein and TAKA parameters used for the study of solvent effects

Ruben Elvas- Leitão^{a,b,*}, Ricardo J. Nunes^a, Filomena Martins^a

^aCentro de Química e Bioquímica, Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Ed. C8, Campo Grande 1749-016 Lisboa, Portugal

^bÁrea Departamental de Engenharia Química, Instituto Superior de Engenharia de Lisboa, Instituto Politécnico de Lisboa, R. Conselheiro Emídio Navarro, 1, 1959-007 Lisboa, Portugal

*rleitao@deq.isel.pt

Grunwald-Winstein plots [1] and TAKA model equation [2] have been used over time to rationalize solvent effects in solvolysis reactions. To our knowledge, very little has been done to correlate both sets of parameters to understand if they deliver the same type of mechanistic information.

One reason for this might be the fact that there are very few, disperse and inconsistent data in the literature regarding solvatochromic parameters for the typical Grunwald-Winstein (G-W) mixtures which involve mainly the following solvent systems: ethanol/water, methanol/water, acetone/water, trifluoroethanol/water and trifluoroethanol/ethanol.

In this work, Kamlet-Taft parameters (π^* , α and β) have been determined at 298.15 K on the basis of the spectroscopic shifts of five solvatochromic probes (betaine (30), 4-nitrophenol, 4-nitroanisole, 4-nitroaniline and *N,N*-dimethyl-4-nitroaniline), for the referred G-W solvent mixtures, in order to establish a coherent matrix of solvatochromic parameters.

Correlations between G-W *N* and *Y* scales and TAKA π^* , α and β scales have been established showing that the former can be in fact expressed as linear combinations of the TAKA parameters, thus confirming the unsuspected similarities between both approaches.

Acknowledgements

The authors are grateful to Fundação para a Ciência e a Tecnologia, Portugal, for financial support under project UID/MULTI/00612/2013.

[1] E. Grunwald, S. Winstein, J. Amer. Chem. Soc. 73 (1951) 2700-2707.

[2] R. W. Taft, J. L. M. Abboud, M. J. Kamlet, M. H. Abraham, J. Sol. Chem. 14 (1985) 153-186.

Synthesis and hepatotoxicity of psychoactive cathinones

Soraia Bronze^{a,*}, Rita Pacheco^{a,b}, Susana Santos^a, Helena Gaspar^a

^a Centro de Química e Bioquímica, Faculdade de Ciências da Universidade de Lisboa, Campo Grande, 1749-016 Lisboa,

^b Área Departamental de Engenharia Química, Instituto Superior de Engenharia de Lisboa, Av. Conselheiro Emídio Navarro, Lisboa, Portugal

*fc42398@alunos.fc.ul.pt

In the last few years, there has been an uprising of new psychoactive substances (NPS) available in “smartshops” and over the Internet. NPS are “new narcotic or psychotropic drugs that are not listed in the Single Convention on Narcotic Drugs of 1961 or the Convention on Psychotropic Substances of 1971, but which may constitute a public health threat comparable to the one posed by substances listed in those conventions”. Since 2005 more than 560 novel NPS have been reported to the Early Warning System (EWS) in Europe, of which 103 are synthetic cathinones being 22 firstly reported in 2015 and 31 in 2014, rendering this class of compounds one of the widest category of NPS known to date [1]. Cathinones are β -keto phenethylamines derivatives, structural analogues of the natural-occurring cathinone, the psychoactive stimulant found in the khat plant.

The neuropharmacological action of cathinones is similar to those of more traditional illicit psychostimulants, such as D-amphetamine, methamphetamine and MDMA, and is related with the increase of synaptic concentration of the monoamines dopamine, norepinephrine and serotonin, either by inhibiting the corresponding transporters (SERT, NET and DAT) or by increasing the pre-synaptic release of the neurotransmitters. In addition it is now known that different synthetic cathinones have diverse mechanisms of action, which strongly depends on their structure [2].

An increasing number of cases of severe liver injury have been reported, as a consequence of khat use or abuse, particularly in countries where *Catha edulis* is legal[3]. Recently a study on the *in vitro* hepatotoxic effects of some individual synthetic cathinones, namely methylone, pentedrone, 4-methylethcathinone (4-MEC) and 3,4-methylenedioxypyrovalerone (MDPV) revealed that MDPV was the more toxic compound[4].

The aim of this work was to evaluate, in a comprehensive way, the hepatotoxicity of a series of synthetic cathinones: methcathinone, buphedrone, pentedrone, mephedrone (4-MMC), 4-MEC, α -PPP α -PBP and α -PVP (Fig. 1), enabling the identification of structural determinants of hepatotoxicity.

Eight cathinones were synthesized and evaluated against human hepatoma cell line Hep G2. Results on *in vitro* hepatotoxicity seem to disclose that cathinones hepatotoxicity varies significantly with their structure. The amine connected to the carbonyl α -carbon seems to be irrelevant to the toxicity, which is dependent of the length of the ketone chain and on methyl substitution of the aromatic ring. The more toxic cathinone was mephedrone with an EC 50= 1.38mM (used as control), followed by 4-MEC with an EC 50= 1.50 mM.


	Methcathinone	R ₁ =R ₃ = H; R ₂ =R ₄ = CH ₃
	Bufedrona	R ₁ =R ₃ = H; R ₂ = CH ₂ CH ₃ ; R ₄ = CH ₃
	Pentedrona	R ₁ =R ₃ = H; R ₂ = CH ₂ CH ₂ CH ₃ ; R ₄ = CH ₃
	α -PPP	R ₁ = H; R ₂ = CH ₃ ; R ₃ ,R ₄ = Pyrrolidine
	α -PBP	R ₁ = H; R ₂ = CH ₂ CH ₃ ; R ₃ ,R ₄ = Pyrrolidine
	α -PVP	R ₁ = H; R ₂ = CH ₂ CH ₂ CH ₃ ; R ₃ ,R ₄ = Pyrrolidine
	4-MMC	R ₁ = R ₂ = R ₄ = CH ₃ ; R ₃ = H
	4-MEC	R ₁ = R ₂ = CH ₃ ; R ₃ = H; R ₄ = CH ₂ CH ₃

Figure 1. General structure of the tested cathinones

Acknowledgments

FCT for the programmatic financing reference number: UID/MULTI/00612/2013

- [1] European Monitoring Centre for Drugs and Drug Addiction and Europol (2016), EU Drug Markets Report: In-Depth Analysis, EMCDDA–Europol Joint publications, Publications Office of the European Union, Luxembourg.
- [2] L.D. Simmler, T.A. Buser, M. Donzelli, Y. Schramm, L.H. Dieu, J. Huwyler, S. Chaboz, M.C. Hoener, M.E. Liechti, Brit J Pharmacol, 168 (2013) 458-470.
- [3] F. Pantano, R. Tittarelli, G. Mannocchi, et. Al. Int. J. Mol. Sci. (2016) 17 580
- [4] A.M. Araújo, M.J. Valente, M. Carvalho, D.D. Silva, H. Gaspar, F. Carvalho, M.L. Bastos, P.G. Pinho, Arch Toxicol (2015) 89 757–771.

Thermodynamic study of aqueous mixtures of 3-butoxypropan-1-amine

Ângela F. S. Santos^a, Maria Luisa C. J. Moita^{b,*}, Luis C. S. Nobre^{a,b}, Isabel M. S. Lampreia^a

^aCentro de Química Estrutural, Faculdade de Ciências, Universidade de Lisboa, Ed. C8, Campo Grande 1749-016 Lisboa, Portugal

^bCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Ed. C8, Campo Grande 1749-016 Lisboa, Portugal

* mlmoita@ciencias.ul.pt

Alkoxyamines are versatile compounds with great affinity for water since they have in their structure the alkoxy and amine functional groups. They have been used in industry for the immobilization of various compounds such as biomolecules by means of condensation reactions with aldehydes or ketones forming oxime bonds, which increase their stability [1]. Additionally, the strong electron-donor functional groups of alkoxyamines have been largely used in polymer chemistry to produce sequence-controlled polymerization and cyclization reactions [2]. Pursuing the volumetric studies [3-5] dedicated to improve comprehension embracing molecular interactions in binary systems water + amphiphilic molecules, the density, ρ , and sound speed, u , in water (1) + 3-butoxypropan-1-amine (BPA)(2), spanning the temperature range (283.15 to 303.15) K and over the entire composition range were determined. A comparison of the present results with those obtained for the system water + 3-ethoxypropan-1-amine (EPA) is made addressing the changes produced by the hydrophobic characteristics of the molecules. These studies are undertaken in order to clarify the influence of structure, chain length and type of hydrophilic groups on the aggregation and hydration schemes.

Derived thermodynamic quantities revealing changes of molecular aggregation with temperature and composition across the whole composition range. The profile of the curves of partial molar properties, which entails first order derivatives of the molar volumes and molar isentropic compressions, shows at least three different regimens of molecular aggregation (Figure 1).




Figure 1 Excess partial molar isentropic compressions

Acknowledgements

Support for this work was provided by FCT under projects UID/MULTI/00612/2013, PEst-OE/QUI/UI0612/2013 and PEst-OE/QUI/UI100/2013.

- [1] M.R. Hill, S. Mukherjee, P.J. Costanzo, B.S. Sumerlin, Polym. Chem. 3 (2012) 1758–1762.
- [2] C. St. Thomas, H. Maldonado-Textle, A. Rockenbauer, L. Korecz, N. Nagy, R. Guerrero-Santos, J. Polym. Sci. Part A: Polym. Chem. 50 (2012) 2944–2956.
- [3] E. Bulemela, P.R. Tremaine, J. Phys. Chem. B, 112 (2008) 5626–5645.
- [4] M.-L.C.J. Moita, L.M.V. Pinheiro, Â.F.S. Santos, I.M.S. Lampreia, J. Chem. Eng. Data 57 (2012) 2290–2295.
- [5] L.M.V. Pinheiro, M.-L.C.J. Moita, Â.F.S. Santos, I.M.S. Lampreia, J. Chem. Thermodyn. 64 (2013) 93–99.

The spin crossover profile in iron(III) Schiff-base compounds: halogen influence

F. F. Martins,^{a*} A. I. Vicente,^a L. P. Ferreira,^{b,c} M. D. Carvalho,^a V. H. N. Rodrigues,^c M. J. Calhorda,^a P. N. Martinho^a

^aCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

^bBioISI, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

^cDepartamento de Física, Faculdade de Ciências e Tecnologia, Universidade de Coimbra, 3004-516 Coimbra, Portugal

*fred.martins94@gmail.com

Technological advances have been pushing the limits of chemistry for the last few years towards creating more efficient and multifunctional molecules and materials. A phenomenon that shows great promise in molecular electronics is spin crossover (SCO).[1] This switching can be harnessed to develop materials with a wide range of possible applications such as memory or sensing nano-devices.[2] Halogen derivatized SCO molecules are of great interest as they can interact with neighboring molecules through either halogen or hydrogen bonds and additionally they can be modified through substitution or coupling reactions conferring additional properties and high versatility to the SCO molecules.[3,4]

Here we report the synthesis and characterization of halogen derivatized SCO compounds with an Fe(III) metallic center coordinated to tridentate (N2O) Schiff-base ligands. We have found that all compounds exhibit SCO with profiles ranging from gradual to abrupt with hysteresis and a detailed study on the halogen influence on these is also being carried out.




Figure 1 – Sample magnetic profile, x-ray crystal structure and picture of synthesized compounds

Acknowledgements

The authors thank Fundação para a Ciência e Tecnologia for financial support UID/MULTI/00612/2013 and PTDC/QEQ-QIN/3414/2014. PNM thanks FCT for financial support (SFRH/BPD/73345/2010).

[1] Gamez, P.; Costa, J. S.; Quesada, M.; Aromí, G., Dalton Trans. 2009, 38, 7845.

[2] Létard, J.; J. Mater. Chem. 2006, 16, 2550.

[3] Devid, E. J.; Martinho, P. N.; Kamalakar, M. V.; Šalitroš, I.; Prendergast, Ú.; Dayen, J. F.; Meded, V.; Lemma, T.; González-Prieto, R.; Evers, F.; Keyes, T. E.; Ruben, M.; Doudin, B.; van der Molen, S. J., ACS nano 2015, 9, 4496.

[4] Gaspar, A. B.; Ksenofontov, V.; Seredyuk, M.; Gütllich, P., Coord. Chem. Rev. 2005, 249, 2661.

The ability of sterols to form liquid ordered domains links membrane biophysical properties to biological processes

Catarina A. C. Antunes^{a,*}, Alena Khmelinskaia^{a,b}, André E.P. Bastos^a, Gerson M. da S. Lobo^a, Rui Malhó^c, M. L. Cyrne^a, H. Susana Marinho^a, Rodrigo F. M. de Almeida^a

^a Centro de Química e Bioquímica, Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

^b Max-Planck Institut for Biochemistry, Department of Cellular and Molecular Biophysics, 82152 Planegg, Germany

^c BioISI, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

* e-mail corresponding author: catarina.alves.antunes@gmail.com

The major sterol present in the plasma membrane of fungi and mammalian cells – ergosterol (erg) and cholesterol (chol), respectively – present small structural differences, given that they share a common metabolic precursor, zymosterol (zym), close to the final step of their biosynthetic route. In this work, we have studied the importance of these structural differences for the formation and properties of sterol-enriched plasma membrane microdomains (lipid rafts) in eukaryotes. To better understand sterol evolution *in vivo*, liposomes containing chol, erg or zym were prepared and different biophysical properties were assessed *in vitro*.

Emission spectra of di-4-ANEPPS, a dye sensitive to sterol-phospholipid interactions, show that chol and erg interact differently with saturated (DPPC, gel phase) and unsaturated (POPC, liquid disordered *l_d* phase) glycerophospholipids, whereas zym interacts similarly with both. The stronger interaction of erg and chol with saturated phospholipids led to the formation of liquid ordered (*l_o*) raft-like domains, while zym showed no capacity to form them. These results were consistent with confocal fluorescence imaging of microdomains in giant unilamellar vesicles

The study of sterol effect on lipid bilayer passive permeability to water and to univalent cations (H^+/K^+ exchange), using binary and ternary mixtures, also corroborated the behavior described above. The presence of erg or chol in DPPC bilayers markedly increased the passive permeability of the membrane to water, concomitant with a change from gel to *l_o* phase. Zym, however, had only a marginal effect. Furthermore, in lipid mixtures containing both saturated and unsaturated phospholipids, zym failed to induce the decrease of passive permeability observed for the raft-forming sterols. These results corroborate that the addition of zym does not change the gel/*l_d* coexistence, a situation where the domain interface packing defects are responsible for most of the observed permeation, whereas the presence of erg and chol shifts this regime to a coexistence of *l_o*-*l_d* domains, with much less interfacial defects.

Additionally, we determined the total expression and distribution in the plasma membrane of Can1p-GFP, a marker of erg-enriched microdomains, in wild-type (wt) and *erg6Δ* yeast cells. The results revealed a similar heterogeneity degree at the plasma membrane but a higher level of expression, possibly to compensate for traffic defects at the level of the Golgi complex also registered in *erg6Δ* cells.

Taken together, our *in vitro* evidence for the inability of zym to promote the formation of *l_o*-like domains provide a biophysical foundation for the observed increased permeability to small water-soluble dyes and stress sensitivity as well as defective lipid-raft dependent traffic observed in zym-enriched *erg6Δ S. cerevisiae* cells, when compared to wt cells. Hence, this work supports the hypothesis that lipid raft formation in fungi and mammalian cellular membranes is a convergent evolutionary trait that assures the establishment of the biophysical properties necessary for biological membrane responses.

Acknowledgements

Fundação para a Ciência e a Tecnologia (FCT), Portugal is acknowledged for grants PTDC/BBB-BQB/6071/2014 and UID/Multi/00612/2013, IF2012 initiative (P.O.P.H., F.S.E.), and Ph.D. fellowship SFRH/BD/88199/2012 to A.E.P.B. We thank Prof. W. Tanner (University of Regensburg, Germany) for the kind gift of Plasmid Ylp211, which was transformed into BY4741 (wt) strain to obtain strain LC4 expressing CAN1-GFP, and Prof. A. Herrmann and Dr. S. Scolari (Humboldt University, Berlin, Germany) for the previous FLIM studies.

[VCl₃{η³-HC(pz)₃}]@CNT as catalysts for the microwave-assisted oxidation of xylenes

Jiawei Wang^{a*}, Luísa M.D.R.S. Martins^{a,b}, Ana P.C. Ribeiro^a, Marta Saraiva^d, Sónia A.C. Carabineiro^c, José L. Figueiredo^c, Armando J.L. Pombeiro^a

^a CQE, Complexo I, IST, UL, Av. Rovisco Pais, 1049-001 Lisboa, Portugal


^b ADEQ, ISEL, IPL, Rua Conselheiro Emídio Navarro, 1959-007 Lisboa, Portugal

^c Lab. Catálise e Materiais, LA LSRE-LCM, Fac. Engenharia, Univ. Porto, 4200-465 Portugal

^d Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal


*jiawei.wang@tecnico.ulisboa.pt

Oxidation is a widely used synthetic route for a large range of chemicals. Important oxidation substrates are alkyl aromatics, such as xylenes [1]. However, the oxidation of xylenes with peroxides, as well as under heterogeneous conditions, is scarce. This prompted us to consider the application of a catalyst able to oxidize alkanes in homogeneous conditions [2] to the heterogeneous oxidation of different xylenes. This was carried out through heterogenisation of a homogeneous complex on a heterogeneous carbon support.



Scheme 1 – Vanadium complex [1].

The C-scorpionate trichloro-vanadium (III) complex [VCl₃{η³-HC(pz)₃}] (pz = pyrazolyl) [1], shown in Scheme 1, was supported on multiwalled carbon nanotubes (CNT) and characterized by SEM, TEM, BET and XPS. The heterogenized complex was used as catalyst for the MW-assisted solvent free oxidation of xylenes with *tert*-butyl hydroperoxide (TBHP), under mild conditions (see Scheme 2 for *m*-xylene, where the main products are *m*-methylbenzylalcohol, *m*-methylbenzaldehyde and *m*-methylbenzoic acid). The effects of different reaction parameters are discussed.



Scheme 2 – Oxidation of xylene under solvent free conditions.

Acknowledgements

FCT for projects UID/QUI/00100/2013, PTDC/QEQ-ERQ/1648/2014 and PTDC/QEQ-QIN/3967/2014. The LA LSRE-LCM was co-financed by FCT and FEDER through COMPETE 2020 (Project UID/EQU/50020/2013 - POCI-01-0145-FEDER-006984). SACC acknowledges Investigador FCT program (IF/01381/2013/ CP1160/CT0007). APCR and JW also acknowledge FCT for SFRH/BPD/90883/2012 and Ph.D. CATSUS PD/BD/114397/2016 grants, respectively. MSS and JW also thank Fundação para a Ciência e Tecnologia for financial support (PEST UID /MULTI/00612/2013). MSS thanks FCT for Grant SFRH / BPD / 88082 / 2012.

[1] R. A. F. Tomás, J. C. M. Bordado, J. F. P. Gomes, Chem. Rev. 113 (2013) 7421–7469.

[2] T. F. S. Silva, E. C. B. A. Alegria, L. M. D. R. S. Martins, A. J. L. Pombeiro, Adv. Synth. Catal. 350 (2008) 706–716.

Force fields for lipid bilayers: How do they work?

Ricardo O. Dias, Diogo Vila-Viçosa, Paulo J. Costa, Miguel Machuqueiro

*Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal
(joseriocardodias@gmail.com)*

The lipid bilayer is the basic structure of biological membranes, which define the boundaries of all cells and regulate the movement of molecules between cells and their environment. Membranes also enable other functions such as inter-cellular communication and energy conservation; in eukaryotic cells, membranes form discrete compartments constituting organelles, the specialized subcellular structures optimized for specific metabolic processes. However, lipids have been found to have more than a simple structural role [1,2].

In recent years, the field of molecular modeling of lipids has evolved significantly due to increases in computational power and improved force fields. Nevertheless, there are still serious limitations in all available force fields used to simulate lipid bilayers, which hinders further developments.

In this work, we studied how the most common lipid force fields are able to retain phosphocholine membranes in their fluid phases (Figure 1). Because lipids tend to gelidify, force fields developed tricks to keep them apart. In the case of GROMOS, we identified that the partial charges in the ester groups are determinant. We are currently investigating if a similar mechanism occurs in AMBER Lipid14 and CHARMM36 force fields.




Figure 1. POPC lipid bilayer in the fluid phase.

Acknowledgements

We acknowledge financial support from FCT projects UID/MULTI/00612/2013 and PTDC/REQCOM/5904/2014.

- [1] Loewen, C.; Gaspar, M.; Jesch, S.; Delon, C.; Ktistakis, N.; Henry, S.; Levine, T. *Science*, 304 (2004), 1644–1647.
- [2] Young, B. P.; Shin, J. J. H.; Orij, R.; Chao, J. T.; Li, S. C.; Guan, X. L.; Khong, A.; Jan, E.; Wenk, M. R.; Prinz, W. A.; Smits, G. J.; Loewen, C. J. R. *Science*, 329 (2010), 1085–1088.

One-step chemical immobilization of Laccase and Fe_3O_4 nanoparticles on gold for biosensing applications

I. Almeida*, F. Henriques, M. D. Carvalho, A. S. Viana

^aCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Ed. C8, Campo Grande, 1749-016 Lisboa, Portugal

**ibalmeida@fc.ul.pt*

Immobilization of enzymes on an appropriate support is a crucial step for the construction of biosensors and many efforts have been made in this field, especially combining enzymes with conductive materials [1]. The attachment of biomolecules on semiconducting metal oxide surfaces, such as magnetic iron oxide nanoparticles (Fe_3O_4 NPs), has been increasingly investigated, since these nanomaterials ensure the preservation of enzyme biological activity, with an amplification of the electrochemical signal [2].

In this work, a simple one-step methodology was explored for the preparation of enzyme-modified electrodes to be employed in biosensing interfaces (Figure 1). Small amine-containing molecules (epinephrine and hexylmethylamine) were used as proof-of-concept of the proposed methodology, whereas the enzyme Laccase is employed aiming the preparation of a sensitive biosensor towards phenolic compounds. This approach relies on the in situ dithiocarbamate formation between carbon disulfide and the amine groups of biomolecules [3,4], and also on the strong affinity between sulfur moieties and metals. The reactivity between carbon disulfide, Fe_3O_4 NPs with two distinct average sizes (ca. 20 and 40 nm) and amine groups, was firstly investigated using the hormone epinephrine, a small electroactive compound. A high amount of immobilized epinephrine and a facilitated redox conversion was obtained for modified electrodes with the larger NPs. Chronoamperometric studies of Laccase functionalized gold electrodes revealed a significant improvement of catalytic activity toward 2,2'-azino-bis(3-ethylbenzothiazoline-6-sulphonic acid (ABTS), when Fe_3O_4 NPs were added into the reaction. This effect was particularly noticed in the presence of the larger magnetite nanoparticles, which showed a very good sensitivity for ABTS oxidation (100 $\text{mA M}^{-1} \text{cm}^{-2}$). UV-visible spectroscopy confirmed the functionalization of nanoparticles by dithiocarbamate groups and atomic force microscopy was used to characterize the morphology of the nanostructured electrodes.




Figure 1 - Schematic representation of the one-step method exploited in this work and an AFM image of modified electrode with Laccase and Fe_3O_4 NPs.

Acknowledgements

This work was funded by FCT (Fundação para a Ciência e a Tecnologia): PhD scholarship (SFRH/BD/70673/2010) and projects: PEst 2015-2020-UID/Multi/00612/2013, IF/00808/2013 (POPH, UE-FSE) and PTDC/CTM-NAN/0994/2014.

- [1] M. M. Rodríguez-Delgado, G. S. Alemán-Nava, J. M. Rodríguez-Delgado, G. Dieck-Assad, S. O. Martínez-Chapa, D. Barceló, R. Parra, Trends in Analytical Chemistry 74 (2015) 21–45.
- [2] X. Shi, W. Gu, B. Li, N. Chen, K. Zhao, Y. Xian, Microchim. Acta 181 (2014) 1-22.
- [3] I. Almeida, A. C. Cascalheira, A. S. Viana, Electrochim. Acta 55 (2010) 8686-8695.
- [4] I. Almeida, V. C. Ferreira, M.F. Montemor, L.M. Abrantes, A. S. Viana, Electrochim. Acta 83 (2012) 311-320.

Anticancer and antioxidant activity from cork waste water

Luís Gomes^{a,b}, Maria Luísa Serralheiro^{b,c}, Miguel Minhalma^a, Rita Pacheco^{a,b}

^a Departamento de Engenharia Química, Instituto Superior de Engenharia de Lisboa, Rua Conselheiro Emídio Navarro, 1, 1959-007 Lisboa, Portugal

^b Centro de Química e Bioquímica, Faculdade de Ciências da Universidade de Lisboa, 1749-016 Lisboa, Portugal

^c Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa. E-mail: a38384@alunos.isel.pt;

This study aims to purify high value compounds from cork effluent fractions obtained using ultrafiltration membranes of cellulose acetate in order to accomplish the identification of the compounds recovered in the fractions and the evaluation of its potential as anticancer and antioxidant.

Cork is bark from the cork oak, is a plant tissue that covers the trunk of the tree and which can be removed every 9 years. In order to be used in industry, the planks have to be cooked in water, after which various cooking cycles must be replaced, leading to the so-called cork wastewater. ¹ This wastewater has a high organic loading rate and a high concentration of tannins and other polyphenols, and as such should be treated before proceeding to discharge. One of the strategies for the treatment of such effluent is the use of membranes, including most importantly the processes of reverse osmosis, microfiltration and ultrafiltration. ²

The pharmacological potential of cork lies in its low molecular weight components, such as the tannins, sterols, flavonoids and simple phenolic compounds. These show potential as anticancer, antimutagenic, antiallergic and anti-aging, and furthermore antioxidants derived from natural products can be less expensive than the synthetic variants. ¹

The effluent was fractionated using two ultrafiltration membranes prepared in the laboratory by reverse-phase method. Fractions were assayed in terms of phenols³, tannins⁴ and sugars⁵. Studies were undertaken such as antioxidant activity⁶, cytotoxicity studies and inhibition of cell proliferation in cell lines MCF-7. Additionally the compounds present in each fractions were also identified by LC-MS.

Some of the fractions collected demonstrated antioxidant activity values and inhibition of cell proliferation promising results. Regarding the cytotoxicity it can be said that none of the fractions has cytotoxic activity. Therefore, fractionation of the cork effluent by membrane process is an alternative to conventional treatment of such effluent, and permits recovery of the fractions, as well as the recovery of important bioactive compounds.



Figure 1 – Cooking stage of cork boards

Acknowledgements

FCT for the programmatic financing reference number: UID/MULTI/00612/2013.

- [1] Silva S. P., Sabino M. A., Fernandes E. M., et al., Int. Mat. Rev., 50 (2005) 345–365.
- [2] Bernardo M., Santos A., Cantinho P., Minhalma M., Water Res.45 (2011) 904–912.
- [3] M. Oktay, I. Gülçin, Ö.İrfan Küfrevioğlu, Food Sci.Technol. 36 (2003) 263-271.
- [4] J. P Van Buren, W. B. Robinson, J. Agric. Food Chem.17 (1981) 772–777.
- [5] X. Yong-Quan, H. Xiong-Fei, T. Ping, J. Yong-Wen, et al., Food Chem. 172 (2015) 831-835.
- [6] P.L. Falé, C. Ferreira, M.H. Florêncio, M. L. Serralheiro, et al. J. Med. Plants Res.7 (2013)1370-1378.

Synthesis of novel nucleos(t)ide and glycosyl phosphate analogs or mimetics possessing therapeutic potential

Nuno M. Xavier,^{a*} Daniela Batista,^a Alexandre Porcheron,^a Vítor J. Martins,^a Andreia Fortuna,^a Paulo J. Costa,^a Stefan Schwarz,^b Anne Loesche,^b René Csuk,^b Radek Jorda^c

Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Ed. C8, 5º Piso, Campo Grande, 1749-016 Lisboa, Portugal

^bBereich Organische Chemie, Martin-Luther-Universität Halle-Wittenberg, Kurt-Mothes-Str. 2, D-06120 Halle (Saale), Germany

^cLaboratory of Growth Regulators, Palacký University & Institute of Experimental Botany AS CR, Šlechtitelů 27, 78371

Olomouc, Czech Republic

*nmxavier@fc.ul.pt

Nucleoside and nucleotide analogs have attracted considerable attention as synthetic targets due to their propensity to display antiviral and antitumor effects.^[1] This bioactivity profile arises from their ability to interfere in biological processes in which natural nucleos(t)ides play key roles, such as nucleic acid synthesis and cell division, which are deregulated in diseases such as cancer or viral infections. Other biological properties have been reported for these classes of compounds, such as antimicrobial efficacies^[2] and cholinesterase inhibitory abilities.^[3] The access to structurally new analogs or mimetics of nucleosides and of nucleotides and the exploitation of their biological potential remains of interest.

In this communication, the synthesis of glucopyranos-6'-yl purines and pyrimidines as new types of regioisomers of nucleosides and that of glucuronamide-derived *N*-glycosyl compounds, including nucleosides and derivatives containing potential bioisostere groups for a phosphate functionality, will be presented.

6'-Isonucleosides were synthesized *via* Mitsunobu coupling between partially protected methyl glucopyranosides and various nucleobase derivatives.^[4,5] The outcome of the reaction was shown to be dependent on the substitution pattern of the sugar moiety when using pyrimidines, leading to uracil/thymine-linked pseudodisaccharides or to products of mono-coupling.

Peracetylated glucuronamides were the glycosyl donors used for N-glycosylation of a nucleobase or a sulfonamide, or were converted into anomeric azides for further access to glycosyl phosphoramidates. Azide-alkyne “click” cycloadditions enabled the synthesis of glucuronamide-based hybrids containing both a benzyltriazole moiety and an anomeric sulfonamide or a phosphoramidate function, as potential nucleotide mimetics.

Biological assays revealed some compounds as good inhibitors of acetylcholinesterase (AChE) or exhibiting cytotoxic effects to tumor cells, with inhibition constants or IC₅₀ values, respectively, in the micromolar concentration range. Molecular docking studies allowed inspecting the binding modes of the best AChE inhibitors to the enzyme.

The synthetic methodologies and the findings of the bioactivity evaluation will be revealed.

Acknowledgements

FCT is acknowledged for funding through the FCT Investigator Program (IF/01488/2013 and IF/00069/2014) and for financial support through the projects IF/01488/2013/CP1159/CT0006 and UID/MULTI/00612/2013.

- [1] L. P. Jordheim, D. Durantel, F. Zoulm, C. Dumontet, *Nat. Rev. Drug. Discov.* 12 (2013) 447–464.
- [2] M. Winn, R. J. M. Goss, K.-i. Kimura, T. D. H. Bugg, *Nat. Prod. Rep.* 27 (2010) 279–304.
- [3] N. M. Xavier, S. Schwarz, P. D. Vaz, R. Csuk, A. P. Rauter, *Eur. J. Org. Chem.* 2014 (2014) 2770–2779.
- [4] N. M. Xavier, S. D. Lucas, R. Jorda, S. Schwarz, A. Loesche, R. Csuk, M. C. Oliveira, *Synlett* 26 (2015) 2663–2672.
- [5] D. Batista, S. Schwarz, A. Loesche, R. Csuk, P. J. Costa, M. C. Oliveira, N. M. Xavier, *Pure Appl. Chem.* (2016) DOI: 10.1515/pac-2016-0102.

Sustainable carbon materials AS support for oxidation catalysts

Cécile Petit, Miguel V. Silva, Ana S. Mestre, Ana P. Carvalho, Carla D. Nunes*

Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

*cmnunes@fc.ul.p

Despite the greater number of studies focused on the immobilization of catalytically active metal complexes onto solid matrices this remains a hot research topic since the use of immobilized catalysts allows an ease separation of the reaction medium enabling successive reuses.

Inorganic materials, such as, silicas, clays or zeolites, and carbon materials, like carbon fibers, are among the most explored solids as catalyst support matrices. On the contrary, the use of activated carbons or other carbon materials prepared by less conventional methodologies is much more restricted. In this context the goal of this work is to test the potential of carbon materials obtained by acid hydrolysis of biomass followed by polycondensation [1], as catalyst supports.

A sisal derived carbon material was derivatized with a Mo complex – $[\text{MoI}_2(\text{CO})_3(\text{MeCN})_2]$ which binds to the surface of the carbon material with displacement of the labile acetonitrile ligands.

Materials were characterized by spectroscopic methods (FTIR, DRX), as well as SEM, TEM, thermogravimetry analysis, and N_2 adsorption at -196°C .

Selective olefin epoxidation is a very important process as epoxides are relevant building blocks across many areas. The synthesized materials have been tested in oxidation catalysis of enantioselective epoxidation of olefins (*cis*-cyclooctene, styrene and R-limonene) with tertiary butyl hydroperoxide at 55°C in dichloromethane during 24 h. Results (conversions, yields and enantiomeric excesses) were analyzed by gas chromatography-mass spectrometry (GC/MS). Kinetic profiles reproduced in Fig. 1 exemplify the results obtained, showing that the carbon material does not hold any intrinsic catalytic activity towards epoxidation of *cis*-cyclooctene but, conversely, Mo-derivatized material showed high catalytic activity towards the epoxidation of this olefin.




Figure 1. Kinetic profile for *cis*-cyclooctene epoxidation catalyzed by carbon material and Mo-derivatized catalyst.

Acknowledgements

The authors thank FCT for financial support (UID/MULTI/00612/2013). ASM thanks the financial support of FCT for the Post-doc grant SFRH/BPD/86693/2012.

[1] L. Wang, Y. Guo, B. Zou, C. Rong, X. Ma, Y. Qu, Y. Li, Z. Wang, *Bioresour. Technol.*, 2011, 102, 1947-1950.

Recovery of phenolic compounds from infusions: Antiacetylcholinesterase and antioxidant activity

João Carvalho^a, Inês Vasco^a, Andreia Maia^a, Tiago Santos^a, Rita Pacheco^{b,c}, Maria Luísa Serralheiro^{a,c}

^a Departamento de Química e Bioquímica, Faculdade de Ciências da Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

^b Área Departamental de Engenharia Química. Instituto Superior de Engenharia de Lisboa. Av. Conselheiro Emídio Navarro, 1959-007 Lisboa, Portugal

email : mlserralheiro@fc.ul.pt

^c Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal; email: mlserralheiro@fc.ul.pt

Infusions of leaves from plants or from peels of fruits is an easy and green way of obtaining compounds with biological activities, requiring only minor purification steps if a mixture of phenolic molecules is the main objective. In this work infusions from leaves and fruit of *Sambucus nigra* (Sabugueiro), Fig1 a, peels from *Annona cherimola*, Fig 1b was used to obtain phenolic compounds. The identification is under evaluation through LC-MS/MS. The antiacetylcholinesterase activity was determined for the non-purified extracts in the case of *S.nigra* and also for the purified peel extracts using DPPH method. The capacity to avoid the lipid peroxidation is also evaluated using the TBARs method. This activity is very important not only to provide, *in vitro*, the antioxidant activity that may happen in biological systems, as well as to develop new non-toxic antioxidants for the food industry. The biological activity of syrups from *S.nigra* are also under study. These syrups are highly recommended for flu and other ailments¹.



Figure 1: Material used in this study; (a) *Sambucus nigra*; (b) *Annona cherimola* fruit peels

This is the work of 4 volunteers from the Biochemistry course.

Acknowledgements

We acknowledge Fundação para a Ciência e Tecnologia (FCT) for financial support to Centro de Química e Bioquímica (Pest-OE/UI/UI0612/2013; UID/MULTI/00612/2013), and to Dr Sofia Maceira for suggesting the «sabugueiro» study and giving us the syrup to analyze.

[1] A.Proença-da-Cunha, A. Pereira-da-Silva, O. Rodrigues-Roque. Plantas e Produtos Vegetais em Fitoterapia, Fundação Calouste Gulbenkian, 2003, p. 568.

Modelling halogen bonds in molecular dynamics simulations of bacteriophage T4 lysozyme

Ricardo J. M. Rosa^{a,b,*}, Diogo Vila-Viçosa^b, Miguel Machuqueiro^b, Paulo J. Costa^b

^aDepartamento de Química, Faculdade de Ciências e Tecnologia, Universidade Nova de Lisboa, 2829-516 Caparica, Portugal

^bCentro de Química e Bioquímica, Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

* rjd.rosa@campus.fct.unl.pt

Halogen atoms are commonly used in drug design to fill hydrophobic cavities in protein binding sites, improve blood–brain barrier crossing or facilitate membrane permeability. Besides these non-specific effects, halogens are able to establish directional, non-covalent interactions, known as halogen bonds ($R-X\cdots B$, where $X = Cl, Br, I$ and $B =$ Lewis base). These bonds are explained by the existence of a positive region at the tip of the halogen atom, called σ -hole, which interacts with Lewis bases. Halogen bonds play an important role in protein-drug interactions and several structures deposited in the RCSB Protein Data Bank show this type of interaction, thus showing its potential for rational drug design. The implementation of halogen bonding in biomolecular force fields is not common and often relies on the use of massless points of charge to emulate the σ -hole. Despite its high popularity, the GROMOS force field does not contain such implementation.

Herein, we present a strategy to include halogen-bonding in the GROMOS 54A7 force field using the bacteriophage T4 lysozyme as a test case. Indeed, the X-ray structures of the bacteriophage T4 lysozyme [1] complexed with several C_6F_5X (where $X = H, F, Cl, Br$ or I) and C_6H_5X (where $X = H$ or I) molecules show evidence for $I\cdots S$ halogen-bonding. Molecular dynamics simulations of this protein bound to benzene or iodobenzene using the standard force field parameters will be compared with simulations performed with our halogen-bond capable implementation for iodobenzene.




Figure 1. Iodobenzene bound in the hydrophobic cavity of T4 lysozyme (3DN4). Protein is shown in cartoon and iodobenzene is shown as sticks.

Acknowledgements

The authors thank FCT for financial support (UID/MULTI/00612/2013). PJC acknowledges FCT for the “Investigador FCT” Program (IF/00069/2014).

[1] L. Liu, W. A. Baase, B. W. Matthews, *J. Mol. Biol.* 385 (2009) 595.

Spontaneous incorporation of silver particles in thiophene-based conducting polymers

Ricardo M. Assunção^{a,b}, Jorge P. Correia^{a*}

^aCentro de Química e Bioquímica, DQB, Faculdade de Ciências, Universidade de Lisboa, Portugal

^bDias de Sousa SA, Alcochete, Portugal

* jorge.correia@fc.ul.pt

Silver dispersions on polymer matrices are among the most studied hybrid systems involving electronically conducting polymers. The metal clusters formation may be carried out by electrodeposition [1], by electroless deposition[2] or, in the special case of polyaniline or polypyrrole, by electroless precipitation [1]. It is well known the capacity that those N-containing polymers have to spontaneously reduce noble metal ions in solution during its dynamic maintenance of the equilibrium [4,5]. Nevertheless, little is known about the mechanisms involved in metal spontaneous reduction by conducting polymers from other families. In this work a systematic study of the silver “electroless” reduction by conducting polymers from the family of thiophene is carried out.

Films of PTh (Polythiophene), P3MeTh (Poly(3-methylthiophene) and PEDOT_h(Poly(3,4-ethylenedioxythiophene)) were electrosynthesized on platinum substrates and immersed, in the reduced or oxidized state, in silver containing solutions under open circuit conditions. The electrode potential was monitored during this process as well as the electrode mass variations by microgravimetry. The modified electrodes were characterized by cyclic voltammetry (before and after exposure to silver solutions) and by hi-resolution microscopy (SEM and AFM). The results point to a galvanic displacement of silver in all the polymers immersed in the neutral state, concomitant with their spontaneous oxidation. However, after this initial process, the quartz crystal microbalance results show that electrodeless precipitation process occurs on the PEDOT_h matrix. It allows an accumulation of higher amounts of the metal at the polymer surface as reflected by the voltammetric results of this hybrid system. Local elemental analysis clearly confirmed the presence of the silver clusters in all the polymers, with varying sizes as monitored by microscopy.

Acknowledgements

This work is funded by FCT (Fundação para a Ciência e a Tecnologia) through the project UID/MULTI/00612/2013

- [1] SS Kim, SI Na, J Jo, DY Kim and IC Nah, *App. Phys. Lett.*, 93 (2008) 073307.
- [2] LM Abrantes and JP Correia, *Electrochim. Acta*, 45 (2000) 4179-85.
- [3] YP Ting, KH Neoh, ET Kang and KL Tan, *J. Chem. Tech. Biotechnol.* 59 (1994) 31-6.
- [4] A Mourato, AS Viana, JP Correia, H. Siegenthaler and LM Abrantes, *Electrochim. Acta*, 49 (2004) 2249-57.
- [5] V Tsakova, *Solid State Electrochem*, 12 (2008) 1421-34.

Mob1, hippo pathway member, is critical for *Toxoplasma gondii* replication

Alexandra Tavares^{1,4}, Inês Delgado¹, Samuel Francisco¹, João Coelho¹, Alexandre Leitão¹, **Helena Soares^{2,3*}**, Sofia Nolasco^{1,2}

¹ Centro de Investigação Interdisciplinar em Sanidade Animal, Faculdade de Medicina Veterinária, Universidade de Lisboa, Lisboa, Portugal

² Escola Superior de Tecnologia da Saúde de Lisboa, IPL, 1990-096 Lisboa, Portugal

³ Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

⁴ present address: Instituto Gulbenkian de Ciência, Oeiras, Portugal

*mhsoares@fc.ul.pt

Toxoplasma gondii is an obligate intracellular parasite of great veterinary and medical importance. It is able to evade the immune system of the host by converting from rapidly proliferating tachyzoites to latent bradyzoite cysts and this parasite number control is a key to the success of the infection. Pathways controlling cell division/proliferation like the Hippo pathway are likely candidates for regulating parasite replication. Human Mob1 participates in this pathway and our recent data suggests it is an excellent candidate in the control of parasite replication/number. Our research group has identified a single mob1 gene in *T. gondii*. A phylogenetic analysis of this gene showed it to be similar to other Apicomplexa but distant from protozoan parasites like the Trypanosomatida. We confirmed that this gene is expressed and our data show that its expression dramatically decreases (94%) during the parasite replication inside the host cell. We have constructed a transgenic parasite strain that overexpresses Mob1 and these parasites show a significant delay in the replication process. Using an in house polyclonal antibody against this protein we observed a very clear polarized localization of the protein in the parasite posterior pole, where the basal complex, a structure involved in cytokinesis in *T. gondii*, is localized. To better understand the role of *Toxoplasma* Mob1 we have created, by using the by CRISPR/Cas9 approach, a strain where Mob1 loss of function can be induced. Our preliminary results, by immunofluorescence microscopy, show that after induction *Toxoplasma* parasites in parasitophorous vacuoles (PV) lose their intrinsic polarity and their normal rosette organization. Indeed, inside of the PV it is difficult to identify the individual dividing parasites that seem to have originated a mass of abnormal cells where multiple nuclei are present. This result suggests that *Toxoplasma* cells have abnormal division and/or fail the cytokinesis. Altogether, the data support that Mob1 is involved in the control of *T. gondii* replication and is a promising candidate to target therapeutic agents against *Toxoplasma* parasites proliferation.

Acknowledgements

Support for this work was provided by FCT: EXPL/CVT-EPI/1945/2013, FRH/BD/101619/2014 (ID) and SFRH/BD/79423/2011 (SF).

Resource allocation to maintenance and stress response: set up of a quantitative experimental model

Miguel A. Rendas, Rafael Bento, Carlos Bernardes, Manuel E.M. da Piedade; Fernando Antunes

Departamento de Química e Bioquímica, Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Lisboa, Portugal

miguel_tgv00@hotmail.com (MR)

Saccharomyces cerevisiae (*S. cerevisiae*), used traditionally in the making of brewers and baking, is a eukaryotic unicellular organism able to grow rapidly in liquid media, making it the simplest model for the study of the eukaryotic cell. The aero-fermentative metabolism of this yeast presents some resemblance with the “Warburg effect” observed in some fast-proliferating cells, making the study of *S. cerevisiae* metabolism of vital importance.

A suitable tool for the study of yeast metabolism is *flow microcalorimetry* as it allows following in real-time the dissipated energy resulting mainly from the catabolism. We found that the dissipated heat associated to the cellular metabolic activity is dependent on the culture medium used. The maximum energy dissipated over time (power, P) obtained from the growth in 2% glucose in 3 different common media - YPD (Yeast Extract, peptone and dextrose), CAA (casein amino acids supplemented medium) and *Synthetic Complete* (SC) - was observed to be significantly higher in the rich medium YPD when compared to the less rich SC medium. Interestingly not only the maximum power changed but also the profile of the P over time; when grown in YPD or CAA, the power increases exponentially until reaching a maximum when occurs the switch from aerobic fermentation to respiratory metabolism and then returns to the baseline; when grown on SC medium the profile presents 2 peaks, the first possibly is caused by the depletion of one or multiple amino acids present in the synthetic medium resulting in a lower P and then a second peak similar to the other media is observed. The P per cell is maximal during the transition from lag to exponential phase but this maximum is higher in the (poor) SC than in a rich media like YPD. Interestingly, during the exponential phase, the P per cell is kept constant in YPD but not on SC, where it decreases continually. This analysis outlines the failure of SC medium, widely used in the selection of auxotrophic mutants in genetic approaches, to provide an unconstrained exponential growth. Unless strictly necessary, the use of SC media should be discouraged when studying the exponential phase as it is associated to a continuously shifting P per cell and, consequently, associated to a non-uniform growth phase.

To optimize exposure conditions to H₂O₂, the effects on the proliferative capacity of yeast cells were investigated. It was found that the killing kinetics of H₂O₂ follows a biphasic exponential decay: a first phase is observed immediately after the addition of 1 mM (lethal dose) H₂O₂ until 3 hours; a second phase follows where the population becomes more resistant to H₂O₂ and less cells die, when exposed to the same concentration of 1 mM, possibly due to adaptation to high concentrations of H₂O₂ or a bet-hedging mechanism. Similar results were observed when cell were previously adapted to H₂O₂, indicative that not all cells in the population become adapted in a typical adaptation protocol. Interestingly cells were able to resume their exponential growth even in the presence of such high concentrations of H₂O₂, implying an enhanced capacity to deal with the oxidative stress.

Finally, we studied the effect of non-lethal doses of H₂O₂ on the metabolism of growing *S. cerevisiae* via 3 *bolus* additions or a *steady state titration* over 90 min (same amount of H₂O₂ applied in both treatments). The P increased slightly immediately after the addition of H₂O₂ in both cases possibly as a result of increased catabolism over anabolism, necessary to the antioxidant response and adaptation. The exposure to the *steady state* is less harsher than with the 3 *bolus* additions since no difference in the heat produced was found compared to control in the first case but was higher in the second, meaning that the cells have to spend more resources to respond to the *bolus* exposure. The effect of several concentrations of H₂O₂ was also studied, observing an immediate decrease in P upon addition, associated with cell death, and a temporary arrest of the P-t curve, possibly associated with an adaptation response that was concentration-dependent, before resuming the typical growth.

Using *flow microcalorimetry* we demonstrate that *S. cerevisiae*'s metabolism is quite dependent on the culture medium and the nutrients contained in the media besides the carbon source, so that extra care must be taken when accessing for quantitative energy balances. Furthermore, the stress response is different depending on the type of administration (*steady state* or *bolus*) of the oxidant.

Acknowledgements

Supported by Fundação para a Ciência e a Tecnologia (FCT), Portugal, through project UID/MULTI/00612/2013) and a Post-Doctoral grant awarded to C. E. S. B. (SFRH/BPD/101505/2014).

Reorganization of plasma membrane lipid domains in *Neurospora crassa* during conidial germination

Filipa C. Santos^{a*}, Andreia S. Fernandes^{b, c}, Catarina A. C. Antunes^a, Filipe P. Moreira^a, Arnaldo Videira^{b, c, d}, H. Susana Marinho^a, Rodrigo F. M. de Almeida^a

^a Centro de Química e Bioquímica, Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

^b IBMC-Instituto de Biologia Molecular e Celular, Universidade do Porto, Rua do Campo Alegre 823, 4150-180 Porto, Portugal

^c Instituto de Investigação e Inovação em Saúde, Universidade do Porto, Rua Alfredo Allen 208, 4200-135 Porto, Portugal

^aICBAS-Instituto de Ciências Biomédicas de Abel Salazar, Universidade do Porto, Rua de Jorge Viterbo Ferreira 228, 4050-313 Porto, Portugal

*e-mail corresponding author: fpsantos@fc.ul.pt

Neurospora crassa, a haploid ascomycete, has been extensively used as a biological model for multicellular eukaryotes, namely in the study of developmental processes, apoptosis, aging and in the molecular basis of the circadian rhythm observed in asexual spore formation, conidiation. *N. crassa* cells have also been used for studies of resistance to antifungal drugs. *N. crassa* cells in the unicellular conidial stage have also ideal features to study sphingolipid (SL)-enriched domains. In this work, two fungal strains in the conidial stage (**Figure 1**), the wild-type (WT) and slime, which is devoid of cell wall, were studied, with different growth times, to evaluate how much their membrane biophysical properties are dynamic and relate with cellular stage. For a thorough biophysical characterization, detecting several types of domains and how they may differ in abundance and organization, the plasma membrane of *N. crassa* was labeled with three fluorescent membrane probes and several photophysical parameters measured, including fluorescence anisotropy and lifetimes, sensitive to distinct biophysical properties were measured. They were also compared with the unicellular fungus *S. cerevisiae*, one of the most widely studied eukaryotes, which has a plasma membrane with high ergosterol levels, whereas *N. crassa* conidia are practically devoid of ergosterol. Conidial growth of *N. crassa* was accompanied by membrane reorganization, namely increase of membrane fluidity, which occurs faster in the supplemented medium (SeM), needed for slime cells growth, than in minimal Vogel's medium (MMV). Gel-like domains, likely enriched in SLs, were found in both *N. crassa* strains. Those domains were particularly compact, rigid and abundant in the case of slime cells, even more than those occurring in *Saccharomyces cerevisiae*. Our results with *N. crassa* suggest that the melting temperature for SL-rich domains occurs near the growth temperature (30 °C) for WT, but at higher temperatures for slime, as previously observed for *S. cerevisiae*. Regarding biophysical properties strongly affected by ergosterol, the slime plasma membrane displays more similarities to *N. crassa* WT than to *S. cerevisiae* cells. The differences encountered during this study gave us new insights about the plasma membrane organization and structure of *N. crassa* strains, and led to a better understanding of the relationships between its lipid composition and dynamics during conidial growth.




Figure 1. Transmission microscopy images for *N. crassa* WT and slime cells at different phases of growth. WT cells in MMV at 5 h (A) and 12 h (D) of growth; WT cells in SeM at 5 h (B) and 12 h (E) of growth; slime cells at 5 h (C) and 12 h (F) growth), with a magnification of 100 $\times$.

Acknowledgements

Fundação para a Ciência e a Tecnologia (FCT), Portugal is acknowledged for PTDC/BBB-BQB/6071/2014, UID/Multi/00612/2013, and IF2012 initiative (POPH, Fundo Social Europeu). F.C.S. acknowledges PhD. Scholarship SFRH/BD/108031/2015.

Insecticidal activity of tropical and subtropical plant essential oils against *Aedes aegypti* larvae

Diara Rocha^{a,f*}, Maria Novo^{a,c}, Olívia Matos^{b,c}, Ana C. Figueiredo^d, Manuel Delgado^e, Cristina Moiteiro^f

^a Instituto de Higiene e Medicina Tropical, UNL, Rua da Junqueira, 100, 1349-008 Lisboa, Portugal. dkrocha@fc.ul.pt

^b Instituto Nacional de Investigação Agrária e Veterinária, INIAV. Quinta do Marquês, Av. República, 2784-505 Oeiras, Portugal

^c Unidade de Parasitologia e Microbiologia Médicas, IHMT, Rua da Junqueira, 100, 1349-008 Lisboa, Portugal

^d Universidade de Lisboa, Faculdade de Ciências, Departamento de Biologia Vegetal, CESAM, Centro de Biotecnologia Vegetal, Campo Grande, 1749-016 Lisboa, Portugal

^e Delegação do Ministério do Desenvolvimento Rural, MDR. Porto Novo, Santo Antão, Cabo Verde

^f Centro de Química e Bioquímica, Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal cmmoiteiro@fc.ul.pt

The use of plants for insect control has increased worldwide, with particular emphasis on search of essential oils (EOs) [1-3]. The emergence and re-emergence of vector-borne diseases, the increasing resistance to chemical insecticides, adverse effects on the environment, among others, it is become crucial to search for new insecticides compounds from natural or synthetic sources. In this perspective, we present the study of the larvicidal property of *Foeniculum vulgare* Mill. and *Mentha pulegium* L., from Cape Verde and Portugal, against *Aedes aegypti* Linnaeus 1762, the main vector of dengue (urban cycle).

EOs were obtained by hydrodistillation of *F. vulgare* and *M. pulegium* aerial parts. The chemical composition of all EOs from Portugal and Cape Verde were analysed by GC, GC-MS and ¹³C NMR. EOs were assayed against *Aedes aegypti* third instar larvae [4]. A dose-mortality effect was observed, allowing the determination of the LC₅₀, LC₉₀ and LC₉₉ by probit regression analysis [5].

Pulegone (61%) and menthone (20%) were the main EO constituents isolated from *M. pulegium* collected in Portugal, while menthol (30%), menthone and menthyl acetate (15% both), and pulegone (4%) dominated Cape Verde EO. Conversely, the EO compositions from *F. vulgare* collected at Cape Verde and Portugal were similar. *trans*-Anethole (32 and 30%), limonene (28 and 18%) and fenchone (10% both) were the main compounds identified in Cape Verde and Portugal EOs, respectively.

The EOs of these plants showed strong larvicidal activity 24h after exposure. This study reveal that the larvae of *Ae. aegypti* have increased susceptibility to the *F. vulgare* EOs from Cape Verde, which can be correlated with the highest percentage of the stereoisomers of limonene (LC₅₀ and LC₉₀ = 13.0 = 23.2 µl L⁻¹) [4].


This study allowed the chemical characterization and activity assessment of plants from two different geographic countries that were not studied before with highly promising results.

Acknowledgments

The authors thank Fundação Calouste Gulbenkian, PhD grant funding of Diara Rocha, Doctor Maria Cândida Liberato, Researcher Emeritus of the Institute of Tropical Scientific Research (IICT), Lisbon for the identification of the plant specimens and UPMM/IHMT funds and National Funds through the FCT under Pest-OE/EQB/LA0023/2011 and ID/MULTI/00612/2013.

References:

- [1] E. L. L. Baldin, A. E. M. Crotti, K. A. L. Wakabayashi, J. P. G. F Silva, G. P. Aguiar, E.S. Souza, R.C.S. Veneziani, M. Groppo, J. Pest Sci. 86 (2013), 301.
- [2] A. Ali, N. Tabanca, B. Demirci, E. K. Blythe, Z. Ali, K. H. C. Baser, I. A. Khan, J. Agr. Food Chem. 63 (2015), 447–456.
- [3] S. Tennyson, D. A. Samraj, D. Jeyasundar, K. Chalieu, Middle-East Journal of Scientific Research 13 (2013) 64–68.
- [4] Rocha D. K., Matos O., Novo M.T., Figueiredo A. C., Delgado M. & Moiteiro C. *Natural Product Communication*, 10 (2015) 677–682.
- [5] WHO (1998). WHO Geneva, Switzerland (28–30 September).



Personalised ICT supported service for independent living and active ageing: The contribution of functional ingredients and the website nutriageing

Alice Martins^{a*}, Marta Sousa Silva^a, António Ferreira^a, Luís Sousa^a, Catarina Dias^a, Ana Marta Matos^a, Vasco Cachatra^a, Daniela Batista^a, Antónia Turkman^b, Feridun Turkman^b, Marília Antunes^b, Helena S. Costa^c, Tânia Albuquerque^c, Ana Sanches^c, Amélia P. Rauter^a

^aCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

^bCentro de Estatística e Aplicações, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

^cInstituto Nacional de Saúde Doutor Ricardo Jorge, Departamento de Alimentação e Nutrição, Av. Padre Cruz, 1649-016 Lisboa, Portugal

*e-mail address (aimartins@fc.ul.pt)

Development of a new service model, to screen for and prevent frailty in older adults, covering nutrition, physical and cognitive domains, has been successfully accomplished with the participation of a multidisciplinary team from five countries, namely Portugal, Spain, Italy, Ireland, and The Netherlands and supported by the European Union under the FP7 program. This communication describes the work developed within the workpackage under the leadership of the Portuguese team, in particular regarding the creation of the innovative NUTRIAGEING website (nutriageing.fc.ul.pt), co-sponsored by IUPAC and by the Lisbon Municipality. It is a unique platform for the transfer of scientific knowledge into advice to the general public. It offers several modules to promote healthier nutrition and to educate the population on how to improve rational food habits. It is structured around healthy eating, cooking recipes with videos, and how to grow a vegetable garden, as a source of cheap and functional food ingredients. Announcing the latest findings reported on the literature, the scientific knowledge offered is complemented by research developed within the Carbohydrate Chemistry Group at CQB on the medicinal plants *Genista tenera*, *Salvia sclareoides* and *Erica australis*. Their non-toxic active principles with antidiabetic, anticholinesterase, anti-inflammatory and antioxidant properties will also be presented and discussed.

Acknowledgements

This work was supported by the European project “PERsonalised ICT supported Service for Independent Living and Active Ageing”, FP7-ICT-2013-10, GA 610359, 2013-2016 and by the IUPAC project “Healthy Life and active ageing – the contribute of functional food ingredients”, IUPAC 2013-054-01, 2013-2016. The authors are also grateful to the European Commission for approval of the INOVAFUNAGEING commitment and thank Fundação para a Ciência e a Tecnologia for financial support of the project UID/MULTI 0612/2013. The Lisbon Municipality is also acknowledged for the support and fruitful partnership.

Synthetic cobalt clays for the storage and slow release of therapeutic nitric oxide

Ana C. Fernandes^{a,*}, Moisés L. Pinto^b, Fernando Antunes^a, João Pires^a

^a Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^b CERENA, Instituto Superior Técnico, Universidade de Lisboa, Av. Rovisco Pais, n.º 1, 1049-001 Lisboa, Portugal


acfernandes@fc.ul.pt

Nitric oxide (NO) is one of the smallest endogenous molecules with particularly interesting roles in biological systems, despite its toxicological potential. [1] Solid carriers have potential biomedical interest in the delivery of exogenous NO for anti-bacterial, anti-thrombic and wound healing applications. [2, 3]

In this work, a smectite clay was successfully synthesized with incorporated cobalt ions in its structure, with the main goal of studying its potential in the field of storage and release of nitric oxide for therapeutic applications. Materials were characterized by X-ray diffraction (Philips PW 1730 diffractometer, nitrogen adsorption at -196 °C (Micrometrics, ASAP 2010), DLS (Zetasizer Nano ZS, Malvern Instruments) and chemical analysis of cobalt contents were obtained by Atomic Absorption Spectroscopy of digested samples using a Unicam 929 (ATI UNICAM) spectrophotometer.

The nitrogen adsorption-desorption isotherms (Fig. 1) show a high specific surface area, as recorded in Table 1, which is the result of a finely divided structure. In this table there are also the values of basal spacing (d_{001}).

The kinetic data for the nitric oxide storage and release was obtained in gas and liquid phases, as recorded in Table 1. The released amounts of NO in the liquid phase were within the biological range and slow release kinetics with a near direct relationship between the released fraction and time was observed. Toxicological assays with HeLa cells indicated that the materials have low cytotoxicity.



FFFig. 1 Nitrogen adsorption-desorption curves at -196 °C

Table 1 – Textural parameters.

	$d_{(001)}$ (Å)	A_{BET} (m ² g ⁻¹)	[NO] _G (mg _{NO} /g _{solid})	[NO] _L (µg _{NO} /g _{solid})
MMT	12.6	20	2.1	8
CoAS-A	14.2	218	8.0	16
CoAS-B	11.9	41	8.5	47
CoOS	13.0	246	6.9	33

Acknowledgements

The authors thank the Foundation for Science and Technology for funding CQB UID/MULTI/00612/2013 and CERENA (UID/ECI/04028/2013) and for the grant SFRH/BD/72058/2010 (ACF).

[1] R.E. Morris, P.S. Wheatly, *Angew. Chem. Int. Ed.*, 47 (2008) 4966-4981.

[2] M.L.Pinto, J. Rocha, J.R. Gomes, J. Pires, *J. Am. Chem. Soc.*, 133 (2011) 6396-6402.

[3] A.C. Fernandes, M.L. Pinto, F. Antunes, J. Pires, *J. Mater. Chem. B*, 1 (2013) 3287-3294.

Capture and activation of CO₂ by metal-organic structures

Janaína C. Almeida^{a*}, Sara Realista^a, Sónia Barroso^b, Maria José Calhorda^a, Paulo N. Martinho^a

^a Centro de Química e Bioquímica, DQB, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^b Centro de Química Estrutural, Instituto Superior Técnico, Universidade de Lisboa, Avenida Rovisco Pais, 1049-001 Lisboa, Portugal

* janacassia_20@hotmail.com

Currently hydrocarbon fuels are the most important source of energy because of their ready availability, stability, and high energy density. The increasing atmospheric concentration of CO₂ has been predicted by models which suggest dramatic and irreversible changes if actions are not taken urgently. It is now well established that CO₂ may be sequestered and stored as liquid CO₂ or solid carbonate but both solutions are energetically demanding and financially costly [1]. Recently reports of reaction of pressurised CO₂ with epoxides to form polycarbonates [2] or cyclic carbonates [3] have attracted both academic and industrial interest, but generally, the recycling of CO₂ to produce high value products via low-cost catalysts has not been exploited. Nelson's cryptands [4] are an example of a dynamic structure with useful applications, which demonstrated the ability to capture and convert CO₂ to carbonate and methyl carbonate following its coordination to encapsulated metal ions. Here we explore the fixation chemistry of CO₂ by derivatised dinuclear Cu(II), Ni(II), Zn(II) and Co(II) cryptands (Figure 1) where the phenyl ring was modified towards engineering these metal-organic structures into supramolecular assemblies. Attaching electron withdrawing groups to the phenyl ring proved to not affect their ability to capture CO₂.




Figure 1 Derivatised cryptands.

Acknowledgements

Support for this work was provided by Fundação para a Ciência e Tecnologia UID/MULTI/00612/2013. PNM and SR thanks FCT for financial support (SFRH/BPD/73345/2010 and PD/BD/52368/2013).

- [1] O.D. Bert Metz, Heleen de Coninck, Manuela Loos and Leo Meyer in: Carbon Dioxide Capture and Storage, UK, 2005, 431.
- [2] G.W. Coates, D.R. Moore, Angew. Chem. Int. Ed., 2004, 43, 6618.
- [3] J. Meléndez, M. North, R. Pasquale, Eur. J. Inorg. Chem., 2007, 3323.
- [4] Y. Dussart, C. Harding, P. Dalgaard, C. McKenzie, R. Kadirvelraj, V. McKee, J. Nelson, J. Chem. Soc., Dalton Trans., 2002, 1704.

Phenolic quantification in different *Camellia sinensis* infusions and relation with their antioxidant activity

Rute Martins ^{a,b}, Daniela Jones Jones ^{a,b}, Ana Marques ^a, Maria Helena Florêncio ^{a,b}, Rita Pacheco ^{a,c}, **Maria Luisa Serralheiro** ^{a,b,*}

^aCentro de Química e Bioquímica e Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa.

^bDepartamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa.

^cÁrea Departamental de Engenharia Química, ISEL, Av. Conselheiro Emídio Navarro, 1959-007 Lisboa, Portugal

*mlserralheiro@fc.ul.pt

Camellia sinensis is a plant largely consumed all over the world, Fig. 1. This plant can be subject to different treatments giving different types of beverages¹. The leaves can be used soon after being collected (green tea) or after being fermented (black tea). A new variety is the use of young leaves to prepare a different infusion (white tea). These dissimilar procedures of preparing the leaves, originate different tastes in the final beverage, due, in part to different content and different type of phenolic compounds¹. These beverages can even be consumed with other liquids like lemon juice or milk, for instance. The objective of this work is the quantification of total phenolic in these different beverages and its correlation with the antioxidant activity and to explain why the black tea changes its colour with the addition of lemon juice. The identification of the phenolic compounds will be carried out by LC-MS/MS and is under evaluation. Phenolics in the infusions as well as its antioxidant activity is indicated in Table 1. This is a small research project from students of the 3rd year of the Chemistry degree.



Figure 1: *Camellia sinensis*

Table 1: Total phenols (equivalent μg of gallic acid/mg of infusion) and antioxidant activity measured as DPPH extinction percentage (μg of extract/mL)

	Total phenols (gallic acid equivalent)	Antioxidant activity (DPPH)
Black tea	257.2 $\pm$ 88.2	0.41 $\pm$ 0.07
Green tea	290.1 $\pm$ 73.1	0.34 $\pm$ 0.17
White tea	97.2 $\pm$ 25.8	-
Black tea Yorkshire	298.0 $\pm$ 80.0	18.4 $\pm$ 1.16
Black tea Yorkshire with lemon juice (100 mL)	666.0 $\pm$ 70.0	8.48 $\pm$ 1.31

Acknowledgements

FCT for the programmatic financing reference number: UID/MULTI/00612/2013

[1] Y.Hilal, U. Engelhardt, J. Verbr. Lebensm. 2 (2007) 414 – 421.

Development of lysosomal-mimicking vesicles to study the effect of sphingosine abnormal accumulation on membrane biophysical properties

Ana C. Carreira^{a,b*}, Rodrigo F. M. de Almeida^b, Liana C. Silva^a

^a*iMed.Ulisboa –Research Institute for Medicines, Faculdade de Farmácia, Universidade de Lisboa, 1649-003 Lisboa, Portugal)*

^b *Centro de Química e Bioquímica, DQB, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal*

**anacarreira@ff.ulisboa.pt*

Sphingosine (Sph) is one of the simplest lipids and one of the most prevalent backbones of sphingolipids in mammals. This lipid plays important bioactive roles in different cellular processes and has been implicated in Niemann Pick type C1 (NPC1), a complex lysosomal storage disease. To understand how the accumulation of this lipid in NPC1 impacts lysosomal membrane structure and biophysical properties, we developed lysosomal-mimicking vesicles displaying internal acidic pH and external neutral pH. Moreover, the lipid composition of the vesicles was modified in order to resemble physiological- or NPC1-like lysosomes. To this end, ternary 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (POPC)/Sphingomyelin (SM)/Cholesterol (Chol) mixtures with, respectively, low and high Chol/SM levels were prepared. The effect of Sph on the membrane permeability and biophysical properties was then evaluated by fluorescence spectroscopy, electrophoretic and dynamic light scattering. Our results showed that Sph has the ability to cause a shift in vesicle surface charge, increase the packing properties of the membrane and promote a rapid increase in membrane permeability. These effects are enhanced in NPC1-lysosomal-mimicking vesicles, i.e., containing higher levels of Chol and SM. These results suggest that lysosomal accumulation of these lipids, as observed under pathological conditions, might significantly affect lysosomal membrane structure and integrity, and therefore contribute to the impairment of cell function.

Acknowledgements

This work was supported by Fundação para a Ciência e Tecnologia (FCT), Portugal: PTDC/BBB-BQB/3710/2014, UID/Multi/00612/2013, SFRH/BD/88194/2012 to ACC and IF2012 and 2014 initiatives (P.O.P.H., F.S.E.) to RFMA and LCS, respectively.

Interaction of *D*-glucose end-capped polylactide ruthenium cyclopentadienyl complexes with GLUT: insights from molecular docking

Carla Antunes^{a*}, Andreia Valente^b, Paulo J. Costa^a

^aCentro de Química e Bioquímica, Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749 - 016 Lisboa, Portugal

^bCentro de Química Estrutural, Faculdade de Ciências da Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

* carlaantunes1994@gmail.com

D-glucose end-capped polylactide ruthenium cyclopentadienyl complexes (RuPMC) show promising cytotoxic activity against several human cancer cells (IC₅₀ values in the μ M range [1, 2] and the sugar moiety in these complexes opens the possibility for a GLUT-mediated mechanism of cellular uptake. Indeed, most mammalian cells import glucose by a process of facilitative diffusion mediated by members of the GLUT family of membrane transport proteins. [3] In particular, GLUT1 is the most common glucose transporter, being overexpressed in many human cancers [4] thus providing a suitable target for cancer therapy.

In the last few years, pharmaceutical companies have been using *Virtual Screening* techniques, in particular, molecular docking, [5] in order to identify and select compounds able to bind a specific target. Docking techniques were also employed in the understanding of GLUT-mediated cellular uptake of carbohydrate-appended $[(\eta^5\text{-C}_5\text{H}_5)\text{Ru}(\text{N-N})(\text{PPh}_3)][\text{PF}_6]$ complexes. [6]

Given the promising nature of these RuPMC complexes, we used molecular docking in order to try to understand their interaction with GLUT1. Model complexes were built varying the length of the polymer chain (see Figure 1, left) and were docked on the cavity of the bacterial xylose transporter Xyle (PDB: 4GBZ), a GLUT1 homolog which was solved in an outward-open conformation thus providing a reasonable initial model for the scenario a given molecule encounters when entering a cell. Our docking results show that Xyle is able to recognize the *D*-glucose moiety of the complex and that the size of the polylactide chain is important.




Figure 1. Left: *D*-glucose end-capped polylactide ruthenium cyclopentadienyl complex (RuPMC) model; right: crystal structure of a bacterial homologue of glucose transporters GLUT1 bound to *D*-glucose.

Acknowledgements

PJC and AV acknowledge FCT, POPH and FSE - European Social Fund for the for the “Investigador FCT” Program (IF/00069/2014 and IF/01302/2013, respectively). PJC thank FCT for the UID/MULTI/00612/2013 project. A.V. acknowledges FCT for project UID/QUI/00100/2013.

- [1] A. Valente, M.H. Garcia, F. Marques, Y. Miao, C. Rousseau, P. Zinck, *J. Inorg. Biochem.* 127 (2013) 79-81.
- [2] M.H. Garcia, A. Valente, T.S.F. Morais, A.I. Tomaz, Patent application PCT/IB2015/002312.
- [3] B. Thorens, M. Mueckle, *Am. J. Physiol. Endocrinol. Metab.* 298 (2010) E141–E145.
- [4] L. Szablewski, *Biochim. Biophys. Acta Rev. Cancer* 1835 (2013), 164.
- [5] Kitchen *et. al.* *Nat. Rev. Drug Discov.* 3 (2004) 935-949.
- [6] P.R. Florindo, D.M. Pereira, P.M. Borralho, P.J. Costa, M.F.M. Piedade, C.M.P. Rodrigues, A.C. Fernandes, *Dalton Trans.* 2016, Accepted (DOI: 10.1039/C6DT01571A)

Using solvatochromic probes for the study of intermolecular interactions in 1,4-dioxane/methanol/acetonitrile solvent mixtures

Ricardo J. Nunes^{a,*}, Nelson Nunes^{a,b}, Ruben Elvas-Leitão^{a,b}, Filomena Martins^a

^a*Centro de Química e Bioquímica, Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Ed. C8, Campo Grande 1749-016 Lisboa, Portugal*

^b*Área Departamental de Engenharia Química, Instituto Superior de Engenharia de Lisboa, Instituto Politécnico de Lisboa, R. Conselheiro Emídio Navarro, 1, 1959-007 Lisboa, Portugal*

*ricardojgrn@gmail.com

Mixtures of different compositions were prepared for the binary systems 1,4-dioxane/acetonitrile and 1,4-dioxane/methanol (9 mixtures each) and for the ternary system methanol/1,4-dioxane/acetonitrile (22 mixtures), in a total of 40 mixtures.

UV-Vis absorption spectra were obtained at 298.15 K for five solvatochromic probes, namely betaine (30), 4-nitrophenol, 4-nitroanisole, 4-nitroaniline and *N,N*-dimethyl-4-nitroaniline. Preferential solvation was studied using the Bosch and Rosès model for the binary mixtures [1], and an extension of the model for the ternary mixture [2], allowing the establishment of the preferential solvation order for all species present in solution, including solvent “complexes” [3].

The applied model enabled the identification of synergistic behaviors in 1,4-dioxane/methanol and 1,4-dioxane/acetonitrile mixtures and therefore to conclude about the existence of solvent “complexes” in solution. No synergism was detected in the ternary system, where the behavior reflected a combination of the contributions of the underlying binary systems.

Kamlet-Taft solvent parameters π^* , α and β , were also calculated for all mixtures and their variation with composition carefully scrutinized.

The addition of small amounts of 1,4-dioxane to the mixtures was seen to cause a considerable variation in π^* . Additionally, in dioxane rich mixtures an accentuated effect in α was observed which was especially visible in the dioxane/methanol mixture.


Acknowledgements

The authors are grateful to Fundação para a Ciência e a Tecnologia, Portugal, for financial support under projects PEst-OE/QUI/UI0612/2013 and UID/MULTI/00612/2013.

[1] E. Bosch, F. Rived, M. Rosés, J. Chem. Soc. Perkin Trans 2 (1996) 2177-2184.

[2] N. Nunes, C. Ventura, F. Martins R. Elvas-Leitão, J. Phys. Chem. B 113 (2009) 3071-3079.

[3] N. Nunes, R. Elvas-Leitão, F. Martins, Spectrochim Acta A 124 (2014) 470-479.



Glycoside antibiotics and mechanism of action

João P. Pais,^{a,b} Ana S. Viana,^a Amélia P. Rauter,^a Luís. Sobral,^b Ricardo Dias,^b Rogério Tenreiro^b

^aCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa Portugal

^bInstituto de Biosistemas & Ciências Integrativas, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal
jppais@fc.ul.pt

The present work aimed to investigate the mechanism of action (MoA) of dodecyl 2,6-dideoxy- α -L-arabino-hexopyranoside, one of the most active compounds within a family of alkyl glycosides, previously discovered by our research group, that exhibited a potent antimicrobial activity against *Bacillus* species, in particular *Bacillus cereus* and *Bacillus anthracis*. Firstly, its antibacterial properties were assessed in vegetative and spore cells and the characteristics of the bactericide activity shown were identified, including the total absence of resistance to this compound even after long exposure.

Compound impact on the bacterial vitality, viability and bacterial metabolism was studied, providing guidelines for the forthcoming studies. Genetic dissection was carried out by random transposition and knock-out of specific membrane related targets.

Given the results obtained, the effects of the compound in different cellular ultra-structures were studied and finally observed by atomic force microscopy.

This study confirms that the mechanism of action of this family of compounds does not rely on molecular interactions with target enzymes, suggesting that membrane interactions should play a role in the selective antimicrobial events observed.

Acknowledgements

The authors thank FCT for financial support (PEST UID/MULTI/00612/2013), as well as the European Commission for funding the project FP7-PEOPLE-2013-IAPP. FCT and Cipan are also gratefully acknowledged for the Ph.D. grant of J. P. Pais (SFRH/BDE/51957/2012).

Biocidal microcapsules: a new strategy for biofouling control

Elisabete R. Silva^{a, b*}, Olga Ferreira^b, Raquel B. Feio^b, João C. M. Bordado^b, Patrícia Rijo^c, Ana C. Marques^{b, d}

^aCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^bCentro de Recursos Naturais e Ambiente, Instituto Superior Técnico, Universidade de Lisboa, 1049-001 Lisboa, Portugal

^cResearch Center in Biosciences and Health Technologies – CBIOS, Universidade Lusófona de Humanidades e Tecnologias, 1749-024

^dDepartamento de Engenharia, Universidade Atlântica, Fábrica da Pólvora de Barcarena, 2730-036 Barcarena, Oeiras, Portugal

*ersilva@fc.ul.pt

The adhesion and growth of micro/macroorganisms on surfaces in contact with water, is one of the most serious problems in water-based systems (e.g. shipping, water purification units, etc.). This undesired bio-attach, known as biofouling can promote substrate deterioration, systems clogging and fluids contamination, resulting on costly maintenance and retrofitting consequences. For instance, on shipping industry the biofouling accumulation on ships hulls can lead to drag friction increases up to 40% and subsequent fuel consumption and Greenhouse gas emissions increases [1]. The most efficient and conventional methods to control biofouling are mainly based on a chemical strategy. They comprise the direct and/or controlled releasing of biocides into the contaminated surface. Most of them employ coatings incorporating biocides which act by a controllable releasing mechanism. However, the main drawback of those systems is the poor control on biocides releasing and the potential degradation of those agents in the coating matrix. This leads to a reduction on the antifouling action of the coating, and thus on its lifecycle. Alternatively, higher biocides content can be used to achieve the required lifetime, but the continued release of those toxic agents into the environment has proven to cause serious side effects on ecosystems [2]. Environmental friendly technologies are therefore sought.

In this work, biocidal polyurethane-polyurea microcapsules (MC's) able to provide a more controllable biocidal action, by promoting partial/total biocide immobilization on MC's shell have been developed. Water-in-oil (W/O) microemulsion method combined with interfacial polymerization has been used for the purpose. Two main innovative strategies for the MC's synthesis were followed: i. the microencapsulation of a biocide (Econea) in the MC's core, together with its chemical immobilization in the MC's shell; ii. and, the chemical immobilization of the biocide in the MC's shell, by using prior modified biocides [3]. Optical microscopy images obtained for the developed MC's show that the first strategy leads to better MC's in terms of their morphologic properties, i.e, well-defined shape and size uniformity. Degradation of their morphology was shown to occur for higher contents of the binding biocide in the MC's shell. FTIR analysis evidenced that the MC's shell composition depends on the MC's core composition (water + surfactant) and biocide type and content. Higher water contents were found to promote polyurea formation within the shell. MC's bioactivity has been assessed using *Staphylococcus aureus* (ATCC 25923) bacteria within different mediums (artificial seawater, DMSO and MilliQ water). All developed MC's, from both strategies, proved to possess bioactivity. The best results in terms of morphological MC's properties and bioactivity were found for MC's prepared from the first strategy and with biocide contents as high as 20 wt.%.

As a proof of concept, MC's with 20 wt.% of biocide obtained from both strategies were incorporated in a marine polyurethane paint to coat acrylic substrates. Similar bioassays performed on those prototypes, evidenced a similar bioactivity for both MC's types. Leaching tests and antifouling assessment at simulated (aquarium) and real conditions (seawater) are in progress to better understand the involved antifouling action, i.e., bioactivity due to biocide leaching or by contact, or both.




Figure 1. Biocidal microcapsules with 20 wt%. biocide.

Acknowledgements

The authors acknowledge the financial support from FCT through SFRH/BPD/88135/2012, SFRH/BPD/96697/2013 and UID/MULTI/00612/2013. The authors also thank HEMPEL A/S for the marine paint providing.

[1] B. Dahlbäck, H. Blanck, M. Nydén, Cost. Mar. Sci. 34 (2010) 212-215.

[2] T. R. Bott, Industrial Biofouling, first ed., Elsevier, United Kingdom, 2011.

[3] E.R. Silva, O. Ferreira, J. C. Bordado, PCT application, PT2015/000050 (108096), 2015.

Biomembrane organization: Biophysical characterization of ordered domains

Gerson M. Lobo^{1*}, Telmo O. Paiva, Ana S. Viana, Rodrigo F.M. De Almeida¹

Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

*email address: gersonlobo@sapo.pt

The present work aims the biophysical study of the coexistence of membrane ordered domains, namely gel (so)/liquid-ordered (lo) phases, in membrane model systems of POPC/PSM/Chol. This system contains the main components of mammalian plasma membrane, and the corresponding phase diagram is already known (**Figure 1**) [1]. Different approaches have been utilized, including time-resolved and steady-state fluorescence measurements, using di-12-ANEPPQ probe labelling liposome suspensions, and solution Atomic Force Microscopy (AFM), to visually detect lateral organization and topographic features in three-component lipid membranes supported by mica substrate [2].

Fluorescence-based techniques reveals peaks at 470 nm and 635 nm, for excitation and emission spectra, respectively, in both pure POPC (liquid-disordered (ld)) and pure DPPC (so) membranes. However, a blue shift occurs in emission spectra, along a ternary so/lo tie-line (**Figure 1**), as well as an increased fluorescence intensity, with profiles that are totally consistent with the so/lo phase boundaries predicted in the phase diagram of **Figure 1**. These results were accompanied by a decrease of steady-state fluorescence anisotropy of di-12-ANEPPQ, suggesting a lower lipid/water interfacial order in lo membranes. The time-resolved fluorescence data reveals three different fluorescence lifetime components, and the mean fluorescence lifetime values of di-12-ANEPPQ are longer for lo membranes.

The membrane/water partition coefficient (K_p) of di-12-ANEPPQ for ld, lo, and so domains was also estimated plotting the intensity values against lipid concentration [3]. The results suggest that di-12-ANEPPQ tends to partition preferentially into an ld phase, but has higher quantum yield in lo phase.

Finally, AFM images reveals the presence of two types of lipid domains for membranes of POPC/PSM/Chol (1:7:2 molar ratio) (**Figure 1**-circle 4). For this lipid composition, domain splitting and thinning were observed, as well as a thickness difference of ca. 1.7nm – 0.9 nm between the two observed domains. It is expected that the more ordered and thicker domains are largely sterol-free and composed mainly by the highest T_m lipid (PSM) [4], as suggested by previous studies *in vivo* using *Saccharomyces cerevisiae* [5].

Acknowledgements

Fundação para a Ciência e a Tecnologia (FCT), Portugal is acknowledged for PTDC/BBB-BQB/6071/2014, PEst 2015-2020 (UID/Multi/00612/2013) and IF2012/2013 initiatives (P.O.P.H., F.S.E.). T. M. Paiva acknowledges fellowship under grant n. IF/00808/2013/CP1159/CT0003.

1. de Almeida, R.F.M. and E. Joly,. Front Plant Sci, 2014. 5.
2. Goksu, E.I., et al., Biochim Biophys Acta, 2009. **1788**(1): p. 254-66.
3. Bastos, A.E., et al., Chem Phys Lipids, 2012. **165**(5): p. 577-88.
4. de Almeida, R.F.M., A. Fedorov, and M. Prieto, Biophys J, 2003. **85**(4): p. 2406-2416.
5. Aresta-Branco, F., et al., J Biol Chem, 2011. **286**(7): p. 5043-54.




Figure 1- 1-palmitoyl-2-oleoyl-*sn*-glycero-3-phosphocoline (POPC)/ *N*-palmitoyl-sphingomyelin (PSM)/ cholesterol (Chol) lipid phase diagram used in the study of coexistence of membrane ordered domains. So/lo tie-line is represented by the black line. The lipid mixtures used in the fluorescence measurements are represented by the green circles numbered from 1 (5 mol% chol) to 7 (40 mol% chol). Adapted from reference [1].

Study of the permeability of phenolic compounds present in cladodes of *Opuntia ficus indica* and their effect on the cholesterol permeability through Caco-2 cell lines

Asma Ressaissi^{a,b}, Pedro L. Falé^{a,c}, Nebil Attia^b, Rita Pacheco^{a,d}, Maria Luísa M. Serralheiro^{a,c}

^a Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^b UR11ES33 'Integrated Physiology', Faculty of Sciences of Bizerte, Carthage University, Tunisia .

^c Departamento de Química e Bioquímica, Faculdade de Ciências da Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

^d Área Departamental de Engenharia Química. Instituto Superior de Engenharia de Lisboa. Av. Conselheiro Emídio Navarro, 1959-007 Lisboa, Portugal

email : mlserralheiro@fc.ul.pt

Currently one of the strategies used to improve health is the use of medicinal plants as functional foods due to their have medicinal properties, nutritional value, and pharmacological activities such as antioxidant, antithrombotic, anti-inflammatory, antiartertherogenic, and cardioprotective effects ^{1,2}. Lots of these studies report a positive correlation between consumption of plant foods rich in phenolic compounds (i.e. flavonoids) and a reduction of the risk to suffer from degenerative diseases. One of these foods is *Opuntia ficus indica* cladodes (nopal), it is a member of *Cactaceae* family. Several studies have reported that nopal exerts beneficial effects on health. In fact, cladodes are used for treating arteriosclerosis, diabetes, gastritis and hypercholesterolemia ^{3,4}. The aim of this work was to study the permeability of the different aqueous extracts from cladodes of *Opuntia ficus indica* through caco-2 cell lines. The effect of these extracts on the permeability of cholesterol was also studied. The study of toxicity on Caco-2, HepG2 and MCF-7 cells of all the aqueous extracts showed no toxicity against the 1 *Opuntia ficus indica* 3 types of cell lines. The permeability of the different compounds present in cladodes' extracts was between 5,3 and 15,6%. For the main compound, piscidic acid (identified by LC-MS/MS), the permeability was between 9,7 and 11,3%. Mixed with cholesterol, the extracts showed a reduction on the cholesterol permeability. The permeability of cholesterol in presence of cladodes extracts was between 10 times and 20 times less than the permeability of cholesterol in absence of extracts.



Figure 1: *Opuntia ficus indica*




Figure 2: piscidic acid

Acknowledgements

We acknowledge Fundação para a Ciência e Tecnologia (FCT) for financial support to Centro de Química e Bioquímica (PEst-OE/QUI/UI0612/2013; UID/MULTI/00612/2013), Asma Ressaissi acknowledges the Grant from Ministry of Higher Education and Scientific Research, Tunisia.

- [1] J. A. Vinson, et al., J. Agric. Food Chem. 43: (1995) 2800–2802.
- [2] C. Manach et al. Curr. Opinion Lipidol., 16 (2005) 77–84.
- [3] E.M. Galati, et al. Pharm. Biol. 41 (2003) 175–179.
- [4] A. Trejo-González, et al. J. Ethnopharmacol. 55 (1996) 27–33.

Characterization and identification of essential oils by GC-MS

J. Aço, M. H. Florêncio, C. Borges, A.P. Marques*

Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal;

*email apamarques@fc.ul.pt

Essential oils are fragrant essences of plants, usually volatile oils obtained by steam distillation or other modified methods from different parts of aromatic plants for example buds, flowers, leaves, roots and wood. In general, these oils are complex mixtures of organic compounds constituted by terpene hydrocarbons, oxygenated terpenes and sesquiterpenes. Essential oils have innumerable applications such as perfumery, cosmetics, aromatherapy, medicine and food. The use of these oils has grown due to its properties, especially anticancer, antiviral, antibacterial and antioxidant [1].

Gas chromatography coupled to mass spectrometry (GC-MS) is an important analytical technique, since it combines the power of chromatography separation with the capacity of mass spectrometric identification. GC-MS is a conventional and useful method for analyzing complex mixtures as essentials oils.

In this work, we characterized and identified by GC-MS the components of three essential oils: *Cananga*, *Ylang Ylang* and *Cinnamon Zeylanicum*. The equipment used was a Trace GC Ultra coupled to an ITQ 900 mass spectrometer with an automatic injector Triplus Rsh from Thermo Scientific. The results were acquired and processed by Xcalibur (version 1.2 from Thermo Scientific). All of the most important peaks were identified by comparison with literature data [2-4] and databases like Wiley 6 and NIST.

Acknowledgements

This work was supported by projects UID/MULTI/00612/2013 and REDE/1501/REM/2005 from Fundação para a Ciência e Tecnologia. We also thank to Plena Natura for providing the essential oils.

References

- [1] G. Buchbauer, in: K.H.C. Bas, G. Buchbauer (Eds.), Handbook of essentials oils: science, technology and applications, CRC Press/Taylor & Francis Group, Boca Raton, 2010, pp 235-280.
- [2] M. Brokl, M.L. Fauconnier, C. Benini, G. Lognay, P. du Jardin, J.F. Focant, , Molecules 18 (2013), 1783-1797.
- [3] R. Wang, R. Wang, B. Yang, , Innovative Food Sci. Emerging Technol. 10 (2009), 289–292.
- [4] E.M. Gaydou, R. Randriamiharisoa, J.P. Bianchinil, J. Agric. Food Chem. 34 (1988), 481-487.

[FeCl₂(HCpz₃)] heterogenized at hierarchical FAU for eco-friendly alkane oxidations

Vanmira Van-Dúnem^a, Ana P. Carvalho^b, Armando J.L. Pombeiro^c, Luísa M.D.R.S. Martins^{a,c}
Angela Martins^{a, b*}


^aADEQ, ISEL, IPL, Rua Conselheiro Emídio Navarro, 1959-007, Lisboa Portugal

^b Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^cCQE, Complexo I, IST, UL, Av. Rovisco Pais, 1049-001 Lisboa, Portugal

*amartins@deq.isel.ipl.pt

The purpose of this work is i) the preparation of hierarchical materials based on FAU zeolite to work as supports for the immobilization of a C-scorpionate iron(II) complex, the [FeCl₂(HCpz₃)] (pz = pyrazolyl) [1], and ii) their use as heterogeneous catalysts for the oxidation of alkanes, namely of cyclohexane to cyclohexanol and cyclohexanone under environmentally friendly conditions (Scheme 1).



Hierarchical FAU supports were prepared according to the procedure from [2] using NaOH or TPAOH in the presence of hexadecyltrimethylammonium bromide under autogenous pressure. Table 1 shows the main properties of the supports.

Table 1. Crystallinity (C_{XRD}) from X-ray patterns and textural parameters calculated from N₂ adsorption isotherms: microporous (V_{micro}) and mesoporous (V_{meso}) volumes.

Sample	C _{XRD} (%)	V _{micro} ^a (cm ³ g ⁻¹)	V _{meso} ^b (cm ³ g ⁻¹)
FAU	100	0.29	0.08
FAU/NaOH	94	0.26	0.15
FAU/TPAOH	92	0.23	0.18

^aEstimated from α_s method; ^bV_{meso}=V_{total}-V_{micro}, where V_{total} is volume adsorbed at p/p⁰=0.95

The heterogeneous catalysts were prepared using two immobilization procedures: the wet impregnation and the incipient wetness impregnation.

The catalytic tests have shown that these heterogeneous systems exhibit a similar catalytic activity to that found for the C-scorpionate complex in homogeneous media [1] with the advantage of easy separation and re-use.

Acknowledgements

Support for this work was provided by FCT through PEst UID/MULTI/00612/2013, UID/QUI/00100/2013, PTDC/QEQ-ERQ/1648/2014 and PTDC/QEQ-QIN/3967/2014 projects.

[1] T.F.S. Silva, E.C.B.A. Alegria, L.M.D.R.S. Martins, A.J.L. Pombeiro, *Adv., Synth. Cat.*, 350 (2008) 706-716.

[2] J. Garcia-Martinez, M. Johnson, J. Valla, K. Li, J.Y. Ying, *Catal. Sci. Technol.*, 2 (2012) 987-994.

Functional protein-based nanoparticles for reduction of cardiovascular diseases risk

André F. Silva^{a,*}, Joaquim T. Marquês^a, Ana S. Viana^a, Rita Pacheco^{a,b,*}, Maria Luísa M. Serralheiro^{a,c}

^aCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^bArea Departamental de Engenharia Química, Instituto Superior de Engenharia de Lisboa, Av. Conselheiro Emídio Navarro, 1959-007 Lisboa, Portugal

c Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande 1749-016 Lisboa.

*Corresponding author: andrefccsilva@gmail.com; rpacheco@deg.isel.ipl.pt.

Cardiovascular diseases are among the highest causes of death in EU ^[1], being the high cholesterol level in the blood one of the risk factors. The treatment often consists in the administration of drugs that are reported as presenting side effects. Previous work showed that plant infusions and decoctions containing flavonoids and other polyphenols can reduce the serum cholesterol levels ^[2, 3]. However, some of the compounds tested demonstrated low permeability (e.g. rutin) and also low stability and solubility in water ^[4]. Encapsulating these flavonoids in proteins can improve this, increasing the uptake by the cells which is expected to have an effect on cholesterol reduction.

As a vital macronutrient in food, proteins possess unique functional properties including the ability to form emulsions which allow them to be an ideal material for the encapsulation of bioactive compounds [5], therefore proteins are suited for the development of delivery systems for bioactive compounds in the form of nanoparticles (NPs). Furthermore, these are considered safe and have nutritional value, and hence the biological products obtained will be applied as functional foods.

The synthesis of the protein-based NPs was made using bovine serum albumin (BSA) loaded with rutin (Rut), as standard, and *Annona cherimola* leaves decoctions, Fig 1. Nanoparticles were prepared in a molar proportion of 10:1 of flavonoid (Rut) to BSA and mixed on a vortex ^[4]. For NPs loaded with *Annona cherimola* leaves decoctions, the extract stock concentration was adjusted in order for the prepared NPs to have the same Rut concentration as standards ^[3]. Obtained NPs were characterized through Atomic Force Microscopy (AFM), their permeability study was conducted using an epithelial cell line (Caco-2), normally used to stimulate the intestinal barrier, grown in Transwell plates. Cytotoxicity was determined with MTT viability test, using the same cell line.

In the characterization of each different nanopreparations, AFM has shown small nanoparticles of BSA with loaded Rut (from 20 nm up to 40 nm) and larger aggregates which corresponds to their association. The preparation with the greater dispersion of aggregates was found to be BSA/Annona, possibly due to the variety of the compounds present in *Annona cherimola decoctions*. Future work is expected to increase the uniformity of the nanopreparations. The results have demonstrated that the NPs are not toxic. Our work has accomplished a great improvement in the cells uptake of rutin when encapsulated in BSA in the intestinal barrier, this is expected to contribute in the increase of the benefits of decoctions in cholesterol reduction as well as in the supply of protein for the development of new food supplies




Figure 1: BSA-based nanoparticles containing rutin.
Image size: 1 μ m x1 μ m, z= 25 nm

Acknowledgements

FCT for the programmatic financing reference number: UID/MULTI/00612/2013.

References

- [1] ESC/EAS Guidelines for the management of dyslipidaemias, European Heart Journal, 32, 1769, 2011.
- [2] P. L. Falé, C. Ferreira, A. M. Rodrigues, F. N. Frazão, M. L. M. Serralheiro. Journal of Medicinal Plants Research, 8, 917, 2014.
- [3] P. L. Falé, C. Ferreira, F. Maruzzella, M. H. Florêncio, F.N. Frazão, M.L. M. Serralheiro. Ethnopharmacol, 25, 718-723, 2013.
- [4] R. Fang, R. Hao, X. Wu, Q. Li, X. Leng, H. Jing. J. Agric. Food Chem, 59, 6292-6298, 2011.
- [5] L. Chen, G. E. Remondetto, M. Subirade. Trends in Food Science and Technology, 17, 272-283, 2006.


CO₂ electroreduction by binuclear Ni(II) and Cu(II) complexes

Paulo N. Martinho,* Sara Realista, Paulo J. Costa, Maria José Calhorda


*Centro de Química e Bioquímica, DQB, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal
pnmartinho@ciencias.ulisboa.pt*

The increasing concentration of CO₂ in Earth's atmosphere and the resulting greenhouse effect have led to a wide effort toward the development of more efficient ways either to its capture, or more interestingly to its utilization in the syntheses of added value chemicals. Despite some reports dealing with the reaction of pressurised CO₂ with epoxides to form polycarbonates [1] or cyclic carbonates [2] and CO₂ conversion by Fe, Ni, Zn and Co based catalysts,[3] many challenges still remain.

Mononuclear (A,B) and binuclear (1, 2, 3, 4) salphen derivatives were synthesized and characterized, and shown to electroreduce CO₂. As depicted in the figure for complex 1, cyclic voltammetry in the presence of CO₂ leads to an increase in current and the appearance of a reduction peak.



The DFT calculated structure (PBE1PBE//6-31G**/LANL2DZ) of the Ni(II) binuclear complex displays a 40° angle between the planes of the two ligands. The HOMO is π^* M-salphen antibonding orbital. However, after oxidation the Singly Occupied MO is localized on two carbon atoms, explaining why these molecules easily polymerize upon oxidation.



The synthesis, electronic structure and reactivity of these complexes will be addressed and a model of the reaction reduction of CO₂ analyzed.

Acknowledgements

Support for this work was provided by Fundação para a Ciência e Tecnologia (FCT), Portugal, through projects UID/MULTI/00612/2013, contract IF/00069/2014 (PJC) and fellowships SFRH/BPD/73345/2010 (PNM), PD/BD/52368/2013 (SR). The authors acknowledge COST Action CM1205.

- [1] G. W. Coates, D. R. Moore, *Angew. Chem. Int. Ed.* 43 (2004) 6618.
- [2] J. Meléndez, M. North, R. Pasquale, *Eur. J. Inor. Chem.* 21 (2007) 3323.
- [3] J. Qiao, Y. Liu, F. Hong, J. Zhang, *Chem. Soc. Rev.*, 43 (2014) 631.
- [4] S. Realista, A. S. Viana, B. P. Cardoso, A. M. Botelho do Rego, P. D. Vaz, A. I. Melato, P. N. Martinho, M. J. Calhorda, *RSC Adv.* 5 (2015) 39495.

Energy-free reduction of silver from dilute solutions by electroless precipitation (re-)using polyaniline films

Inês J. Pereira^{a*}, Jorge P. Correia^a

^a*Centro de Química e Bioquímica, DQB, Faculdade de Ciências, Universidade de Lisboa, Portugal*

**ijordaopereira@gmail.com*

It is well known that the fully oxidized state of polyaniline (pernigraniline form) reduces to produce the protonated emeraldine in acid medium at open circuit potential. It is so expected that in the presence of metallic ions that can act as oxidizing species, spontaneous deprotonation occur with simultaneous reduction of the metal ion - eventually to zero oxidation state (figure 1) – continuing the process while the polymer is exposed to the solution [1]. The high electrochemical potentials of noble metals make them suitable for this electroless precipitation process in polyaniline (PAni) films [2]. This spontaneous, selective (to noble metals) and sustained reduction of metal ions is of particular importance in the field of extractive metallurgy [3].




Figure 1. Electroless precipitation mechanism

In this work the process of electroless precipitation of silver from acidic dilute solutions of silver ions is investigated. Thin PAnifilms were electrochemically synthesized on vitreous carbon and stainless steel electrodes and exposed to 1 mM silver solutions for different periods at ambient temperature. The amount of reduced metal in each experiment was assessed by atomic absorption spectroscopy. The effect of film thickness and immersion time in the silver extraction efficiency was evaluated by optical microscopy and electrochemical characterization of the pristine films and after exposure to the silver containing solutions. It was observed that the amount of reduced silver increases with polymer thickening and the metal presence doesn't affect the electroactivity of the polyaniline film. The selectivity of the electroless precipitation methodology to reduce noble metal ions was also evaluated by adding significant amounts of engineering metal ions to the silver solutions. The results clearly demonstrate the effectiveness of the methodology employed in this work to selectively reduce the noble metal ions in solution. The re-use of the polymer films for multiple extraction runs was explored as well.

Acknowledgements

This work is funded by FCT (Fundação para a Ciência e a Tecnologia) through the project UID/MULTI/00612/2013

[1] LM Abrantes and JP Correia, Mat. Sci. Forum, 1995, 191, 235-240

[2] V Lyutov and V Tsakova, J. Solid State Electrochem., 2011, 15, 2553-2561

[3] YP Ting, KH Neoh, ET Kang and KL Tan, J. Chem. Tech. Biotechnol., 1994, 59, 31-36

Luminescent hydrophobic surfaces for smart windows

J. B. Aldeias, S. Realista, M. J. Calhorda, P. N. Martinho

Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal
joanabal19@gmail.com

The general properties of a material and its wettability may have a huge impact on its usage and application in everyday life. Recently hydrophobic surfaces have attracted research teams to put efforts into combining wettability/hydrophobicity with other interesting properties.

For that purpose, we synthesized three new Zn(II) Schiff-base binuclear complexes displaying both hydrophobic and luminescent properties and studied their deposition on glass surfaces by the low-cost drop-casting method. The surfaces were prepared by dissolving the new Zn(II) Schiff-base complexes in three different solvents (chloroform, toluene and tetrahydrofuran) using different evaporation rates. Several parameters were taken into account: concentration of the Zn(II) complexes, solvent, temperature, number of layers and surface wettability. The techniques used to characterise both complexes and surfaces were ^1H and COSY NMR spectroscopy, FTIR spectroscopy, UV-vis spectroscopy, elemental analysis and contact angle microscopy. We have found that the best solvent to combine hydrophobicity, luminescence and transparency with our Zn(II) Schiff-base compounds was toluene.




Figure 1. Zn(II) Schiff-base hydrophobic luminescent surfaces.

Acknowledgements

We thank FCT for financial support UID/MULTI/00612/2013. PNM thanks FCT for financial support (SFRH/BPD/73345/2010).

References

- 1 Q. Wei, C. Schlaich, S. Prévost, A. Schulz, C. Böttcher, M. Gradzielski, Z. Qi, R. Haag and C. a. Schalley, *Adv. Mater.*, **26** (2014) 7358–7364.
- 2 S. Subhash Latthe, *J. Surf. Eng. Mater. Adv. Technol.*, **02** (2012) 76–94.
- 3 L. Y. Liu Y, Zhao X, Cai B, Pei T, Tong Y, Tang Q, *Pub.Med.*, (2014).

Synthesis and photocatalytic activity evaluation of new titanate nanotubes modified by co-doping with cobalt and rhenium

B. Barrocas*, O.C. Monteiro


Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal.

**btbarrocas@fc.ul.pt*

Nowadays, pharmaceuticals and personal care products (PPCPs), are a source of water contamination, and they create several environmental pollution problems. PPCPs include pharmaceutical drugs, cosmetics, food supplements and other personal care products. Unfortunately, after use, these products are disposed in the environment, usually without any treatment. Photocatalysis technology offers an efficient and environmental friendly approach for the elimination of different kinds of pollutants [1]. In particular, the use of nanocrystalline semiconductors on the photocatalytic treatment of wastewaters has been generating great interest.

Titanate nanotubes (TNTs) are very interesting since they combine the properties and applications of conventional TiO_2 nanoparticles, with the properties of layered titanates, such as photocatalytic activity and ion-exchange ability, respectively. In addition, the intrinsic properties of these 1D materials, mainly surface area, physical and chemical adsorption ability, optical and photocatalytic properties, can be tailored and adjusted to a specific application/interaction. However their high charge recombination rate and wide band gap limit their practical application. Therefore, the synthesis of TNT-based materials with either a broader range of light absorption or a lower charge recombination rate would be an important achievement toward the development of successful photoactive catalytic materials. For instance, the modification of TNTs with a transition metal (e.g. Co and/or Re), could reduce the charge recombination, and extend their optical absorption and photocatalytic performance of the pristine material.

In this work, nanocrystalline TNTs materials modified by cobalt and/or rhenium doping (CoTNT, ReTNT and CoReTNT) were successfully prepared, by hydrothermal method [2], and characterized using XRD, XPS, TEM and DRS. The influence of the Co and Re content on the optical and photocatalytic activity of the samples was investigated using the terephthalic acid as the probe molecule to study the catalytic production of hydroxyl radical ($\bullet\text{OH}$). The photocatalytic performance was subsequently studied using PPCPs (caffeine and paracetamol) as model pollutants. The best catalytic sample was then tested on a mixture of the two pollutants. The results show that the Co and/or Re modified TNTs samples (CoTNT, ReTNT and CoReTNT) are better catalysts than the unmodified TNTs.



TEM image of doped TNTs.

Acknowledgements

The authors thank FCT for financial support (UID/MULTI/00612/2013 and SFRH/BD/101220/2014).

- [1] D. Zhang, R.M. Gersberg, W.J. Ng, S.K. Tan, Environmental Pollution 184 (2014) 620-639.
- [2] E.K. Ylhäinen, M.R. Nunes, A.J. Silvestre, O.C. Monteiro, J. Mater. Sci. 47 (2012) 4305-4312.

Synthesis of new CNS-targeted drug-like leads against Alzheimer's disease: the role of the sugar moiety in C-glucosyl flavonoid analogues

Ana M. Matos,^a Joana Cristóvão,^b Cláudio Gomes^b, Amélia P. Rauter^a

^aCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Ed C8, Piso 5, Campo Grande, 1749-016 Lisboa, Portugal

^bInstituto de Biosistemas & Ciências Integrativas, Faculdade de Ciências, Universidade de Lisboa, Ed C8, Piso 5, Campo Grande, 1749-016 Lisboa, Portugal

Alzheimer's disease (AD) is the most common form of dementia, which currently affects over 46 million people worldwide.¹ Yet, despite continuous efforts over the past 30 years, no cure is possible with the drugs delivered to the market so far.

Natural polyphenols such as quercetin, apigenin and chrysin have been reported for their striking neuroprotective effects, including the ability to decrease amyloid- β (A β)-induced neurotoxicity while reducing β -secretase-mediated amyloid precursor protein (APP) processing, improving cognitive function and memory retention in rodents.²⁻⁴ Due to the multifactorial nature of AD, we envisage that the broad activity of polyphenols comes across as a desirable feature of new multitarget drug leads, that ought to tackle several pathological features, thus maximizing the chances of blocking neurodegenerative progression.

Preliminary assays conducted by our group have shown that flavones tend to be more effective in inhibiting the formation of A β_{1-42} amyloid fibrils than the analogue flavanones or isoflavones. Quercetin, in particular, showed an astonishing anti-amyloidogenic effect, which prompted us to synthesize flavone analogues with drug-like properties and to inspect the role of C-C linked glucose units, that stabilize amyloid polypeptides in their disaggregated state.⁵ Ultimately, it is our goal to unveil structural requirements for the anti-amyloidogenic and neuroprotective effects, and to find new blood brain barrier (BBB)-permeable molecules that combine key features of the flavone scaffold with the benefits of the sugar moiety.

In this communication, the synthesis of C-glycosylated and non-glycosylated flavone analogues will be presented and their therapeutic potential against AD discussed. Moreover, given the tight association between type 2 diabetes and AD, the usefulness of this type of compounds in type 2 diabetic patients with cognitive impairment will also be debated.

Acknowledgements

The authors thank Fundação para a Ciência e a Tecnologia for financial support (PEST UID/MULTI/00612/2013) and for funding the Ph.D. grant of A. M. Matos (SFRH/BD/93170/2013). The European Commission is also gratefully acknowledged for funding the project FP7-PEOPLE-2013-IAPP.

- [1] Prince M., Wimo A., Guerchet M., Ali G.C., Wu Y.T., Prina M. World Alzheimer's Report **2015**.
- [2] Sabogal-Guáqueta A.M., Muñoz-Manco J.I., Ramírez-Pineda J.R., Lamprea-Rodeiguez M., Osorio E., Cardona-Gómez G.P. *Neuropharmacology*, **2015**, 93, 134.
- [3] Zhao L., Wang J.L., Liu R., Li X.X., Li J.F., Zhang L. *Molecules*, **2013**, 18(8), 9949.
- [4] Li. R. Zang A., Zhang L., Zhang H., Zhao L., Qi Z. Wang H. *Neurol Sci*, **2014**, 35, 1527.
- [5] Ladiwala A.R.A., Mora-Pale, Lin C., Bale S.S., Fishman Z.S., Dordick J.S., Tessier P.M. *ChemBioChem*, **2011**, 12(11), 1749.

Opi1p translocation to the nucleus is regulated through oxidation by hydrogen peroxide in *Saccharomyces cerevisiae*

C. Camelo^a, F. Vilas-Boas^a, A. Cepeda^a, C. Real^a, F. Pinto^{c,b}, G. da Costa^a, H. Soares^{a,d}, H. S. Marinho^{a,b}, L. Cyrne^{a,b,*}

^a Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

^b Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^c BioISI- Biosystems & Integrative Sciences Institute, Campo Grande, Lisboa, Portugal

^d Escola Superior de Tecnologia da Saúde de Lisboa, 1990-096 Lisboa, Portugal

*mlcyrne@fc.ul.pt

Adaptation of yeast cells to hydrogen peroxide (H₂O₂) leads to a rapid change in membrane permeability accompanied by a decrease of membrane fluidity and the alteration of its lipid composition, allowing cells to survive to higher doses of H₂O₂ (1,2). During adaptation to H₂O₂ several genes that contain the regulatory element UAS_{INO}, and which codify for enzymes involved in phospholipid and fatty acid metabolism, are repressed (2,3). This repression is due to the translocation into the nucleus of the endoplasmic reticulum-bound transcriptional repressor Opi1p (4). However, the mechanisms of regulation of this translocation regulated by H₂O₂ are still unknown.

Oxidation of particular cysteine residues in proteins by H₂O₂ is involved in signaling cascades that culminate in the regulation of transcription (5). Opi1p has in its structure four cysteine residues that may be targets of oxidation. Such oxidation might be responsible for the H₂O₂-dependent translocation of Opi1p to the nucleus and subsequent transcriptional repression of target genes. Opi1p oxidation when cells were exposed to adaptive doses of H₂O₂ was confirmed by a protein electrophoresis (SDS-PAGE), after tagging oxidized protein cysteine sulphhydryl groups with methoxy-polyethylene glycolmaleimide (MAL-PEG) (6). To determine whether Opi1p cysteine residues are responsible for H₂O₂-mediated translocation of Opi1p to the nucleus, yeast strains with individual mutations in cysteine residues were prepared (cysteine to alanine substitutions). These mutations did not compromise the function of Opi1p as a transcriptional repressor, since all cells presented similar levels of expression of a reporter gene containing UAS_{INO}. As expected, the wild-type strain displayed Opi1p mainly in the endoplasmic reticulum, which translocated to the nucleus in the presence of H₂O₂. In clear contrast, Opi1p translocation to the nucleus in cells treated with H₂O₂ was impaired in the C159A mutant, with Opi1p being equally distributed in the periphery and inside the nucleus. These results identify cysteine residue 159 of Opi1p as responsible for H₂O₂-mediated translocation of this protein to the nucleus.

Acknowledgements

Supported by Fundação para a Ciência e a Tecnologia (FCT), Portugal (PEst-OE/QUI/UI0612/2013).

- (1) Branco, M. R., Marinho, H. S., Cyrne, L., and Antunes, F. (2004) *J. Biol. Chem.* **279**, 6501-6506;
- (2) Pedroso, N., Matias, A. C., Cyrne, L., Antunes, F., Borges, C., Malhó, R., de Almeida, R. F., Herrero, E., and Marinho, H. S. (2009) *Free Radic. Biol. Med.* **46**, 289-298;
- (3) Cepeda, A. (2010) Estudo da regulação da expressão do sintase de ácidos gordos de *Saccharomyces cerevisiae* na aquisição de resistência ao peróxido de hidrogénio. MSc. thesis, FCUL;
- (4) Loewen, C. J., Gaspar, M. L., Jesch, S. A., Delon, C., Ktistakis, N. T., Henry, S. A., and Levine, T. P. (2004) *Science* **304**, 1644-1647;
- (5) Marinho, H.S., Real, C., Cyrne, L., Soares, H., Antunes, F. (2014) *Redox Biol.* **2**, 535-562.
- (6) Wu H-H, Thomas J.A., and Momand J., (2000) *Biochem. J.* **351**, 87-93

Novel BiOCl-Bi₂S₃ nanostructures synthesized from deep eutectic solvents and its photocatalytic properties

Virgínia C. Ferreira*, Olinda C. Monteiro

Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

*vcferreira@fc.ul.pt

The development of new materials displaying improved chemical and physical properties have been subject of intense research for application in (photo)catalysis, supercapacitors and energy storage devices [1-4]. The possibility of preparing hybrid materials allows to combine the individual properties of its constituents in materials with improved performance which can be tuned, by design, through adequate combination [5,6].

In this work, a novel route for the preparation of bismuth oxychloride (BiOCl) nanoparticles sensitised with Bi₂S₃ in a one-step method, at room temperature from an environmentally friendly deep eutectic solvent, is reported. The influence of the synthesis conditions, temperature, sulphur source, concentration of reactants and presence of water, on the structural, morphological, optical and photocatalytic properties of the synthesized nanoparticles is analysed and discussed.

Under the experimental conditions used here, it was found that the sulphur source and the Bi:S ratio in the synthesis solution play important roles in the preparation and properties of such materials. The photocatalytic activity was tested towards the rhodamine B degradation under UV-vis light irradiation with the hybrid materials showing promising results.




Figure 1. BiOCl-Bi₂S₃ TEM image and its photocatalytic performance for RhB degradation under UV-vis light irradiation.

Acknowledgements

The authors gratefully acknowledge the financial support from Fundação para a Ciência e a Tecnologia (SFRH/BPD/77404/2011 and UID/MULTI/00612/2013).

- [1] G. Li, B. Jiang, S. Xiao, Z. Lian, D. Zhang, J. C. Yu, H. Li, Environ. Sci.: Processes Impacts 16 (2014) 1975-1980.
- [2] Z. Jiang, Y. Liu, T. Jing, B. Huang, Z. Wang, X. Zhang, X. Qin, Y. Dai, RSC Adv. 5 (2015) 47261-47264.
- [3] J. Yuan, J. Wang, Y. She, J. Hu, P. Tao, F. Lv, Z. Lu, Y. Gu, J. Power Sources 263 (2014) 37-45.
- [4] Y. Zhang, S. Lu, M.-Q. Wang, Y. Niu, S. Liu, Y. Li, X. Wu, S.-J. Bao, M. Xu, Mater. Lett. 178 (2016) 44-47.
- [5] C. Cao, L. Xiao, C. Chen, Q. Cao, Appl. Surf. Sci. 357 (2015) 1171-1179.
- [6] Y. Yang, F. Zhou, S. Zhan, Y. Liu, Y. Yin, J. Inorg. Organomet. Polym. 26 (2016) 91-99.

Development of a LC-MS/MS method for quantitative analysis of nine pharmaceutical compounds in surface water samples

Rodrigo A. Osawa^{a*}, Maria Helena Florêncio^a, Maria Rosário Bronze^{b,c}

^a Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^b Institute for Experimental Biology and Technology, Apartado 12, 2780-901 Oeiras, Portugal

^c Faculty of Pharmacy University of Lisbon, Av. Prof. Gama Pinto, 1649-019 Lisbon, Portugal

*rodrigo.osawa@gmail.com

Emerging contaminant, are by definition chemical compounds that are not monitored because they have unknown toxicological effects or have no regulatory legislation. Emerging compounds may be of synthetic or natural origin, found in products consumed by a large population and end up in natural ecosystems [1]. Pharmaceutical drugs are included in this group. Many of these drugs can cause aquatic toxicity, genotoxicity and endocrine interference on biota and humans. These compounds are found in superficial waters, underground waters and sewage effluents[2,3].

This study aimed to develop a method for the analysis, in an environmental matrix, of antidepressant drugs as amitriptyline (AMI), bupropion (BUP), venlafaxine (VEN), citalopram (CIT), trazadone (TRA) and duloxetine (DUL). Anticancer drugs as methotrexate (MET), ifosfamide (IF), cyclophosphamide (CYP) were also studied.

LC-ESI-MS/MS (electrospray ionization-tandem mass spectrometry) in MRM (Multiple Reaction Mode) was performed for the detection and quantification of those drugs. Sample preparation used a nylon membrane (0.45 µm) for filtration and added 0.1% formic acid to acidify the sample to pH 3 or 0.1M sodium hydroxide to pH 7 for comparison of the influence of pH on the extraction. Solid phase extraction was performed on a cartridge of hydrophilic-lipophilic balance, Oasis HLB (200mg, 6 mL, Waters) previously conditioned with 5 ml methanol and 5 ml water, washed with 5 ml 5% methanol in water and eluted with 5 ml methanol. The eluate was evaporated to dryness under a gentle stream of nitrogen and the dried extracts were stored at -5 °C until analysis.

The method showed good selectivity (Table 1), with detection limits between 0.6 and 1.2 ng L⁻¹ and quantification limits were between 2 and 4 ng L⁻¹. Superficial waters were collected and examined for the presence of possible contaminants. Recoveries were calculated by spiking the samples before extraction at a concentration of 40 ng L⁻¹ and for matrix effects, standards with blank water extracts spiked, were compared, after extraction. The results of the recoveries were better at pH 3 in the range of 72-110% and matrix effects for pH 3 were in the range of 59-103%. The method is adequate to detect and quantify these pharmaceutical drugs in environmental matrices. For future studies it is intended to collect samples of effluent treatment plants to determine the presence and confirm the removal of these compounds.

Table 1 - Conditions for the quantification of pharmaceuticals compounds.

Drugs	L.D (ng L ⁻¹)	L.Q (ng L ⁻¹)	Linearity (ng L ⁻¹)	r ²	Rt	m/z (quantification)	m/z (confirmation)	Re pH 3 (%)	Re pH 7 (%)	ME pH 3 (%)
CIT	1.2	4	4-400	0.9974	5.93	325>108.8	325>262	101.76	92.68	89.47
VEN	0.6	2	2-400	0.9954	5.44	278.1>57.7	278.1>260	105.02	51.38	62.55
AMI	1.2	4	4-400	0.9980	6.21	278.1>90.8	278.1>233	72.86	28.57	58.85
BUP	1.2	4	4-400	0.9962	5.32	240>184	240>166	118.58	108.0	81.49
TRA	0.6	2	2-400	0.9976	5.7	372>176.1	372>147.9	97.49	79.05	103.33
DUL	1.2	4	4-400	0.9952	6.21	298>153.9	298>43.7	109.86	59.58	96.01
MET	1.2	4	4-400	0.9930	5.56	454.9>308	454.9>175	96.21	107.8	78.32
CYP	0.6	2	2-400	0.9952	7.17	261>139.9	261>119.8	101.05	106.1	75.88
IF	1.2	4	4-400	0.9927	7.08	260.9>153.9	260.9>91.8	110.98	108.5	89.51

L.D = Limit of detection; L.Q = Limit of quantification; r² = Correlation coefficient; Rt = Retention time; R e= Recovery; ME = Matrix Effect.

Acknowledgements

The authors wish to thank Coordenação de Aperfeiçoamento de Pessoal de Nível Superior for PhD grants (BEX 0845-14-0) and CQB's PEST UID/MULTI/00612/2013.

[1] Stuart, M et al., The Science of the Total Environment 416 (2012), p.1–21.

[2] Giebulowicz, J et al., Ecotoxicology and Environmental Safety 104 (2014), p.103–109.

[3] Golovko, O et al., Chemosphere 111 (2014), p.418–426.

AChE inhibition and antioxidant activity of *Centaurium erythrae* (Fel da terra) infusions

Laura Guedes^{a,b}, Samuel Barata^{a,b}, Maria Helena Florêncio^{a,b}, Rita Pacheco^{a,c}, Miguel Machuqueiro^{a,b}, Maria Luísa Serralheiro^{a,b,*}

^a Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa.


^b Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa.

^c Área Departamental de Engenharia Química, Instituto Superior de Engenharia De Lisboa, Av. Conselheiro Emídio Navarro.

*mlserralheiro@fc.ul.pt

Centaurium erythrae is a plant (Fig.1 left) with laxative effect, used to treat lack of appetite and hyperglycemia. Acetylcholinesterase (AChE) is an enzyme localized in the neurosynaptic junctions¹ and its inhibition can increase intestinal motility². Aqueous extracts of leaves from *C. erythrae* showed an inhibitory activity towards AChE of 0.8 mg/mL (IC₅₀). The presence of gentiopicroside and secoiridoid glycoside was detected by LC-MS/MS (Fig. 1 right). The aqueous extract was fractionated by HPLC-DAD in order to separate the compounds present in the mixture: gentiopicroside (corresponding to the peak of the graph) and the compounds before and after this peak, corresponding to the first and third fraction, respectively. Docking studies revealed gentiopicroside as a mild AChE inhibitor. Indeed, the experimental results demonstrated that 300 µg/mL of gentiopicroside inhibited the enzyme in 23%, while the same amount of the third fraction, containing flavonoid derivatives, showed 40% of AChE inhibition, highlighting the relevant role of these compounds, and the first fraction inhibited in 29%. Further studies are ongoing to identify the remaining flavonoids present in these fractions. The aqueous extract showed an IC₅₀ of 1.7 mg/mL towards HepG2 and MCF7 cell lines. This modest value indicates that no toxicity was detected with the infusion. On the other hand, 0.5 mg/mL of gentiopicroside had a percentage of cytotoxicity of 15%, while the first and third fractions showed values of 11% and 37%, respectively, to the same concentration towards MCF7 cell line. The extracts, even after digestion with gastrointestinal enzymes, retained their AChE inhibitory activity and no modifications in the chromatogram were observed. The extracts when subject to the gastrointestinal enzymes did not lose the AChE inhibitory activity and when analysing the chromatogram no modifications were noticed. Once again third fraction had the best values regarding the antioxidant activity (EC₅₀ = 0.134 mg/mL), the first fraction had an EC₅₀ of 0.495 mg/mL and 0.05 mg/mL of gentiopicroside showed an antioxidant activity of 13%. In conclusion gentiopicroside has a moderate AChE inhibitory activity as well as a low antioxidant activity.

Figure 1. Picture of *C. erythrae* plant (left) and HPLC chromatogram of the extract (right) showing gentiopicroside identified as the main peak.



Acknowledgements

FCT for the programmatic financing reference number: UID/MULTI/00612/2013.

- [1] Devlin TM. Muscle contraction. In: Devlin, T.M. Eds. Textbook of Biochemistry with Clinical Correlations. Wiley-Liss, New York, 1997, pp. 954.
- [2] S. Cellek, R. Thangiah, E.M. Jarvie, S Vivekanandan, O. Laludew, G.J. Sanger, Neurogastroenterol Motil. 20 (2008) 539–545.

Novel titanate nanotubular materials modified with transition metals for the photocatalytic degradation of PPCPs

J. C. Cardoso*, O. C. Monteiro

Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal.

**joelcanhoto@gmail.com*

In the later years, the consumption of pharmaceuticals and personal care products (PPCPs) such as, antibiotics, stimulants, endocrine disruptors, food supplements and others, has increased. In result, the release of these compounds into the environment have been rapidly increasing and, with them of some phenomena such as, the appearance of antibiotic resistant bacteria or the increase of some endocrine diseases has also increased [1]. Several PPCPs removal methodologies have already been proposed but the definitive answer to solve this issue is far from being found.

The use of nanocrystalline semiconductors as photocatalysts, on the treatment of industrial wastewaters, has generated great interest, due to their unique physicochemical properties. TiO_2 has been the most extensively used semiconductor in the removal of a large number of organic pollutants from water systems. However, TiO_2 has a major drawback in processes associated with solar photocatalysis due to its wide bandgap (3.2 eV), making it difficult to implement an overall technological process based on TiO_2 [2]. The high recombination rate of photo-charge carriers is also a handicap for this semiconductor large scale application. Therefore, the synthesis of TiO_2 -based materials, *e.g.* titanate nanotubes (TNT), with a broader range of light absorption and a lower charge recombination rate would be an important achievement towards the development of successful photoactive materials [2].

This work reports on the synthesis of nanocrystalline TNT-based materials modified by both transition metal and ion-exchange (TNT/M) and doping (M-TNT), prepared via a hydrothermal treatment of an amorphous undoped and M-doped precursors recently reported. The transition metals selected for this study were Mn, Cu, Ni and Cr. The influence of the transition metal position in the TNT structure on the optical and photocatalytic properties of the materials was studied. The photocatalytic activity of the samples was firstly investigated using the terephthalic acid (TA) as probe molecule to study the catalytic production of hydroxyl radical ($\bullet\text{OH}$). Their photocatalytic performance was subsequently studied using caffeine and sulfamethazine as model pollutants. The results show that either M-TNT or TNT/M modified samples are better catalysts than the pristine TNT, being the photocatalytic performance dependent on the transition metal used and on its position in the TNT crystalline structure.

Acknowledgements

The authors thank FCT for financial support (UID/MULTI/00612/2013)

[1] B. Kasprzyk-Hordern, R. Dinsdale, A. Guwy, *Water Research* 43 (2009) 363-380.

[2] E. K. Ylhäinen, M. R. Nunes, A. J. Silvestre e O. C. Monteiro, *Journal of Materials Science* 47 (2012) 4305-4312

Exploiting 2-deoxy glycosides towards new antimicrobial and neuroprotective compounds

Catarina Dias, A. P. Rauter

Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Ed C8, Campo Grande, 1749-016 Lisboa, Portugal.

**catarinastdias@fc.ul.pt*

The increasing average life expectancy in developed countries has led to an escalating concern associated with the emergence of age-related diseases, such as dementia, a condition that is characterized by the loss of memory and other intellectual abilities interfering with daily life. Alzheimer's disease (AD) is the most common type of dementia noteworthy for its devastating nature and unsuccessful treatment options. The main characteristics of AD patients' brains include the extracellular deposition of amyloid- β peptides in senile plaques, the deposition of neurofibrillary tangles, the loss of synaptic function, inflammation and neuronal death.¹

Alkyl 2-deoxy/2,6-dideoxy-*arabino*-hexopyranosides with a potent antimicrobial activity in some Gram positive bacteria have been previously described by our research group,² and their mechanism of action was recently unravelled (unpublished results). Preliminary results have also shown that some of these glycosides interact with cystatin B amyloid fibrils, inspiring us to investigate 2-deoxy glycosides embodying natural neuroprotective polyphenols as aglycones. Rosmarinic acid was previously identified by our research group as the active principle of the neuroprotective plant *Salvia sclareoides*, as it prevents amyloid aggregation, and reduces a number of other events underlying AD pathology.³ Much attention has been paid also to resveratrol, due to its ability to inhibit A β oligomeric cytotoxicity and reduce neuronal cell death.⁴ Envisioning bioavailability enhancement of these products and taking into account the physico-chemical properties of carbohydrates/amyloid interaction, methodologies towards a series of rosmarinic acid and resveratrol glycosylated analogues were developed and will be presented. Their key structural features for neuroprotective properties will be proposed according to A β fibrillization assay results monitored by ThT fluorescence.

Acknowledgements

The authors thank FCT for financial support (PEST UID/MULTI/00612/2013), as well as the European Commission for funding the project *FP7-PEOPLE-2013-IAPP*. FCT and Cipan are also gratefully acknowledged for the Ph.D. grant of C. Dias (SFRH/BDE/51998/2012).

- [1] D.J. Selkoe. *Physiol. Rev.* 81(2) (2001) 741.
- [2] A. Martins, M.S. Santos, C. Dias, P. Serra, V. Cachatra, J.Pais, A.P. Rauter et al. *Eur. J. Org. Chem.* (2013) 1448.
- [3] a) F. Marcelo, C. Dias, A. Martins, P.J. Madeira, T. Jorge, M- H. Florencio, J. F. Canada, E. Cabrita, J. Jimenez-Barbero, A. P. Rauter. *Chem. Eur. J.* 19(21) (2013) 6641. b) C. Airoidi, E. Sironi, C. Dias, F. Marcelo, A. Martins, A.P. Rauter, F. Nicotra, J. Jimenez-Barbero. *Chem. Asian J.* 8 (2013) 596. c) T. Alkam, A. Nitta, H. Mizoguchi, A. Itoh, T. Nabeshima. *Behav. Brain Res.* 180(2) (2007) 139.
- [4] S.D. Rege, T. Geetha, G.D. Griffin, T.L. Broderick, J.R. Babu. *Front. Aging Neurosci.* 6 (2014) 1.

Are pH-sensitive amino acids in their most common protonation states at the water/membrane interface?

Pedro B. P. S. Reis, Vitor H. Teixeira, Diogo Vila-Viçosa, Miguel Machuqueiro

Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal (pdreis@ciencias.ulisboa.pt)

The study of biological membranes has been for quite some time a challenge for researchers. A detailed description of the water/membrane interface has to take in consideration all important factors and pH is recognizably one of them, even though it is usually ignored due to its high complexity in terms of modelling. The pK_a values of typical titrable amino acids can be significantly influenced by changes in the environment, i.e., peptide insertion into a lipid bilayer [1-2].

The main objective of this work is a comprehensive study of how a membrane environment can shift the pK_a values of common pH-sensitive amino acids (Asp, Glu, His, Lys, Cys, Tyr, and the N- and C-termini). For this, we used our recently developed CpHMD-L methodology [3] with a DMPC membrane and the model Ala-based pentapeptides that have already been well characterized in water by Pace and co-workers [4]. With this approach, we intend to capture the coupling between conformation/configuration/ insertion and protonation at the membrane interface, taking into consideration that if complete insertion occurs, the peptides will probably no longer be able to exchange protons with the solvent (Figure 1).




Figure 1. The Ala₅ pentapeptide at the water/membrane interface.

Acknowledgements

We acknowledge financial support from FCT projects UID/MULTI/00612/2013 and PTDC/QUECOM/5904/2014.

- [1] J. F. Hunt, P. Rath, K. J. Rothschild, D. M. Engelman, *Biochemistry* 36 (1997) 15177.
- [2] P. R. Magalhães, M. Machuqueiro, A. M. Baptista, *Biophys. J.* 108 (2015) 2282.
- [3] D. Vila-Viçosa, V. H. Teixeira, A. M. Baptista, M. Machuqueiro, *J. Chem. Theory Comput.* 11 (2015) 2367.
- [4] R. L. Thurlkill, G. R. Grimsley, J. M. Scholtz, C. N. Pace, *Protein Sci.* 15 (2006) 1214.

Mn(III) single ion magnet with a tridentate Schiff-base ligand

S. Realista^{a*}, A. J. Fitzpatrick^b, G. Santos^a, L. P. Ferreira^{c,d}, S. Barroso^e, L. C. J. Pereira^f, N. A. G. Bandeira^{a,g}, P. Neugebauer^h, G. G. Morgan^b, J. van Slageren^b, M. J. Calhorda^a, P. N. Martinho^a]

^aCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

^bSchool of Chemistry, University College Dublin, Belfield, Dublin 4, Ireland

^cBioISI, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

^dDepartment of Physics, University of Coimbra, 3004-516 Coimbra, Lisboa

^eCentro de Química Estrutural, Instituto Superior Técnico, Universidade de Lisboa, 1049-001 Lisboa, Portugal

^fC2TN, Centro de Ciências e Tecnologias Nucleares, Instituto Superior Técnico, Universidade de Lisboa, Estrada Nacional 10, ao Km 139.7, 2695-066 Bobadela LRS, Portugal

^gInstitute of Chemical Research of Catalonia (ICIQ), Avda. Paisos Catalans, 16- 43007 Tarragona, Spain

^hInstitut für Physikalische Chemie, Universität Stuttgart, Pfaffenwaldring 55, 70569 Stuttgart, Germany

*smrealista@ciencias.ulisboa.pt

Single ion magnets (SIMs) are a class of materials with potential application as high-density magnetic memories and quantum-computing devices in spintronic field.[1] The size of the barrier of the reversal magnetization (U_{eff}) is the determining factor to the suitability of a single ion magnet to be applied in data storage devices. Efforts aiming at maximizing the anisotropy by an appropriate ligand field have been made to achieve high barriers.[2] Manganese(III) is a d^4 metal ion displaying a Jahn-Teller (JT) distortion when in an octahedral coordination environment. This feature turns Mn(III) into an promising ion to study its magnetic properties, namely spin crossover[3] and single ion magnet[2]. We report the synthesis of Mn(III) Schiff base cationic complexes (Figure 1) using different counter anions. SQUID magnetometry showed that all compounds are in the high-spin state with one pair of bond lengths (Mn-Namine) considerably longer than the others (Figure 1). Both magnetic and HF-EPR measurements reveal that the complex has the largest axial zero field splitting ($D = -4.6 \text{ cm}^{-1}$) known to date for a Mn(III) single ion magnet. AC magnetic measurements at 2000 Oe allowed to determine the energy barrier for spin reversal (10.19 K) and spin reversal relaxation time ($1.476 \times 10^{-6} \text{ s}$) for the Mn(III) ion. Computational studies were used to characterise the electronic structure and substantiate the zero field splitting in the Mn(III) complex.




Figure 1. Cation structure

Acknowledgements

This work was supported by Fundação para a Ciência e a Tecnologia (FCT), Portugal (Projects UID/MULTI/00612/2013, UID/MULTI/04046/2013 and UID/Multi/04349/2013). FCT is gratefully acknowledged by S.R. for a PhD grant (PD/BD/52368/2013) and by P.N.M. for a postdoctoral grant (SFRH/BPD/73345/2010). N.A.G.B. gratefully acknowledges the COFUND 291787-ICIQ-IPMP grant. A Science Foundation Ireland Investigator Project Award 12/IP/1703 is gratefully acknowledged by G.G.M. and funding from the National University of Ireland, University College Dublin and the Cultural Service of the French Embassy in Ireland for scholarships is gratefully acknowledged by A.J.F. The COST action CM1305 is acknowledged by P.N.M. and G.G.M.

- [1] J. Vallejo, A. Pascual-Álvarez, J. Cano, I. Castro, M. Julve, F. Lloret, J. Krzystek, G. De Munno, D. Armentano, W. Wernsdorfer, R. Ruiz-García, E. Pardo, *Angew. Chemie Int. Ed.*, 2013, 52, 14075.
- [2] G. A. Craig, M. Murrie, *Chem. Soc. Rev.*, 2015, 44, 2135.
- [3] P. N. Martinho, B. Gildea, M. M. Harris, T. Lemma, A. D. Naik, H. Müller-Bunz, T. E. Keyes, Y. Garcia and G. G. Morgan, *Angew. Chemie - Int. Ed.*, 2012, 51, 12597.

Polymorphism or thermal disorder in 4-hydroxyheptanophenone?

Cátia S.D. Lopes^a, M. Fátima M. Piedade^{a,b}, Ricardo G. Simões^a, Herminio Diogo^{a,b}, Carlos E. S. Bernardes^a, Manuel E. Minas da Piedade^{a*}

^a Centro de Química e Bioquímica e Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^b Centro de Química Estrutural, Complexo Interdisciplinar, Instituto Superior Técnico, 1049-001, Lisboa, Portugal

* memp@ciencias.ulisboa.pt

Studies of polymorphism involving families of structurally related molecules are particularly interesting to understand how the interplay of molecular size, shape and interactions may affect the packing architectures and relative stability of different crystal forms. One such families consists of 4-hydroxybenzoyl compounds, $\text{HOC}_6\text{H}_4\text{C(O)R}$ ($\text{R} = \text{H}$, alkyl), differing only in the length of the alkyl chain bonded to the carbonyl group. Based on these compounds it is, for example, possible to investigate how the hydrogen bond pattern sustaining the packing is affected by changes in the alkyl chain length.

Results of single crystal X-ray diffraction (SCXRD) analysis on 4-hydroxyheptanophenone will be reported. These results evidenced that on increasing temperature in the range 150-298 K, the crystal system is unaltered and the observed structural changes are essentially related with modifications in the conformational freedom of the alkyl chain (Figure 1). These changes are likely to be responsible for heating rate sensitive thermal events detected by differential scanning calorimetry. The 4-hydroxyheptanophenone packings corresponding to the low and high ends of the temperature range probed for will be compared with those also obtained for other $\text{HOC}_6\text{H}_4\text{C(O)R}$ compounds that were also studied by SCXRD.




Figure 1. Overlay of molecular structures of 4-hydroxyheptanophenone at 150 K (■), 190K (■), 220K (■), and 290K (■)

Acknowledgements

This work was supported by Fundação para a Ciência e a Tecnologia, Portugal (Project UID/MULTI/00612/2013 and UID/QUI/00100/2013). A post-doctoral grant from FCT is gratefully acknowledged by Carlos Bernardes (SFRH/BPD/101505/2014).

Photocatalytic degradation of carbamazepine and ibuprofen in water using novel TiO₂-MoO₃ materials


J. Diniz, C.D. Nunes, O.C. Monteiro

*Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal
ojorgediniz@gmail.com*

The increase of the amount of pharmaceutical and cosmetic products in lakes, rivers and even drinking water has become a concerning matter for the world's health. Such chemicals are classified as emerging pollutants. [1] Different treatment methods have been reported but heterogeneous photocatalysis gained increased attention, with titanium oxide (TiO₂) being the most known and used photocatalyst in water treatment. Such choice is due to its strong photocatalytic activity, chemical stability, nontoxicity and low cost. However, it presents a critical drawback: the wide band gap of TiO₂ makes only possible the use of the ultraviolet fraction of the solar light (3–5%) [2]. Great efforts have been made for shifting its light absorption towards the visible region and thus improve its photocatalytic efficiency. Similarly, molybdenum oxides are also attractive due to their unique structural and optical properties and they are broadly employed in electrochromic and photochromic devices, gas sensors and lithium batteries.

In this study, new photocatalytic materials were prepared by combination of TiO₂ and MoO₃ particles. Nanocrystalline particles and nanocomposites were prepared using a hydrothermal approach. The new hybrid materials were structural, morphological, and optically characterized by X-ray diffraction (XRD), transmission and scanning electron microscopy (TEM/SEM), diffuse reflectance spectroscopy (DRS) and Diffuse Reflectance Infrared Fourier Transform Spectroscopy (DRIFTS).

Regarding their future photocatalytic application, the materials were first used to evaluate the catalytic quantum yield of hydroxyl radical (•OH) using terephthalic acid as probe molecule. The main purpose of the new hybrid materials was achieved by testing their photocatalytic performance concerning the degradation of carbamazepine and ibuprofen as water pollutants and preliminary results show their promising application in water treatments.



Acknowledgments

The authors thanks Fundação para a Ciência e Tecnologia for financial support (UID/MULTI/00612/2013).

[1] T. A. Ternes, *Water Res.*, 32(1998), 3245–3260.

[2] M. Kaneko and I. Okura (Eds.), *Photocatalysis: Science and Technology*, Biological and Medical Physics Series, Springer, Berlin 2010.


pH-dependent insertion of pHLIP peptide into lipid bilayers: pK_a values of key residues

Diogo Vila-Viçosa*, Vitor H. Teixeira, Miguel Machuqueiro

Centro de Química e Bioquímica, Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

* diogo.vicosa@fc.ul.pt

The pH (low) insertion peptide (pHLIP)¹⁻³ is a family of peptides that are able to insert into a lipid bilayer at acidic pH. These peptides are based in a transmembranar sequence of bacteriorhodopsin that is unstructured in solution (stage I), interacts with lipid bilayers remaining unstructured at neutral pH (stage II) and inserts into the bilayer with a significant α -helical content at acidic pH (stage III). This family of peptides have already been used to target tumor cells in vivo since acidosis is an hallmark of these tissues.⁴ These events are difficult to study at the molecular level, in particular, the relation between the pK of insertion of pHLIP peptides and the pK_a of some key residues is yet to be clarified. In this work, we used a linear response approximation to determine the pK_a of these residues. We studied four different pHLIP variants to understand the importance of the ASP positions and its mutation to GLU. For these variants, there are experimental data available that we used to validate our approach. Finally, we also propose a mutation to a HIS residue in the sequence, expecting to turn off the pHLIP peptide insertion into the membrane (stage III) at too low pH values.



We acknowledge the financial support from FCT (PTDC/QEQCOM/5904/2014 and UID/MULTI/00612/2013).

- [1] J. F. Hunt, P. Rath, K. J. Rothschild, D. M. Engelman, *Biochemistry*, 36:15177 (1997)
- [2] M. Musial-Siwek, A. Karabadzak, O. A. Andreev, Y. K. Reshetnyak, D. M. Engelman, *BBA*, 1798:1041 (2010)
- [3] J. Fendos, F. N. Barrera, D. M. Engelman, *Biochemistry*, 52:4595 (2013)
- [4] D. Weerakkody, A. Moshinoka, M. S. Thakur, V. Moshinoka, J. Daniels, D. M. Engelman, O. A. Andreev, Y. R. Reshetnyak, *PNAS*, 9:5834 (2013).

New intermetallic electrocatalysts for hydrogen production

D. Siopa^a, A. P. Gonçalves^b, J. Branco^b, A. Gomes^a

^aCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^bC²TN, Instituto Superior Técnico, Universidade Técnica de Lisboa, Estrada Nacional 10, 2685-953 Sacavém, Portugal

dasiopa@fc.ul.pt

Clean and sustainable molecular hydrogen (H₂) production can be achieved through electrochemical water splitting [1].

The hydrogen evolution reaction (HER) requires an electrocatalyst which lowers the overpotential and promotes higher exchange current density values. Rare-earth-based AB₂ intermetallic compounds can be seen as potential hydrogen electrocatalysts due to their high activity towards HER [2].

The intermetallics studied were based on the earth abundant, inexpensive Cu element combined with different lanthanides to prepare three distinct compounds – PrCu₂, GdCu₂ and YbCu₂.

These materials were prepared in the form of ingots by induction melting using high-purity metals in the adequate proportion. The melting process was repeated at least three times in order to ensure a perfect homogeneity. Working electrodes were assembled using a PTFE cavity electrode filled with a mixture of powdered intermetallic compound and conductive carbon paste (CP), in a 1:1 mass ratio, for optimal current collection.

In order to assess the electrocatalytic properties of these intermetallics and their activity for the hydrogen evolution reaction (HER) in alkaline medium at different temperatures, electrochemical techniques like linear voltammetry (LV) and chronoamperometry (CA) have been applied. The analysis of the polarization curves allow the determination of the kinetic parameters - Tafel slope, transfer coefficient and exchange current density - which are crucial to understand the hydrogen formation mechanism.

From Figure 1, it could be concluded that the intermetallic compounds can be ordered as PrCu₂ > GdCu₂ > YbCu₂ due to their ability to the hydrogen reaction since a current of $I = -2$ mA is reached at -1.34 , -1.41 and -1.47 V vs Hg/HgO, respectively.




Figure 1. Linear voltammograms recorded for the different electrodes under study at room temperature. Sweep rate of 1 mVs⁻¹.

Acknowledgements:

The authors would like to thank Fundação para a Ciência e Tecnologia for funding through strategic project UID/MULTI/00612/2013 and project PTDC/AAG-TEC/3324/2012.

References

1. C. G. Morales-Guio, L. A. Stern and X. Hu, *Chem. Soc. Rev.* (2014) 43, 6555;
2. M. Tliha, C. Khaldi, S. Boussami, N. Fenineche, O. El-Kedim, H. Mathlouthi and J. Lamloumi, *J. Solid State Chem.* (2014) 18 (3), 577.

Bioactivity and phytochemical studies on Tunisian globe artichoke (*Cynara scolymus*) wastes

Faiza Mejri^{a,b,c,*}, Alice Martins^a, Karim Hosni^c, Hervé Casabianca^d, Amélia P. Rauter^a

^aCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa

^bFaculté des Sciences de Bizerte, TUNISIA

^cInstitut National de Recherche et d'Analyse Physico-Chimique, Biotechpole de Sidi Thebet, TUNISIA

^dInstitut des Sciences Analytiques, Lyon, FRANCE

*e-mail address: faizamejri@hotmail.fr

Globe artichoke (*Cynara scolymus*) is an ancient herbaceous perennial plant original from the Mediterranean Basin, but it is extensively cultivated all over the world. From this vegetable only the inner leaves and hearts are considered edible, whereas external bracts, leaves, and stems are food industrial by-products. These wastes represent more than 60% of the fresh weight, and they are generally used as animal feedstuff and for fiber production or discarded, with additional waste treatment costs in compliance with environmental laws. Over the last years, artichoke is attracting the scientific, industrial, and public communities by its beneficial effects such as antibacterial, antioxidant, anti-HIV, hepatoprotective, urinate, anti-inflammatory, anticarcinogenic, anti-hypercholesterolemic and cardiovascular protective properties.

Analytical studies on the phytochemical profile of Tunisian artichoke wastes showed that phenolics, like caffeoylquinic acid and flavonoids, are major constituents. However, little attention has been paid to their potential as a source of functional ingredients. Therefore, the present work highlights the biological activities of artichoke extracts, giving special attention to the *in vitro* antioxidant, antibacterial, and anti-inflammatory activities of leaves and floral stem residues of Tunisian globe artichoke cultivar “Violet d’Hyères”. Our results indicate that phenolic compounds may be responsible for the strong antioxidant, antimicrobial and anti-inflammatory properties denoted by the studied extracts, fractions, and isolated compounds. So, these wastes can be regarded as a cheap source of new functional ingredients for food and pharmaceutical industries.

Acknowledgements

This work is supported by Fundação para a Ciência e a Tecnologia, (Project Pest-OE/QUI/ UI0612/2013; UID/MULTI/00612/2013).

Why is INH-C₁₀ more active than INH against *Mycobacterium tuberculosis*? Decoding the puzzle...

Filomena Martins^{a*}, Miguel Machuqueiro^a, Peter C. Loewen^b

^aCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Ed. C8, Campo Grande 1749-016 Lisboa, Portugal

^bDepartment of Microbiology, University of Manitoba, Winnipeg, MB Canada R3T 2N2

* filomena.martins@ciencias.ulisboa.pt

Tuberculosis (TB) is still nowadays a serious global threat. Its causative agent, *Mycobacterium tuberculosis* (*Mtb*), has become increasingly resistant to the most effective 1st line antitubercular drugs, in particular to isoniazid, INH (Figure 1). New INH-based compounds have been proposed to circumvent *Mtb* resistance and therefore improve drug activity. To become active against *Mtb*, INH needs to be first activated by the catalase-peroxidase KatG enzyme. Among the most promising compounds, INH-C₁₀ (Figure 1), a new acylated INH derivative was shown to have a similar activity to INH against the wt strain but a MIC value six times lower against a *katG* S315T mutated strain [1]. It was recently observed using MD simulations that this increased activity could not be assigned to changes in the steric environment of the access channel to the heme site, since they were not found to be significant, but rather to subtle electrostatic changes in the vicinity of the heme pocket which will, in the end, influence its reactivity [2].

In an effort to understand the twists of INH and INH-C₁₀ in KatGs, crystallographic studies were undertaken. Having seen no evidence of binding in the soaking experiments with INH-C₁₀ in the enzyme, NBT assays were also carried out to assess the amount of free radicals produced in the first step of the reaction of INH-C₁₀ in KatG. Results showed that this reaction was much slower than that of INH which was not expected from the experimental binding constants and MIC values.

To shed some light into these findings, a computational study was performed where the energetic barriers of radical formation for INH and several acylated INH derivatives (INH-C_x, with $x = 2, 4, 6, 8, 10$) were calculated at the quantum level. In this study, a simple free energy difference between products and substrates in the activation reaction was evaluated. The difference for each acylated compound was compared with INH ($\Delta\Delta G$) and it was noted that it is always positive and nearly constant (around 9-10 kcal mol⁻¹). Therefore, the radical formation in these derivatives is thermodynamically less favoured (deactivated) than in INH and independent on the size of the alkyl chain, assuming that KatG deals similarly with all compounds. These results suggest that, despite its smaller reactivity, the hydrophobic nature of INH-C₁₀ may promote a better trafficking through the *Mtb* membrane, leading to higher concentrations in the vicinity of KatG, thus resulting in lower MIC values.




Figure 1. Optimized structures of INH (left) and INH-C₁₀ (right).

Acknowledgements

The authors are grateful to Fundação para a Ciência e a Tecnologia, Portugal, for financial support under projects PEST-OE/QUI/UI0612/2013 and UID/MULTI/00612/2013.

- [1] F. Martins, S. Santos, C. Ventura, R. Elvas-Leitão, L. Santos, S. Vitorino, M. Reis, V. Miranda, H. F. Correia, J. Aires-de-Sousa, V. Kovalishyn, D.A.R.S. Latino, J. Ramos, M. Viveiros, Eur. J. Med. Chem. 81 (2014) 119-138.
- [2] V.H. Teixeira, C. Ventura, R. Leitão, C. Rafols, E. Bosch, F. Martins, M. Machuqueiro, Mol. Pharm. 12 (2015) 898-909.

Identification of psychoactive substances in seized products in Portugal

Christophe Gonçalves^a, Cláudio R. Queirós^a, Sara Ciríaco^a, M. João Caldeira^b, Helena Gaspar^a

^aCentro de Química e Bioquímica, Faculdade de Ciências da Universidade de Lisboa, Campo Grande, 1749-016 Lisboa

^bLaboratório de Polícia Científica da Polícia Judiciária, Novo edifício-sede da Polícia Judiciária,

Rua Gomes Freire, 169-007 Lisboa, Portugal

chris.goncalves@hotmail.fr

In the past few years, and in particularly in the last 5 years, there has been an uprising of New Psychoactive Substances (NPS) available in “smartshops” and over the internet, with a rate of two new substances detected every week. NPS are psychoactive substances not internationally controlled but that may pose a risk similar to traditional drugs. In 2015, 100 new substances were detected for the first time, bringing the total number of NPS monitored to 568. Now, there are more than twice new substances on the market as compounds controlled under international conventions. Synthetic cathinones and synthetic cannabinoids are the most abundant classes of NPS reported in EU. Since 2005, 160 synthetic cannabinoids and 103 synthetic cathinones have been monitored [1].

Three years ago, a new law has been implemented in Portugal (Dec-Lei 54/2013 de 17 de Abril)[2], which forbids the production and commercialisation of about 159 NPS, being liable to fast updates, in order to keep up with the everyday appearance of new substances.

This project is being developed within the scope of the collaboration between the *Faculdade de Ciências da Universidade de Lisboa* (FCUL) and the *Laboratório de Polícia Científica da Polícia Judiciária* (LPC/PJ), as a result of the need to create effective analytical databases that will facilitate the quick identification of these drugs in a forensic context in Portugal.

This study describes the identification and characterisation of five psychoactive substances (**Figure 1**) detected in seized products in Portugal: three synthetic cannabinoids (STS-135, 5F-AKB-48, 5F-PB-22); one synthetic cathinone (MPHP) and a tryptamine (4-OH-MIPT). Additionally, raw materials such as 8-quinolinol and adamantylamine used in the synthesis of some of those substances were also detected.




Figure 1. Structure of the five psychoactive substances identified in seized product in Portugal.

Using the complementary information from different analytical techniques (1D/2D NMR, GC-EI-MS and ESI-MS), it was possible to determine unequivocally the structures of the psychoactive substances present in the samples without using chemical standards or any kind of purifications. However, in order to use those compounds as qualitative standards for further identifications of NPS, the 3 cannabimimetic substances identified were also isolated and purified from the seized samples.

Acknowledgments

The authors thank FCT for financial support through project UID/MULTI/00612/2013.

[1] European Monitoring Centre for Drugs and Drug Addiction, *European Drug Report 2016 - Trends and Developments*, Publications Office of the European Union, Lisbon, 2016.

[2] Decreto-Lei 54/2013, *Diário da República*, 1.ª série, No. 75, 17 de abril de 2013.

Conducting polymers exhibiting pseudo cationic doping for cathodes in Li-ion batteries

Ariana R. Silva ^a, Jorge P. Correia ^{a*}

^a Centro de Química e Bioquímica, DQB, Faculdade de Ciências, Universidade de Lisboa, Portugal

*jorge.correia@fc.ul.pt

The unique properties of conducting polymers have been explored for a variety of applications and these materials have been well succeeded in many fields [1,2]. Poly(3,4 ethylenedioxythiophene) doped with poly(styrene-4-sulfonate) (PEDOT:PSS) exhibits characteristics like, among others, the porosity and the pseudo-cationic doping, making it an attractive choice for electrodes in Li-ion batteries [3,4]. Besides the stability, the character and easiness of mass transfer, in particular that of the Li⁺, during the charge/discharge processes of the polymer, is a crucial parameter for the performance of such devices [5].

In this work the mass transfer phenomena taking place during the redox conversion of PEDOT:PSS films is investigated by coupling electrochemical methods with microgravimetry (EQCM), Probe Beam Deflection (PBD) and in-situ ellipsometry in two different organic solvents containing LiClO₄. Polymeric films were potentiostatically and potentiodynamically synthesized onto platinum electrodes from aqueous solutions containing the monomer and PSS. The combined information of the PBD and EQCM allows distinguishing the solvent mass fluxes from that of the ions that can be resolved individually. The ellipsometric information acquired both in the course of the synthesis and during the redox conversion process was mathematically treated and fitted to a suitable physical model in order to monitor the film thickness and dielectric constants change imparted by the mass fluxes in the polymer matrix. The data revealed a quite fast quasi-ideal pseudo-cationic doping being the perchlorate anion participation in the conversion process almost negligible. Long runs of charging/discharging conversions of the PEDOT:PSS demonstrated the high stability of electric and mass transfer responses of the modified electrodes.

Acknowledgements

This work is funded by FCT (Fundação para a Ciência e a Tecnologia) through the project UID/MULTI/00612/2013

- [1] PM Dziwowski, M Grzeszczuk, *Electrochim. Acta*, 55 (2010) 3336.
- [2] N Oyama, T Sarukawa, Y Mochizuki, T Shimomura, S Yamaguchi, *J. Power Sources*, 189 (2009) 230.
- [3] AL-Oleksiak, K Kazubowska, A Kupniewska, *J. Electroanal. Chem.*, 501 (2001) 54.
- [4] D Han, G Yang, J Song, L Niu, Alvaska, *J. Electroanal. Chem.*, 602, (2007) 24.
- [5] AL-Oleksiak, A Kupniewska, *Solid State Ionics*, 157 (2003) 241.

Tuning the pore size distribution of mesoporous carbons: the key role of alkaline chlorides in mixtures with zinc chloride

Ana S. Mestre^{a,b*}, Raquel Gonçalves^a, Valentine Vigneron^a, Cristina Freire^b, Ana P. Carvalho^a

^a Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^b REQUIMTE/LAQV, Dep. de Química e Bioquímica, Faculdade de Ciências, Universidade do Porto, 4169-007 Porto, Portugal

*asmestre@fc.ul.pt

Activated carbons are unique materials due to their high adsorption properties, tuneable pore size distribution and rich surface chemistry, being so crucial in various industrial processes. But, the sustainability, stability and versatility of these materials still stimulates continuous efforts in the development of improved or new synthetic routes leading to better performances or even new properties. Particularly, the synthesis of mesoporous carbons has had significant advances in the last decades, due to their importance in applications involving large molecules, such as, adsorbents for dyes, catalyst supports for biomolecules, or as electrodes for biosensors [1].

A recent study reported the synthesis of highly mesoporous carbons from biomass using a eutectic salt mixture KCl/ZnCl₂, attaining $A_{\text{BET}} \approx 1300 \text{ m}^2 \text{ g}^{-1}$ with more than 90 % of mesopores [2]. The present work aims to explore the potentialities of alkaline and zinc chlorides mixtures, as porogens for the control of the mesopore size distribution in glucose-derived activated carbons. The activation was made with eutectic mixtures of LiCl/ZnCl₂ (LiZn), NaCl/ZnCl₂ (NaZn) and KCl/ZnCl₂ (KZn), and also only with ZnCl₂ for comparison purposes. Briefly, 1 g of glucose was milled with 3 g of the salt mixture being activated under N₂ flow (5 cm³ s⁻¹): up to 240 °C (10 °C min⁻¹), which was kept for 2 h, further increased up to 800 or 1000 °C (10 °C min⁻¹), then held for more 2 h. The furnace was cooled (N₂ flow), sample was washed with distilled water until no detection of chlorine ions in the supernatant, dried and stored. Characterization was made by N₂ adsorption isotherms at -196 °C, ash content, determination of the pH at the point of zero charge (pH_{PZC}) and DRIFT.

Salt mixtures play a crucial role in the porosity development as illustrated in the N₂ adsorption isotherms shown in Fig. 1(a). All curves belong to the type IV(a) according to the IUPAC classification [3], being characteristic of mesoporous solid materials. Samples obtained with the eutectic salt mixtures LiZn and NaZn present type H2(b) hysteresis loops associated with pore blocking and broad distribution of neck widths, sample prepared with KZn has a type H4 hysteresis loop usually found in micro-mesoporous carbons.

In all cases superactivated carbons were obtained (total pore volumes ($V_{\text{total}} \geq 1 \text{ cm}^3 \text{ g}^{-1}$ and $1000 < A_{\text{BET}} < 2000 \text{ m}^2 \text{ g}^{-1}$) (Fig. 1(b)). The materials have well developed micro and mesopore networks, with V_{meso} accounting to 45-60 % of the V_{total} when LiZn and NaZn were used, and reaching 80 % when KZn was employed. Sample prepared only with ZnCl₂, in the same amount of ZnCl₂ used in the eutectic mixture KZn, has the double of the A_{BET} but only 18 % of V_{meso} , proving that alkaline metals play a fundamental role in the development of pores in this methodology. The results obtained so far reveal that the mesopore size distribution can be tuned by the salt mixture: LiZn leads to materials with high V_{meso} with small diameters (2 – 10 nm; NaZn originates mesopores up to 20 nm, and KZn results in high volume of large mesopores (20 – 50 nm).

Acknowledgements

This study was funded by projects PEst-OE/QUI/UI0612/2013 (CQB) and UID/QUI/50006/2013 - POCI/01/0145/FERDER/007265 (REQUIMTE) from FCT/MEC through national funds and co-financed by FEDER, under the Partnership Agreement PT2020. ASM thanks FCT for Post-doc grant (SFRH/BPD/86693/2012).

[1] J. Lee, J. Kim, T. Hyeon, Adv. Mater. 18 (2006) 2073-2094.

[2] Z. Ma, H. Zhang, Z. Yang, Y. Zhang, B. Yu, Z. Liu, J. Mater. Chem. A 2 (2014) 19324-9.

[4] M. Thommes, K. Kaneko, A.V. Neimark, J.P. Olivier, F. Rodriguez-Reinoso, J. Rouquerol, K.S.W. Sing, Pure Appl. Chem. 87 (2015) 1051-1069.




Figure 1. Textural characterization of materials obtained at 800 °C with the mentioned eutectic mixtures (a) N₂ isotherms; (b) textural parameters (bars: volumes; line: A_{BET}).

Influence of abiotic effects on antioxidant and polyphenol content of some economically valuable Mediterranean crops

Basma Mabrouk^a, Barbara Oliveira^b, Leila Bettaieb Ben Kaab^a, Maria Eduarda Araújo^{b*}

^a Faculty of Sciences of Tunisia, University Tunis El Manar, Tunis, Tunisia

^bCQB and Department of Chemistry and Biochemistry, Faculty of Sciences, University of Lisbon, 1749-016 Lisboa, Portugal

*mearaujo@fc.ul.pt

Many economically valuable plants are used in folk medicine due to its antioxidant properties which are usually a consequence of its high content in phenolic compounds. However, the antioxidant properties depicted by many medicinal plants are not an intrinsic property but are usually highly affected by abiotic factors.

Fenugreek (*Trigonella foenum-graecum*) is an annual Leguminosae crop, cultivated throughout the world especially in India, in Middle Eastern countries, and widely in the Mediterranean basin including Tunisia. It has many effects on health. Leaves and seeds of Fenugreek are used in ethnomedicine due to its anti-oxidant, antidiabetic, antimicrobial, anti-inflammatory, etc, activities. Nowadays, contamination by heavy metals (such as arsenic, zinc, cadmium, cobalt, aluminium, lead...) has adverse effects on the mobilization of mineral, organic reserves and growth of plants and causes many problems on health. In this communication it is presented the effect of zinc and arsenic on the antioxidant properties and phenol content of methanol extracts of fenugreek cultivated under these metal stress and also the effect of salicylic acid used to counteract these adverse conditions. Antioxidant properties were investigated following the bleaching of the DPPH radical during 60 minutes (Figure 1). As it can be seen in Figure 1 all extracts were mildly active, have either been prepared from the leaves or the roots. Metal stress (ZR, AR, ZF and AF extracts) increased the production of antioxidant compounds comparing with non treated plants (TR and TF). Polyphenol content was highly affected by treatments and results will be presented and discussed.

Sumac, *Rhus coriaria* L. is another important plant that grows in the Mediterranean basin. Is a small shrub from the Anacardian family whose fruits are used as condiment and whose leaves were traditionally milled and used in leather production and as a mordant in textile dyeing due to its high content in gallotannins. Two samples of sumac, one collected in North of Portugal and the other a commercial one from Italian origin were compared (Table 1).

Tannin content was obtained integrating the ATR-FTIR absorbance band at 1209 cm⁻¹ of aqueous solutions of tannic acid with several concentrations. The band at 1209 cm⁻¹ is a band common to all tannins, and very important in gallotannin spectra (Figure 2) [1].

Sumac, *Rhus coriaria* L. is another important plant that grows in the Mediterranean basin. Is a small shrub from the Anacardian family whose fruits are used as condiment and whose leaves were traditionally milled and used in leather production and as a mordant in textile dyeing due to its high content in gallotannins. Two samples of sumac, one collected in North of Portugal and the other a commercial one from Italian origin were compared (Table 1).

Tannin content was obtained integrating the ATR-FTIR absorbance band at 1209 cm⁻¹ of aqueous solutions of tannic acid with several concentrations. The band at 1209 cm⁻¹ is a band common to all tannins, and very important in gallotannin spectra (Figure 2) [1].




Figure 1. Bleaching of DPPH radical by methanol extracts of leaves and roots of non treated (TR, TF) and treated plants




Figure 2. ATR-FTIR spectra of tannic acid and sumac

Table 1. Antioxidant, total phenolic content and tannin content of sumac

Sumac origin	EC ₅₀ (mg/mL, DPPH test)	TPC (GAE)*	TC**
Portuguese	4.2	62.7	558
Commercial	9.3	39.9	386

* TPC: total phenolic content; GAE: galic acid equivalents, mg/g extract; **TC: tannin content, tannic acid equivalents mg/g extract

Acknowledgements

This work was founded by Fundação para a Ciência e Tecnologia (FCT) Portugal (UID/MULTI/00612/2013).

[1] L. Falcão, M. E. M. Araújo, Vibrational Spectroscopy 74 (2014) 98.

Step by step development of a lipid-based immunosensor platform with high capability to inhibit nonspecific adsorption

Telmo O. Paiva^{a*}, Inês Almeida^a, Joaquim T. Marquês, Yan-Yan Chen^b, Niu Yu^c, Gang Jin^c, Rodrigo F. M. de Almeida^a, Ana S. Viana^a

^aCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^bSuzhou Institute of Nano-Tech and Nano-Bionics (SINANO), Chinese Academy of Sciences, 215123, Suzhou, China

^cInstitute of Mechanics, Chinese Academy of Sciences, 100190, Beijing, China

*tmpaiva@fc.ul.pt

To detect the presence of a particular bioanalyte (e.g. a tumor biomarker), usually in low concentration and in complex matrices, a biosensor with both high affinity and selectivity must be employed, avoiding nonspecific interactions between those molecules and the biosensor components [1]. These requirements can be achieved using immunosensor interfaces, where the sensitive biological elements are antigens or antibodies immobilized on a surface and the formation of the complex antigen-antibody generates a measurable signal [2]. Regarding nonspecific interactions, our recent work has given support to the developing notion that lipid biointerfaces are very effective to suppress them, and that this feature, allied with their biocompatibility, represents an additional advantage to the development of immunosensing devices [3].

Here, we present the design and assembly of a highly organized and stable immunosensing interface with special capability to inhibit the nonspecific adsorption of proteins, enhancing the signal generated by the recognition event and thus the limit of detection. Several characterization techniques were employed during the implementation of this bottom-up approach. Conventional ellipsometry was used to estimate gold substrates thickness as well as for the screening of each step of the immunosensor development. Atomic force microscopy allowed the direct visualization of the modified surfaces during the process. Total internal reflection imaging ellipsometry, as well as surface plasmon resonance, were selected to perform the biosensing assays with the antibodies in order to evaluate the performance of the platform developed.

The developed interface consists of a gold surface modified with a self-assembled monolayer of 11-amino-1-undecanethiol, which contains terminal amine groups that were used to covalently link modified lipid molecules (1-myristoyl-2-(14-carboxymyristoyl)-sn-glycero-3-phosphocholine), through the terminal carboxylic groups previously activated. 2-hydroxyoleic acid molecules were added to the lipid monolayer to provide oriented carboxylic groups to covalently link Immunoglobulin G (IgG). The performance of the interface was evaluated by the real-time detection of Anti-IgG-IgG complex formation.

Acknowledgments

Fundação para a Ciência e a Tecnologia (FCT) is acknowledged for funding the projects: PEst 2015-2020 (UID/Multi/00612/2013), IF/00808/2013/CP1159/CT0003, IF2012/2013 initiatives (POPH, FSE), 7th Sino-Portugal S&T Cooperation 2013-2015, PTDC/CTM-NAN/0994/2014.

[1] S. Gs, A. Cv, and B. B. Mathew, *Journal of Instrumentation Technology*, 2 (2014) 26-39.

[2] Y. Niu, A. I. Matos, L. M. Abrantes, A. S. Viana, and G. Jin, *Langmuir*, 28 (2012) 17718-25.

[3] I. Almeida, J. T. Marques, W. Liu, Y. Niu, R. F. de Almeida, G. Jin, A. S. Viana, *Colloids Surf B: Biointerfaces*, 136 (2015) 997-1003.

Thermochemistry of peroxy radicals: a combined photoacoustic calorimetry, quantum chemistry and molecular dynamics study

Rui M. Borges dos Santos^{a,b,*}, Elsa M. Gonçalves^a, Filipe Agapito^a, Nuno Galamba^b, José A Martinho Simões^a

^aCentro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^bCCMAR and CBMR, Universidade do Algarve, Campus de Gambelas, 8005-139 Faro, Portugal

*rmsantos@fc.ul.pt

Reaction 1, where $R^\bullet$ denotes a carbon radical and $ROO^\bullet$ the product peroxy radical, is a key step in the interaction mechanism of oxygen with organic matter, hence its enormous importance. Its study is essential, for instance, to optimize the production of energy (and thus limit emissions) from internal combustion engines, or to develop antioxidants to avoid the deterioration of cellular membranes due to lipid peroxidation.



To fully describe the thermodynamics of reaction 1, we need to know the stability of both the carbon and peroxy radicals. However, the energetics of radicals can be notoriously difficult to study. While the thermodynamic stability of long-lived organic molecules is well established for a large number of substances [1], only a few hundreds of carbon radicals are listed in a recent compilation of thermochemical data [2], and the situation is particularly critical for peroxy radicals: the same compilation includes less than 30! The need for further research on the subject is evident but, to study such demanding species, one needs to use advanced experimental or theoretical techniques (or preferably, both).

Time-resolved photoacoustic calorimetry (TR-PAC) is a recent technique suited to study radical reactions, in which they are started with a laser (photo-) and then followed via a microphone (-acoustic). In this way the radicals can be “temporally isolated”, which allows probing their thermochemistry with chemical accuracy (ca. 5 kJ/mol). Regarding computational chemistry methods, the accuracy of composite approaches together with the explicit treatment of electronic correlation [3] is comparable to the best conventional methods but, unlike those, it can be used for much “larger” systems (in electron number, such as the peroxy radicals). Finally, molecular dynamics (MD) is essential for the interpretation of the above results at a molecular level, bridging the experimental and electronic structure data.

In this work we will present our methodology to study the above reaction, together with experimental and computational results for the thermochemistry of selected carbon and peroxy radicals. The interpretation of the TR-PAC results for peroxy radicals is complicated by some still poorly understood issues, namely the volume change and the solvation effects of reaction 1. Both these problems can be conveniently addressed by MD. Still, fundamental aspects of this procedure need to be addressed, such as the development of suitable force-fields for the radicals. These subjects, which were neglected in previous studies of reaction 1 [4], will also be discussed and presented as part of our complete and accurate methodology for the experimental and computational study of the thermochemistry of peroxy radicals.

Acknowledgements

Support for this work was provided by Fundação para a Ciência e a Tecnologia (FCT), Portugal, through CQB PEST-UID/MULTI/00612/2013. E.M.G., F.A. and N.G. acknowledge financial support from FCT post-doctoral grants (SFRH/BPD/88607/2012, SFRH/BPD/74195/2010, and SFRH/BPD/110483/2015, respectively).

- [1] P. J. Linstrom, W. G. Mallard. NIST Chemistry WebBook, NIST Standard Reference Database Number 69, National Institute of Standards and Technology Gaithersburg MD, <http://webbook.nist.gov>.
- [2] Y.-R. Luo, Comprehensive Handbook of Chemical Bond Energies. CRC Press: Boca Raton, 2007.
- [3] A. Karton, J. M. L. Martin, J. Chem. Phys. 136 (2012) 136 124114.
- [4] M. Kranenburg, M. V. Ciriano, A. Cherkasov, P., Mulder, J. Phys. Chem. A 104 (2000) 915-921.

The impact of ligand derivatisation on Mn(III) thermal spin transitions

Patrícia S. Arranja^{a*}, Liliana P. Ferreira^{b,c}, Sónia Barroso^d, Maria José Calhorda^a, Paulo N. Martinho^a

^a Centro de Química e Bioquímica, DQB, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^b BioISI, Faculdade de Ciências, Universidade de Lisboa, Campo Grande, 1749-016 Lisboa, Portugal

^c Departamento de Física, Faculdade de Ciências e Tecnologia, Universidade de Coimbra, 3004-516 Coimbra, Portugal

^d Centro de Química Estrutural, Instituto Superior Técnico, Universidade de Lisboa, Avenida Rovisco Pais, 1049-001 Lisboa, Portugal

*isabelpassoares@gmail.com

Spin crossover (SCO) compounds which can switch between two magnetic states are magnetic materials with good potential for incorporation into spintronics.[1] Magnetism in transition metal ions arises from unpaired spins residing in *d*-orbitals. Careful ligand design produces molecules able to reversibly switch between the two possible electronic spin states, HS and LS. Switching between different magnetic states in a single molecule is then achievable by heat, light or pressure stimulus. Hysteretic SCO complexes represent a very rare example of bistability in single molecules which normally have just one stable electronic state under ambient conditions.[2]

In Mn(III) complexes with $N_4(O^-)_2$ donating Schiff-bases the modulation of the orientation of the $N_4(O^-)_2$ donors in a series of hexadentate ligands had a dramatic effect on the spin state of their Mn(III) complexes. The continuous search for new examples of SCO complexes with spin transitions around room temperature, and also for systems showing a high degree of cooperativity, led us to further investigate new examples of d^4 Mn(III) complexes with this ligand series. Like Fe(III), Mn(III) also may present two paramagnetic spin states with LS ($S = 1$) and HS ($S = 2$) states but with permanent distortion in the HS state due to a strong Jahn-Teller effect.

We report a series of complexes derived from the polyamine 1,2-bis(3-aminopropylamino)ethane which reacted with salicylaldehyde to form the hexadentate Schiff-base ligands. This ligand formed the mononuclear Mn(III) complexes by reaction with different $MnY_2 \cdot xH_2O$ salts ($Y = BF_4^-$, ClO_4^- , BPh_4^- and PF_6^-) after air oxidation. All compounds were characterized by FTIR, UV-Vis and NMR spectroscopies, elemental analysis and single crystal X-ray crystallography. Magnetic measurements on all compounds were also performed.




Figure 1 – Schiff-base ligand used in this study (left) and single crystal X-ray structure of the Mn(III) BF_4^- salt (right).

Acknowledgements

The authors would like to thank Fundação para a Ciência e Tecnologia for financial support UID/MULTI/00612/2013 and PTDC/QUE-QIN/3414/2014. PNM thanks FCT for financial support (SFRH/BPD/73345/2010).

[1] Feringa, B. L. *Molecular Switches*; Wiley-VCH Verlag GmbH, 2001.

[2] Martinho, P. N., Gildea, B., Harris, M. M., Lemma, T., Naik, A. D., Müller-Bunz, H., Keyes, T. E., Garcia, Y., Morgan, G. G., *Spin Transitions*, 2012, 51, 12597-12601.

Recovery of palladium from a spent industrial catalyst: A hydrometallurgical approach

A. P. Paiva^{a*}, P. P. Abreu^a, C. A. Morgado^a, G. I. Carvalho^a, O. Ortet^{a,b}, C. A. Nogueira^c

^a Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal

^b Departamento de Ciência e Tecnologia, Universidade de Cabo Verde, Praia, Cabo Verde

^c Laboratório Nacional de Energia e Geologia (LNEG), I.P., Estrada do Paço do Lumiar, 22, 1649-038 Lisboa, Portugal.

* appaiva@ciencias.ulisboa.pt

Platinum-group metals (PGMs) are nowadays a trademark of several top technological devices, their unique properties being extensively explored in medicine applications, in the manufacture of electronic materials and fuel cells, and particularly as catalysts (in automotive catalytic converters, oil refining and fine chemistry processing). To prevent ore exhaustion due to current and extensive industrial demand, since mineral deposits of PGMs are scarce in the earth surface, the development of recycling practices suited to end-of-life materials (the so-called urban mining) has become a real necessity [1].

In this communication the development of an integrated hydrometallurgical treatment applied to a spent hydrogenation catalyst from a Portuguese petrochemical industry – Figure 1 – is described. The spent catalyst, composed by alumina with 0.03% palladium and 0.03% chromium, has been leached under different experimental conditions, for which the parameters taken into account were the nature and concentration of the leaching agents, temperature, time, liquid/solid ratio (L/S) and particle size. The system showing the maximum Pd solubilisation and the minimal dissolution of Al and Cr consisted on 2M HCl and 1M H₂O₂, 27°C, 10 min, L/S=2 L/kg and an average 176 µm of particle size – Table 1.



Figure 1. Spent catalyst sample used in this work.

Table 1. Composition of the “best” leaching solution.

	[Pd] / mgL ⁻¹	[Al] / gL ⁻¹	[Cr] / mgL ⁻¹
2M HCl +1M H ₂ O ₂	63.00 ± 2.00	2.92 ± 0.10	0.40 ± 0.05

The leaching solution was subject to solvent extraction (SX) with two thiodiglycolamide derivatives in toluene. It is well known that *N,N'*-dimethyl-*N,N'*-dicyclohexylthiodiglycolamide (DMDCHTDGA) [2] and *N,N'*-dimethyl-*N,N'*-dibutylthiodiglycolamide (DMDBTDGA) [3] are able to efficiently and selectively extract Pd(II) from HCl solutions, 0.1M thiourea in 1M HCl being used as stripping agent to transfer the PGM to new aqueous phases.

The practical usefulness of these SX systems has been accessed by: a) DMDCHTDGA and DMDBTDGA consecutive Pd extraction-stripping cycles with the leaching solution, to test their robustness upon reutilization; b) The plot of the correspondent equilibrium extraction isotherms, to evaluate the maximum Pd(II) loading capacity. The overall results obtained show that both thiodiglycolamide derivatives are re-utilizable, maintaining their efficiency for Pd(II) recovery after several cycles, although the progressive accumulation of Al in the solvents may be a drawback. The co-extraction of Al also affects the loading capacity of the organic solvents towards Pd(II), since they are not so high as expected, due to the negative interference of Al.

Acknowledgements

Support for this work was provided by FCT [UID/MULTI/00612/2013], and [O. Ortet] acknowledges the financial support through his PhD grant as well [SFRH/BD/78289/2011]. Thanks are also due to Repsol Polímeros SA [Eng. André Vilelas] for the spent catalyst sample.

- [1] S. Steinlechner, J. Antrekowitsch, JOM 67 (2015) 406–411.
- [2] O. Ortet, A.P. Paiva, Sep. Purif. Technol. 156 (2015) 363–368.
- [3] O. Ortet, M.E. Martins, A.P. Paiva, Metals 5 (2015) 2303–2315.

“Healthy Life”: Interaction of polyphenols with lipid bilayers and their effects in human cells

Hugo A. L. Filipe^{a*}, Catarina Peneda^a, Joaquim T. Marquês^a, Miguel Machuqueiro^a, João C. Ramos^a, Maria da Soledade Santos^a, H. Susana Marinho^a, Ana S. Viana^a, Helena Soares^{a,b}, Rodrigo F. M. de Almeida^a

^aCentro de Química e Bioquímica, Departamento de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, Ed. C8, Campo Grande, 1749-016 Lisboa, Portugal

^bEscola Superior de Tecnologia da Saúde de Lisboa, IPL, 1990-096 Lisboa, Portugal

**hafilipe@fc.ul.pt*

This work concerns the transversal project of the CQB thematic line: “Healthy Life: Molecular Interventions and Regulation Mechanisms”.

Biologically active plant phytochemicals have a broad range of pharmacological effects including anticarcinogenic, antimicrobial, antioxidant, and anti-inflammatory activity. [1] Notwithstanding the possibility of having a specific target, phytochemicals must interact and permeate through cell membranes in the body. Indeed, it was suggested that those molecules insert into the membranes and thereby may have a promiscuous activity by changing structural properties of lipid bilayers. [2]

Some well-known phenolic acids such as caffeic (CA), rosmarinic (RA) and chlorogenic (CGA) acids, whose identification in plant extracts has been achieved by CQB research groups, were selected to be addressed in first place (Figure 1).

All the phenolic acids studied have low lipophilicity and among them, RA was the only one with a partition to biological membrane models measurable by fluorescence spectroscopy, as opposed to CA and CGA. Cyclic voltammetry measurements using an electrode modified with a supported lipid bilayer, also indicated a higher affinity of RA to lipid membranes. In addition, oxidation/reduction of the phenolic acids displayed higher reversibility in the lipid milieu than in the aqueous bulk. Indeed, the reduced form of phenolic acids was unstable in aqueous solution. In particular, in DMEM/F-12 cell culture media, a colour change observed after incubation with each compound could be reverted by the addition of a reducing agent. The higher reversibility of phenolic acids oxidation/reduction, once they were inserted in the lipid membrane, may contribute to the stability of the compounds and prevent the formation of degradation products. Molecular dynamics (MD) simulations are being performed to probe the location and orientation of these and other selected compounds in lipid bilayers.

The influence of the phenolic acids in the cytoskeleton organization, both actin filaments and microtubules, of a human retinal pigment epithelial cell line (RPE1) was also investigated. All compounds induced concentration and time dependent effects, translated in structural alterations mainly at the cell periphery, and also in the perturbation of cell division. Moreover, it was not evident that these compounds induce apoptosis under the conditions tested. RA seemed to induce evident effects at earlier times and at lower concentrations, as compared to CA and CGA. This higher sensibility of RPE1 cells to RA correlates with the higher affinity of this compound to the lipid bilayer.





Figure 1. Effects in the actin filaments (red) and microtubules (green) of RPE1 cells, after 8 h incubation with 100 μ M of phenolic acid.

Acknowledgements

Support for this work was provided by F.C.T. through IF2012/2013 initiatives (P.O.P.H., F.S.E.) and Project UID/MULTI/00612/2013. H.A.L.F. acknowledges post-doc fellowship under the same Project. M.L. Serralheiro is acknowledged for the phenolic acids.

- [1] A. Stefansson and M. Bakovic, *Nutrients* 6 (2014) 3777-3801.
[2] H. I. Ingólfsson, P. Thakur, K. F. Herold, E. A. Hobart, N. B. Ramsey, X. Periole, D. H. de Jong, M. Zwama, D. Yilmaz, K. Hall, T. Maretzky, H. C. Hemmings, C. Blobel, S. J. Marrink, A. Koçer, J. T. Sack and O. S. Andersen, *ACS Chem. Biol.* 9 (2014) 1788-17



“Designing” porous networks for water treatment: Removal of atenolol and antibiotics by activated carbons

Susana Marques^{a,*}, Stephane Gonçalves^b, Ana S. Mestre^a, Andreja Z. Gotvajn, Ana P. Carvalho^a

^a*Centro de Química e Bioquímica, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisboa, Portugal*

^b*Université Pierre et Marie Curie – 4 Place Jussieu, 7500 Paris, France*

^c*Faculty of Chemistry and Chemical Technology, University of Ljubljana, Slovenia*

**scrmarques@fc.ul.pt*

Continuous release of antibiotics into the environment could exert pressure on ecosystem by developing microbial antibiotic resistance leading to public health important issues. These drugs are only partially absorbed by the body, reaching domestic wastewater and, since the conventional wastewater treatment processes are ineffective in removing these compounds, they are continuously introduced in the aquatic system [1-3]. To solve this problem, advanced technologies, such as adsorption onto carbon materials, have been developed. However, high cost restricts the application of these materials, boosting researchers to explore the preparation of carbons from waste materials or renewable sources [4].


The objective of this work was to prepare activated carbons from the char of apple tree bark residues and evaluating the influence of different experimental conditions in the textural properties of the materials. Selected samples were tested as adsorbents of a β -blocker medicine (atenolol), and two antibiotics (tiamulin fumarate and oxytetracycline hydrochloride).

Acknowledgements

The authors thank FCT, Portugal, for financial support to CQB center (Project UID/MULTI/00612/2013) and fellowships to SM (SFRH/BD/91767/2012) and ASM (SFRH/BPD/86693/2012). Salmon & Cia (Norit) and Quimitejo are also acknowledged by kindly supplying of the commercial carbon samples.

REFERENCES

- [1] A.M. Pereira, L.J. Silva, L.M. Meisel, A. Pena, *J Toxicol Env Health* 78 (2015) 959-975.
- [2] J. L. Sotelo, A. Rodríguez, S. Álvarez, J. García, *Int J Environ Res* 6(4) (2012) 961-968.
- [3] M. Pomorska-Mól, C. Kowalski, *J Liq Chromatogr* R T 32 (2009) 1023-1031.
- [4] A.S. Mestre, E. Tyszko, M.A. Andrade, M. Galhetas, C. Freire, A. P. Carvalho, *RSC Adv* 5 (2015) 19696-19707.



Science at CQB

Index of Authors



Abreu, P. P.	P79
Aço, J.	P40
Agapito, F.	P77
Ahmad, S. M.	O8
Albuquerque, T.	P27
Aldeias, J. B.	P46; P73
Aleixo, R.	O4
Almeida, I.	P13; P76
Almeida, J. C.	P30; P39
Almeida, L. C.	P4
Almeida, R. F. M. de	O2; P3; P10; P24; P32; P34; P76; P80
Alves, A. F.	O6
Antunes, C.	P33
Antunes, C. A.	P3
Antunes, C. A. C.	P10; P24
Antunes, F.	P23; P28; P29
Antunes, M.	P27
Araújo, M. E. M.	P75
Arranja, P.	P78
Ascensão, L.	P67
Assunção, A.	P14
Assunção, R. M.	P21
Atti, N.	P38; P57

Bandeira, N. A. G.	P39; P59
Barata, S.	P53
Barrocas, B.	P47
Barros, J. M.	P2
Barroso, S.	P30; P59; P78
Bastos, A. E. P.	P10
Batista, D.	P16; P27
Bettaie, L.	P75
Bento, R.	P23
Bernardes, C.	P23; P60
Bordado, J. C. M	P36
Borges, C.	P40
Branco, J.	P65
Brandão, P.	P29
Brigas, A.	O4
Brito, E.	P67
Bronze, M. R.	P52
Bronze, S.	P7

Cachatra, V.	O10; P27
Caldeira, M. J.	P69
Calhorda, M. J.	P9; P18; P30; P39; P43; P44; P46; P59; P73; P78
Câmara, A.I.	O9
Camelo, C.	O9; P50
Carabineiro, S. A. C.	P11
Cardoso, B. P.	P44
Cardoso, J. C.	P54
Carmona. B.	O9
Carreira, A. C.	P32
Carvalho, A. P.	O4; O5; P17;

Carvalho, G. I.	P79
Carvalho J.	P19
Carvalho, M. D.	O6; P9; P13; P39
Casabianca, H.	P66
Cepeda, A,	P50
Chen, Y-Y	P76
Ciríaco, S.	P69
Coelho, J.	P22
Colabufo, N.	O10
Condinho, A.	P18
Correia, J. P.	P1; P4; P21;

Costa, G.	P50
Costa, H. S.	P27
Costa, P. J.	P12; P16; P20;
P43	
Coyaud, E.	O9
Cristóvão, J.	P48
Cruz, M. M.	O6
Csuk, R.	P16
Cyrne, M. L.	P10; P50

Delgado, I.	P22
Delgado, M.	P25
Dias, C.	P27; P56
Dias, R.	O5; P35
Dias, R. O.	P12
Diniz, J.	P61
Diogo, H.	P60
Duarte, T. A. G.	P72

Martins, R.	P31	Pais, J. P.	P35
Martins, V. J.	P16	Paiva, A. P.	P79
Matos, O.	P25	Paiva, T. O.	P37; P76
Mejria, F.	P66	Peneda, C.	O2; O9; P80
Melato, A. I.	P44	Pereira, I.	P45
Mendo, S. G.	O6	Pereira, L. C. J.	P59
Mendonça, M. H.	O6	Petit, C.	P17
Mestre, A. S.	O5; P17; P74	Piedade, M. E. M.	P23; P60
	P81	Piedade, M. F. M.	P60
Minhalma, M.	P15	Pinto, F.	O9; P50
Moita, M. L. C. J.	P8	Pinto, M. L.	P28; P29; P49;
Moiteiro, C.	P25		P62
Monteiro, O. C.	P47; P51; P54;	Pinto, R. V.	P29
	P61; P73	Pires, J.	P28; P29; P49;
Moreira, F. M.	P3		P62
Moreira, F. P.	P24	Pombeiro, A. J. L.	P11; P41; P64;
Morgado, C. A.	P79		P72
Morgan, G. G.	P59	Porcheron, A.	P16
Mota, P.	P73		
Mourato, A.	O3		
		Q	
N		Queirós, C. R.	P69
Neugebauer, P.	P59		
Neng, N. R.	O8	R	
Nobre, L. C. S.	P8	Ramgi, P. C.	P44
Nogueira, C. A.	P79	Ramos, J. C.	O ;P802
Nogueira, J. M. F.	O8	Raught, B.	O9
Nolasco, S.	P22	Rauter, A. P.	O10; P2; P5;
Novo, M.	P25		P27; P35; P48;
Nunes, C. D.	O1;P17; P61;		P56; P66
	P71	Real, C.	P50
Nunes, N.	O4; P34	Realista, S.	P30; P39; P43;
Nunes, R.	P5		P44; P46; P59;
Nunes, R. J.	P6; P34		P73
		Reis, J. C.R.	P55
O		Reis, P. B. P. S.	P58
Oliveira, B.	P75	Rendas, M. A.	P23
Oliveira, M. L.	O6	Ressaissi, A.	P38; P57
Ortet, O.	P79	Ribeiro, A. P. C.	P11
Osawa, R. A.	P52	Ribeiro, L.	P26
Ottaviani, D.	P64	Rijo, P.	P36
		Rocha, D.	P25
P		Rodrigues, V. H. N.	P9
Pacheco, M.	O5	Rosa, R. J. M.	P20
Pacheco, R.	O7; P7; P14;		
	P15; P19; P31;	S	
	P38; P42; P53;	Sampaio, R. S.	P1
	P57; P67	Sanches, A.	P27
Pais, J.	P5	Santos, A. F. S.	P8
		Santos, F. C.	P24

107