

# ARE AIR-BORNE MYCOTOXINS A PUBLIC HEALTH CONCERN IN PORTUGAL?

C Viegas<sup>1</sup>, S Viegas<sup>1,2</sup>, R Sabino<sup>3</sup>, E Casimiro<sup>4</sup>, C Veríssimo<sup>3</sup>

<sup>1</sup> Escola Superior de Tecnologia da Saúde de Lisboa - IPL; <sup>2</sup>CIESP – Centro de Investigação e Estudos em Saúde Pública, Escola Nacional de Saúde Pública, ENSP, Universidade Nova de Lisboa; <sup>3</sup>Instituto Nacional de Saúde Dr. Ricardo Jorge - Laboratório de Referência de Doenças Sistémicas e Zoonoses, Departamento de Doenças Infecciosas; <sup>4</sup> INFOTOX Environmental Risk Consultants, Lisbon Portugal

## Introduction

Microscopic filamentous fungi, under suitable environmental conditions, can lead to the production of highly toxic chemical substances, commonly known as mycotoxins. The most widespread and studied mycotoxins are metabolites of some genera of moulds such as *Aspergillus*, *Penicillium* and *Fusarium*. Quite peculiar conditions may influence mycotoxin biosynthesis, such as climate, geographical location, cultivation practices, storage and type of substrate. Toxicity has been extensively investigated for the most important mycotoxins, such as aflatoxins, ochratoxin A and *Fusarium* toxins, and much information derived from toxicokinetics in animal models has also been obtained. The adverse effects are mainly related to genotoxicity, carcinogenicity, mutagenicity, teratogenicity and immunotoxicity.

## Aim of the Study

To identify fungal species able to produce important mycotoxins in different Portuguese settings.

## Methodology

Descriptive studies were developed to monitor air fungal contamination in different settings such as 10 gymnasiums with swimming pools, two elementary schools, one maternity, one hematological unit, 10 hospitals' food units, two companies' food units and one poultry.



Fig. 1 Air sampler

Air samples were collected through impaction method. Surface samples were collected by using pre-moistened swabs and a 10 by 10 cm square stencil.

## References

- Brandi, G, Sisti, M, Paparini, A, Gianfranceschi, G, *et al.*, Swimming pools and fungi: An environmental epidemiology survey in Italian indoor swimming facilities. *International Journal of Environmental Health Research*. 17 (2007) 197 – 206.
- Daisey, J, Angell, W, Apte, M., Indoor air quality, ventilation and health symptoms in schools: an analysis of existing information. *Indoor Air*. 13 (2003) 53 – 64.
- Groopman, J, Kensler, T, Wild, C, Protective Interventions to Prevent Aflatoxin-Induced Carcinogenesis in Developing Countries. *Annu. Rev. Public Health*. 29 (2008) 187–203.
- Hedayati, M, Pasqualotto, A, Warn, P, Bowyer, P, Denning, D, *Aspergillus flavus*: human pathogen, allergen and mycotoxin producer. *Microbiology*. 153 (2007) 1677–1692.
- Hodgson, M, Morey, P, Leung, W, Morrow, L, Miller, D, Jarvis, B, Robbins, H, Halsey, J, Storey, E, Building associated pulmonary disease from exposure to *Stachybotrys chartarum* and *Aspergillus versicolor*. *J Occup Environ Med*. 40, (1998) 241 – 249.
- Hoog, C, Guarro, J, Gene, G. Figueiras, M. (2th ed). (2000). Atlas of Clinical Fungi. Centraalbureau voor Schimmelcultures.
- London, K, Coke, A, Burne, J, Shaw, A, Oppenheim, B, Morris, C, Kitchens as a source of *Aspergillus niger* infection. *J Hosp Infect*. 32 (1996) 191 – 198.
- Lues, J, Theron, M, Venter, P, Rasephei, M, Microbial composition in bioaerosols of a high-throughput chicken-slaughtering facility. *Poultry Science*. 86 (2007) 142 – 149.
- Lugaskas, A, Filamentous fungi isolated in hospitals and some medical institutions in Lithuania: *Indoor Built Environm*. 13, (2004)101 – 108.
- Rao, C, Burge, H, Chang, J, Review of quantitative standards and guidelines for fungi in indoor air. *J Air Waste Manage Assoc*. 46 (1996) 899 – 908.
- Sarica, S, Asan, A, Otkun, M, Ture, M, Monitoring indoor airborne fungi and bacteria in the different areas of Trakya University Hospital, Edirne, Turkey. *Indoor Built Environment*. 11 (2002) 285 – 292.
- Sedmíková, M, Reisnerová, H, Dufková, Z, Bartá, I, Jílek, F, Potential hazard of simultaneous occurrence of aflatoxin B1 and ochratoxin A. *Vet. Med*. 46 (2001) 169-174.
- Speijers, G, Speijers M, Combined toxic effects of mycotoxins. *Toxicology Letters*. 153 (2004) 91–98.
- Sun, C, Wang, L, Chen, C, *et al.*, Genetic polymorphisms of glutathione S-transferases M1 and T1 associated with susceptibility to aflatoxin-related hepatocarcinogenesis among chronic hepatitis B carriers: a nested case-control study in Taiwan. *Carcinogenesis*. 22 (2001) 1289-1294.
- Wang, J, Groopman, J, DNA damage by mycotoxins. *Mutation Research*. 424 (1999) 167–181.
- White, P, Collins, J, McGill, K, Monahan, C, O'Mahony, H., Distribution and prevalence of airborne microorganisms in three commercial poultry processing plants. *J. Food Products*. 64 (2001) 388 – 391.

## Results

Air and surfaces of seven different types of settings were analyzed in order to detect fungal species potential producers of mycotoxins (Table 1).

Table 1. Results obtained on air and surfaces

| Setting                                   | Most frequent fungi  | <i>Aspergillus</i> species  |
|---|--|---|
| Gymnasiums with swimming pools (Air)      | <i>Cladosporium</i> sp. (36.6%)<br><i>Penicillium</i> sp.(19.0%)   | <i>A. flavus</i> , <i>A. niger</i> , <i>A. glaucus</i> , <i>A. fumigatus</i> , <i>A. parasiticus</i> , <i>A. restrictus</i> and <i>A. sydowii</i>   |
| Elementary schools (Air)                  | <i>Cladosporium</i> sp. (52.2%)<br><i>Penicillium</i> sp. (27.5%)  | <i>A. flavus</i> , <i>A. niger</i> and <i>A. ochraceus</i>  |
| Hematological unit (Air)                  | <i>Penicillium</i> sp. (44.6%),<br><i>Aspergillus</i> sp. (28.2%)  | <i>A. flavus</i> , <i>A. niger</i> , <i>A. versicolor</i>   |
| Maternity and hospitals' food units (Air) | <i>Cladosporium</i> sp. (28.4% - 23.2%)<br><i>Penicillium</i> sp. (41.5% - 43.6%)  | Maternity ( <i>A. versicolor</i> , <i>A. fumigatus</i> , <i>A. ochraceus</i> and <i>A. niger</i> )<br>Hospital food units ( <i>A. versicolor</i> , <i>A. niger</i> , <i>A. flavus</i> , <i>A. ochraceus</i> , <i>A. candidus</i> , <i>A. fumigatus</i> , and <i>A. niveus</i> ) |
| Companies' food units (Air)               | <i>Cladosporium</i> sp. (71.2%)<br><i>Penicillium</i> sp. (13.0%)  | <i>A. glaucus</i>   |
| Poultry (Air)                             | <i>Cladosporium</i> sp. (40.5%)<br><i>Alternaria</i> sp. (10.8%)   | <i>A. fumigatus</i> , <i>A. flavus</i> and <i>A. niger</i> .  |
| Sampled surfaces                          | <i>Penicillium</i> sp. was the most frequent isolated genus with an exception to gymnasiums with swimming pools and the hematological unit where the most frequent genus or species detected were <i>Fusarium</i> sp. and <i>A. flavus</i> respectively. | Hematological Unit<br><i>A. versicolor</i> , <i>A. niger</i> , <i>A. fumigatus</i> , <i>A. glaucus</i> , <i>A. nidulans</i>   |

## Conclusions

All the analyzed settings are contaminated by fungi known as mycotoxins producers. Considering the public health risk due to possible air contamination and exposure to mycotoxins by inhalation, preventive measures must be taken.

It was also possible to perform the detection of *Aspergillus* species. Although some *A. flavus* strains do not produce mycotoxins, aflatoxin presence is possible to occur in some of the studied settings. Furthermore, Aflatoxin B1 has been classified as a human carcinogen (hepatocellular carcinoma) by the International Agency for Research on Cancer with a sufficient evidence in humans and a strong support that main mechanism is genotoxicity

Additionally, in some of the studied settings, risk assessment must be performed considering the toxicological interactions between mycotoxins and the sensibility of the exposed population.