

# Fungal Contamination in Lisbon's Primary Schools - Sampling Insights and Analytical Approaches

Renata Cervantes<sup>1,2</sup>, Pedro Pena<sup>1,2</sup>, Bianca Gomes<sup>1,3</sup>, Marta Dias<sup>1,2</sup>, Bruna Riesenberger<sup>1</sup>, Margarida Rodriguez<sup>1</sup>, Liliana Marques<sup>1</sup>, Carla Viegas<sup>1,2</sup>

<sup>1</sup> H&TRC—Health & Technology Research Center, ESTeSL—Escola Superior de Tecnologia e Saúde, Instituto Politécnico de Lisboa, 1990-096 Lisbon, Portugal;

<sup>2</sup> Public Health Research Centre, Comprehensive Health Research Center, NOVA National School of Public Health, CHRC, REAL, CCAL, NOVA University Lisbon, 1099-085 Lisbon, Portugal

<sup>3</sup> CE3C—Center for Ecology, Evolution and Environmental Change, Faculdade de Ciências, Universidade de Lisboa, 1749-016 Lisbon, Portugal



## Introduction

Climate change is posing challenges for Portugal due to intense weather changes, affecting public health and causing pathogens to adapt and spread, increasing the global risk of infectious diseases [1,2].

Azole fungicides are less effective against resistant fungi, raising concerns for children [1,2].

Warm and humid conditions promote the growth of pathogenic fungi and the production of mycotoxins, impacting health by causing gastrointestinal problems, organ damage and chronic diseases. Even after fungi removal, mycotoxins continue to pose risks [3,4,5].

## Objectives

- Identifying fungal species present in indoor environments.
- Assessing spatial distribution and concentration levels within classrooms and other areas.
- Investigating factors influencing fungal proliferation, such as building characteristics and seasonal variations.
- Evaluating the effectiveness of existing cleaning protocols and providing insights into proactive management strategies to protect students' and staff members' health and well-being.


## Methodology

### Sampling Methods

#### Active




#### Passive




#### Particulate Matter



### Analytical Methods



### Metabolites Assessment



## Results and discussion

The expected results are that seasonal variations in fungal load show complex environmental interactions[1,2].

Examining fungal load distribution in DGI8 media at 27°C and 37°C helps assess growth preferences at different temperatures[6].

Methods used to assess azole resistance and mycotoxin provide essential insights into the resilience and potential harm of fungal species under varying environmental conditions[3,4,5].

Addressing fungal exposure risks requires a comprehensive approach for an accurate risk assessment and to target mitigation strategies on educational environments [6].

## Acknowledgements

This project was supported by FCT/MCTES UIDP/05608/2020 (https://doi.org/10.54499/UIDP/05608/2020) and UIDB/05608/2020 (https://doi.org/10.54499/UIDB/05608/2020). This work is also supported by national funds through FCT/MCTES/FSE/UE, 2023.01366.BD; UI/BD/153746/2022 and CE3C unit UIDB/00329/2020 (https://doi.org/10.54499/UIDB/00329/2020); UI/BD/151431/2021 (https://doi.org/10.54499/UIDB/151431/2021); and Instituto Politécnico de Lisboa, national support through IPL/2022/InChildHealth/BI/12M; IPL/IDI&CA2023/FoodAllEU\_ESTeSL; IPL/IDI&CA2023/ASPRisk\_ESTeSL; IPL/IDI&CA2023/ARAFSawmills\_ESTeSL. This project was partly funded by EU Horizon 2021 grant no. 101056883 and co-funding from author's organizations and/or Ministries. Funding from Swiss SERI grant 22.00324, UKRI grant 10040524, and NHMRC grant APP2017786 and APP2008813. Views expressed are of the author(s) and do not necessarily reflect those of EU, Swiss SERI, UKRI, or NHMRC

## Conclusions

- **Standardized protocols** need to be defined and implemented for **effective risk assessment**.
- it is essential to consider **climate changes** and **seasonal influences** into **health policies** to mitigate the risks associated with fungal exposure.

## References

- [1] Seidel et al., 2024 [https://doi.org/10.1016/S2666-5247\(24\)00039-9](https://doi.org/10.1016/S2666-5247(24)00039-9)
- [2] Xiao et al., 2022 <https://doi.org/10.1007/s00253-022-12119-2>
- [3] WHO. 2023b <https://www.who.int/teams/environment-climate-change-and-health/air-quality-and-health/health-impacts>
- [4] Adams et al., 2021 <https://doi.org/10.1111/ina.12865>
- [5] Reham & Gamaleldin et al., 2020 <https://microbiologyjournal.org/prevalence-of-bacteria-in-primary-schools/>
- [6] Viegas et al. 2019 <https://doi.org/10.3390/microorganisms7080234>