

Article

Development and Characterization of Nutritious and Sustainable Canned Fish Meal Prototype for Different Population Segments

Helena Oliveira ^{1,2,*}, Maria Cristóvão ^{1,2,3,†}, Alexandre Nabais ^{1,2,3}, Carolina Camacho ^{1,2}, Maria Sapatinha ^{1,3}, Carla Pires ^{1,2}, Helena M. Lourenço ¹, Maria H. Fernandes ⁴, Maria J. Fernandes ⁴, Maria J. Fraqueza ⁴, Helga C. Augusto ⁵, Cláudia Viegas ⁶, Maria P. Duarte ³, António Marques ^{1,2}, Amparo Gonçalves ^{1,2} and Maria L. Nunes ²

- ¹ IPMA, IP—Portuguese Institute for the Sea and Atmosphere, I.P., Division of Aquaculture, Upgrading and Bioprospection, Av. Doutor Alfredo Magalhães Ramalho 6, 1495-165 Algés, Portugal; a.nabais98@gmail.com (A.N.); carollcamacho@ipma.pt (C.C.); m.sapatinha@gmail.com (M.S.); cpirez@ipma.pt (C.P.); helena@ipma.pt (H.M.L.); amarques@ipma.pt (A.M.); amparo@ipma.pt (A.G.)
 - ² CIIMAR/CIMAR-LA—Interdisciplinary Centre of Marine and Environmental Research, University of Porto, Terminal de Cruzeiros do Porto de Leixões, Av. General Norton de Matos S/N, 4450-208 Matosinhos, Portugal; nunes.leonor@gmail.com
 - ³ METRICs—The Mechanical Engineering and Resource Sustainability Centre, Chemistry Department, NOVA School of Science and Technology, Universidade NOVA de Lisboa, Campus de Caparica, 2829-516 Caparica, Portugal; mpcd@fct.unl.pt
 - ⁴ CIISA—Centre for Interdisciplinary Research in Animal Health, AL4AnimalS—Associate Laboratory for Animal and Science, Faculty of Veterinary Medicine, University of Lisbon, Av. Universidade Técnica, Polo Universitário do Alto da Ajuda, 1300-477 Lisboa, Portugal; helenafernandes@fmv.ulisboa.pt (M.H.F.); mjfernandes@fmv.ulisboa.pt (M.J.F.); mjoaofraqueza@fmv.ulisboa.pt (M.J.F.)
 - ⁵ COFISA—Conservas de Peixe da Figueira, S.A., Terrapleno do Porto de Pesca—Gala, 3090-735 Figueira da Foz, Portugal; helgacaugusto@cofisa.pt
 - ⁶ H&TRC—Health & Technology Research Centre, ESTeSL—School of Health Technology, Polytechnic University of Lisbon, Av. D. João II, Lote 4.69.01, 1990-096 Lisboa, Portugal; claudia.viegas@estesl.ipl.pt
- * Correspondence: helena.oliveira@ipma.pt
- † These authors contributed equally to this work.



Academic Editors: Jaspreet Singh and Francesco Desogus

Received: 7 February 2025

Revised: 4 April 2025

Accepted: 22 May 2025

Published: 30 May 2025

Citation: Oliveira, H.; Cristóvão, M.; Nabais, A.; Camacho, C.; Sapatinha, M.; Pires, C.; Lourenço, H.M.; Fernandes, M.H.; Fernandes, M.J.; Fraqueza, M.J.; et al. Development and Characterization of Nutritious and Sustainable Canned Fish Meal Prototype for Different Population Segments. *Sustainability* **2025**, *17*, 5050. <https://doi.org/10.3390/su17115050>

Copyright: © 2025 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Canning is an excellent solution to provide convenient, affordable, nutritious, and safe seafood with a long shelf life. However, many canned products use tuna, sardines, and Atlantic chub mackerel, species that raise sustainability concerns and whose overuse can put additional pressure on them. Hence, this study aimed to i) develop and assess a nutritious and sustainable canned meal prototype using the Atlantic horse mackerel (*Trachurus trachurus*) (whose EU allowable-catch amounts have increased, particularly, in Iberian waters) and vegetables in light brine in terms of stability, sterility, and physicochemical and sensory properties over a 4-month period at room temperature and ii) evaluate its nutritional contribution for different population groups. After preparation, the meal was stored for one month at $\approx 20^\circ\text{C}$ and $\approx 40^\circ\text{C}$ (to simulate the 4 months). Although the pH was not stable, the meal was considered commercially sterile according to the challenge accelerated tests. Moreover, aging did not significantly affect the meal's physicochemical and sensory properties. This innovative meal prototype can be claimed to be “low-fat”, “reduced in NaCl/Na”, a “source of protein, phosphorus, iron, selenium and vitamin D”, and “high in vitamin B12”. It proved to be both nutritious and appealing for consumption, with potential to be scaled up.

Keywords: Atlantic horse mackerel; eco-innovative product; accelerated aging; stability and sterility; physicochemical properties; nutritional contribution; sensory attributes

1. Introduction

The regular consumption of seafood is strongly recommended by most dietary guidelines as it is a rich source of essential nutrients such as high-quality proteins, long-chain omega-3 polyunsaturated fatty acids (*n*-3 PUFAs; healthy lipids), minerals (e.g., iodine—I, iron—Fe, zinc—Zn, and calcium—Ca), and vitamins (A, D, and B12) for all population groups (children and adults, including pregnant women and seniors) [1]. For instance, the most common recommendation in Europe is an intake of one to two portions of fish (approximately 100–150 g) (depending on the country) per week to ensure the supply of essential nutrients [2,3]. A diet rich in seafood contributes not only to the prevention of many diseases (e.g., cardiovascular, cerebrovascular, and diabetes) but also to the consumer's well-being [1,4]. Additionally, many vegetables, including grain legumes such as beans, are also nutritionally valuable, providing protein with essential amino acids, complex carbohydrates, dietary fiber, vitamins, minerals, and phytochemical compounds with several bioactive properties. They are cholesterol-free and generally low in fat, being considered excellent options to complement healthy and less monotonous meals [5,6].

On the other hand, the demand for ready-to-eat/ready-to-cook (convenient) foods is growing globally, pushing processors to integrate new approaches and culinary concepts in the production of traditional products. Among traditional technologies, canning represents a key strategy in seafood processing [7]. In this process, the seafood flesh (e.g., an edible fish portion cooked and drained) is placed in cans with other ingredients, such as vegetables and salt, and covered with a filling medium, which can be an edible oil (e.g., olive oil or sunflower oil), water, or a sauce. Then, the cans are sealed, washed, and steam-sterilized [7,8]. This process allows the market to distribute convenient products on a global scale, taking advantage of the extended shelf life for distribution, storage, and consumption (without refrigeration) under controlled food safety conditions [7]. It is expected that the canned food market in Europe will grow at a compound annual rate (CAGR) of 3.75% between 2024 and 2029, thus increasing the production volume by approximately 20% during this period [9].

The fish most commonly used in the production of canned products are small pelagics (anchovies, sardines, Atlantic mackerel, and chub mackerel) and the top predator tuna [7,8]. However, the population status and spatial distribution of small pelagic fish are highly susceptible to environmental changes and economic development [10]. Therefore, due to the sustainability of these fishery resources, it is becoming important to diversify by preparing canned products with other less-consumed and abundant species, such as the Atlantic horse mackerel (*Trachurus trachurus*). The good condition of its existing stocks, mainly due to the fact that it is a species that grows rapidly in its early life stages and has an extended breeding season, has led to increases in allowed catch amounts (e.g., proposed EU Total Allowable Catches (TAC) for horse mackerel in Iberian waters (2024): +5%) [11–13]. Since this species presents a sustainable exploitation pattern in the Iberian Peninsula and is well known to have a high nutritional value, several marketing initiatives have been implemented to encourage its consumption and valorization [11,14–16]. Moreover, it is well known that legumes are also appreciated all over the world as sustainable and economical alternatives to meat [5]. Therefore, given that many consumers consider canned fish to be tasty and versatile and often consume it with cooked or canned vegetables, the authors believe that offering a sustainable canned meal that includes both fish and vegetables could be a good option to meet their needs. The attractiveness of a new product can be strengthened if it is affordable, tastefully appealing, and nutritionally balanced (e.g., low/reduced in sodium chloride (NaCl)/sodium (Na), which is associated with high blood pressure and other health problems) [17,18]. This last fact is extremely important

because according to [19], around 11 million deaths globally are associated with a poor diet, 3 million of which are related to high Na intake.

However, no studies were found in the literature addressing the preparation, nutritional characterization, commercial stability and sterility, physicochemical and sensory properties, and performance of canned meals based on fish (e.g., Atlantic horse mackerel) and vegetables during storage. For this reason, the focus of this work was to i) develop and assess a nutritious, sustainable, convenient, and appealing canned meal prototype using Atlantic horse mackerel, a selection of grain legumes, and other vegetables in a light brine in terms of stability, sterility, and physicochemical and sensory properties over a 4-month period of storage at room temperature and ii) evaluate its nutritional contribution for children (7–10 years), healthy adults (≥ 18 years), and pregnant women.

2. Materials and Methods

2.1. Selection of Grain Legumes and Other Vegetables

The vegetables used were selected for their versatility, availability, and nutritional value. Specifically, the chickpea was mainly chosen as an ingredient since it (i) is rich in fiber and protein; (ii) contains several key vitamins and minerals [16], and (iii) has a well-accepted color. The choice of green bean and carrot was made because they are vegetables that are part of most consumers' diets and are also well accepted due to their vibrant colors. Finally, the courgette was chosen for its appearance and to smooth out the filling medium (Table 1).

Table 1. Canned meal recipe ingredients.

Ingredients	Quantity (%)
Steamed horse mackerel fillet	19.7
Sautéed onion and garlic (in olive oil) *	8.7
Blanched chickpea	17.5
Peeled blanched courgette	6.6
Peeled blanched carrot	6.6
Thawed green bean	4.4
Xanthan gum (in powder)	0.04
Brine (14% NaCl)	3.0
Water (to the top)	33.6
Total	100

* Cooked for around 30 min at 100–105 °C.

2.2. Preparation of Raw Material (Fish) and Other Ingredients

Fresh Atlantic horse mackerel (188.8 ± 29.4 g total weight; 25.7 ± 1.1 cm total length; mean $\pm$ standard deviation) caught off the FAO 27 area in Portuguese waters was supplied by a local wholesaler and transported to the laboratory under controlled temperature conditions. Upon arrival, fish were beheaded, gutted, washed, drained, and steamed for 5 min at 100 °C in an oven (Combi-Master CM6, Rational Grossküchen Technik, GmbH, Landsberg am Lech, Germany). The fish were then cooled at room temperature, filleted by hand, and distributed among 32 glass jars (with a volume of approximately 290 cm³). The list of ingredients and quantities used to prepare a balanced canned meal is shown in Table 1.

The chickpeas, courgettes, and carrots were bought fresh at a local supermarket and blanched in the laboratory to avoid losses of both color and flavor as well as texture degradation that are usually caused by enzymatic action [20]. More specifically, the chickpeas were water-soaked overnight in an airy place and blanched in the next day for 10 min in boiling water (2:5 *m/v*; with 0.7% NaCl). After draining the blanching water, the chick-

peas were cooled with cold water and ice to stop the blanching process, drained, and frozen at $-20\text{ }^{\circ}\text{C}$. The carrots and courgettes were cleaned, peeled, and cut into slices ($\approx 0.4\text{ cm}$ of thickness) and then into semicircles and quarters, respectively. Afterward, these pieces were blanched (1 min) and cooled with boiling and iced water, respectively. The green beans were purchased frozen also at a local supermarket (already blanched). Additionally, xanthan gum (xanthan gum clear, Sosa, Barcelona, Spain) was added as a thickener; the brine concentration (NaCl, 14%) was selected based on the NaCl contents indicated by companies aiming to supply healthy products, i.e., lower than $1\text{ g}/100\text{ g}$, and the sensory results obtained in a preliminary test carried out with canned Atlantic horse mackerel fillets in olive oil, which revealed that a NaCl concentration of around $0.6\text{ g}/100\text{ g}$ ($\text{Na} = 230 \pm 22\text{ mg}/100\text{ g}$) is sufficient to impart a slight-to-moderate salty taste to the product. Finally, water was selected as filling medium since it was found (in a previous assay) that olive oil used for this purpose can make vegetables very oily and unappealing to consumers.

2.3. Canning and Accelerated Aging

The 32 glass jars were sealed and sterilized (Uniclave 88, AJ Costa, Agualva-Cacém, Portugal) for 75 min at $115 \pm 1\text{ }^{\circ}\text{C}$ (well established and common practice of industrial units for this type of product based on previous F0 calculations [21]) at the Faculty of Veterinary Medicine of the University of Lisbon. After cooling, all glass jars were stored for 30 days (usual procedure to stabilize canned products) [21] in two different conditions: (i) 16 were kept at room temperature ($\pm 20\text{ }^{\circ}\text{C}$) in a dry place not exposed to direct sunlight and (ii) 16 were maintained at $38.9 \pm 0.8\text{ }^{\circ}\text{C}$ (to simulate a 4-month aging period at $20\text{ }^{\circ}\text{C}$). The calculation of accelerated aging was based on the Arrhenius equation, which indicates that an increase of $10\text{ }^{\circ}\text{C}$ at a given temperature doubles the rate of chemical reactions and/or microbiological growth [22].

2.4. Analytical Methods

The proximate composition and histamine, NaCl, Na, and K contents were assessed in the raw material ($n = 3$ homogenized fish; Grindomix GM200, Retsch, Haan, Germany; 5000 rpm). Furthermore, the canned samples stored at the two different conditions were evaluated in terms of both stability and sterility ($n = 9$ jars/condition) as well as physico-chemical ($n = 3$ jars/condition) (with the exception of histamine, which was only assessed in the raw material) and sensory analyses ($n = 3$ jars/condition) through the methods described below (Sections 2.4.1–2.4.4). For chemical analyses and instrumental color, the filling medium of each jar was drained and the mix of fish and vegetables was homogenized (Grindomix GM200). The analytical quality control of chemical analyses is presented in Supplementary Table S1. All chemical results are expressed in wet weight.

2.4.1. Stability and Sterility

The stability test was carried out in accordance with the Portuguese standard NP 4404-1 [23], which consists in the evaluation of packages stored at three different temperatures ($n = 3$ /condition for each temperature), $20\text{ }^{\circ}\text{C}$ (21 days), $30\text{ }^{\circ}\text{C}$ (21 days), and $55\text{ }^{\circ}\text{C}$ (seven days), in relation to (i) external examination of the package (glass jar and lid) during the storage time; (ii) internal examination of the glass jar content after opening, i.e., evaluation of the appearance (color and texture) and odor of the meal; and (iii) pH measurement of content (pH meter: HI99163; pH electrode: FC230B, HANNA Instruments, Póvoa de Varzim, Portugal). Thin smears of each glass jar were also prepared for fresh examination and stained using the Gram method. The microscopic observation of the microorganisms (average number per field) was focused in 20 optical fields [23].

On the other hand, the canned samples' sterility was determined according to the Portuguese standard NP 2309-2 [24] for low-acid food. This test involved the detection of aerobic and anaerobic mesophiles and thermophiles, with the media being incubated at 30 and 55 °C (i.e., at the same temperatures that the packages were stored for 21 and 7 days, respectively).

2.4.2. Chemical Analyses

Proximate Composition and Energy Value

The contents of moisture, ash, and free fat were determined according to the Association of Official Analytical Chemists methods [25]. Briefly, moisture was assessed by oven (ULE 500, Memmert, Schwabach, Germany) drying of each sample overnight at 105 ± 2 °C while ash was obtained by incineration of previously dried sample in a muffle furnace (M104, Heraeus, Hanau, Germany) for 16 h at 500 ± 25 °C. Free fat was quantified using a Soxhlet extraction method (in a Soxhlet apparatus, Behr Labor-Technik, Dusseldorf, Germany), with diethyl ether as a solvent (at approximately 40 °C; 7 h), by weighing the fat residue after drying (105 ± 2 °C) in an air oven. Total nitrogen was determined following the Dumas method [26] in an automatic nitrogen analyzer (FP-528, LECO Corp., St. Joseph, MI USA) calibrated with EDTA. Nitrogen was released by combustion at 850 °C and detected by thermal conductivity. Crude protein was calculated from total nitrogen through the conversion factor of 6.25 [27]. Total carbohydrates were calculated by difference and the energy value was estimated using Food and Agriculture Organization factors [27].

NaCl and Macroelements (Na and K)

The measurement of salt (i.e., chlorine, Cl^- as NaCl) content was performed by volumetric titration of sample with silver nitrate (AgNO_3) according to Mohr's method [28]. Additionally, the Na and potassium (K) contents were quantified by flame atomic absorption spectrophotometry (Spectr AA 55 B spectrophotometer, Agilent Technologies, Santa Clara, CA, USA) with a background deuterium correction, following the method reported by [29]. The concentrations were calculated through linear calibration obtained from absorbance measurements of, at least, five different concentrations of standard solutions (NaNO_3 and KNO_3 , both dissolved in 0.5 M HNO_3).

Histamine and Total Volatile Basic Nitrogen (TVB-N)

The determination of histamine was carried out on samples previously homogenized (POLYTRON® PT 3100 D, Kinematica AG, Malzers, Switzerland) with 10% (m/V) trichloroacetic acid (1:2, m/V) and filtered through a 0.2 µm porosity filter. The amine was separated by high-performance liquid chromatography (HPLC, Agilent 1100 Series, Santa Clara, CA, USA) on a reverse-phase column, LiChrospher® 100 RP-18 (5 µm) LiChroCART® 250-4, according to an elution gradient (mobile phase, consisting of two eluents: 0.1 M sodium acetate buffer (pH 4.5, adjusted with glacial acetic acid, containing 1-octanesulfonic acid) and 0.2 M sodium acetate buffer (pH 4.5, adjusted with glacial acetic acid, containing acetonitrile and 1-octanesulfonic acid)) at a flow rate of 1.0 mL/min and post column derivatization with ortho-phthalaldehyde reagent (OPA), following the method described by [30]. The detection of the compound was performed by fluorescence in an Agilent 1100 Series fluorimeter ($\lambda_{\text{Excitation}} = 340$ nm and $\lambda_{\text{Emission}} = 455$ nm). Histamine was identified and quantified by comparison with a calibration curve (hexanediamine was used as internal standard) in the working range of 1.25–10 µg/mL using the Software Agilent Chem Station for LC (version A.10.02 [1757]).

TVB-N content was determined in trichloroacetic acid (5%) extracts of homogenized samples (POLYTRON® PT 3100 D) by the microdiffusion Conway method [31] using a drying oven (INCO 2/108, Memmert, Schwabach, Germany).

2.4.3. Instrumental Color

The instrumental color was analyzed in all samples using a colorimeter (CR-400, KONICA MINOLTA, Tokyo, Japan), at standard illuminant D65 and a 2-degree observer, previously calibrated with a white standard tile ($L^* = 93.47$; $a^* = -0.16$; $b^* = 3.32$). The L^* , a^* and b^* coordinates from CIELAB system were recorded. In this system, L^* values describe lightness on a scale of 0 (black) to 100 (white); the a^* values denote the intensity from green (negative) to red (positive); the b^* values range from blue (negative) to yellow (positive). The chroma (saturation; $C^*_{a,b}$) was calculated according to the following equation [32]:

$$C^*_{a,b} = \sqrt{a^{*2} + b^{*2}}$$

2.4.4. Sensory Evaluation

The sensory assessment was performed after sterility confirmation of samples in the sensory laboratory of IPMA equipped with testing room according to [33]. A quantitative descriptive method was applied using seven trained panelists (both sexes) and a 5-point scale (0—absent; 1—slight; 2—moderate; 3—intense; 4—extreme) to rate the intensity of the selected attributes [34]: canned food odor, other odors, color of ingredients (fish, chickpea, green bean, courgette, and carrot) and filling medium, salty taste, other flavors, fish firmness, fish succulence, meal oiliness, filling medium consistency, ingredient cooking degree (fish, chickpea, green bean, courgette, and carrot), and mouth feeling. The jars were opened 15 min before testing and about 60 g of each sample (stored at 20 °C and 40 °C) was presented to each panelist in white coded plates. The panelists rinsed their mouths with water between portion assessments to reduce the residual flavor effects.

2.5. Estimated Nutritional Composition of Other Nutrients

Other nutrient levels present in the meal recipe were estimated with @Nutrisoft—@Nutrition Information Software (version 1, Registration n°. 3208/2012) [35]. This software allowed the estimation of some macro- and micronutrients (i.e., phosphorus (P), Ca, magnesium (Mg), Fe, Zn, I, selenium (Se), and vitamins B12 and D) based on the amounts of ingredients used and their nutritional values. The calculations were carried out using the food composition table published by [16].

2.6. Nutritional Contribution

The meal nutritional contribution (i.e., 190 g, which corresponded approximately to the drained product) in terms of proteins, Na, K, P, Ca, Mg, Fe, Zn, I, Se, and vitamins B12 and D was calculated considering the dietary reference values (DRVs) recommended by [17,36] for different population groups (children aged 7–10 years old and adults, including pregnant women). The calculations (%) were made according to the following equation:

$$\text{Nutritional Contribution(\%)} = \frac{C \times M}{\text{DRV}} \times 100$$

Here,

C—the mean concentration of the nutrient (mg/kg) (obtained as indicated in previous sections). In the case of protein, Na, and K, only the contents observed after 1 month at room temperature were used since the impact of aging on these parameters was nil (as can be seen in Section 3).

M—the weight of the defined meal (kg).

DRV—the PRI, Population Reference Intake (mg/day), in the case of protein, Ca, Fe, and Zn, or the AI, Adequate Intake (mg/day), in the case of the other specific nutrients.

2.7. Statistical Analysis

Statistical analysis was performed using the STATISTICA© software version 12 (Stat-Soft Inc., Tulsa, OK, USA). All data were checked for normality of distribution and homogeneity of variances using Shapiro–Wilk and Levene’s tests, respectively. The influence of storage at different conditions on proximate composition, energy value, NaCl and macroelement (Na and K) contents, TVB-N level, instrumental color, and sensory attributes of the canned meal was tested by Student’s t-test. Statistical significance was considered at $p < 0.05$ for all analyses [37].

3. Results and Discussion

3.1. Stability and Sterility

The external visual examination of the packages (glass jars and lids) analyzed did not reveal any physical damage or leakage of the filling medium. In both tested conditions, the appearance (including color and texture) and odor of the samples were considered characteristic of a canned product. However, it was observed that some vegetables (particularly the courgette) showed signs of overcooking. Furthermore, some pH differences of more than 0.5 were detected between the samples maintained at ≈ 20 °C and those kept at 30 and 55 °C in both conditions (Table 2). These differences can be explained by the pH variation in the courgette based on previous experiments without this vegetable. This change could be ascribed to accelerated chemical reactions triggered by heating, such as possible Maillard reactions or other heat-induced transformations like decarboxylation [16,38]. Finally, the fresh smear and Gram coloration tests, in 20 fields, also had negative results, meaning that there was no microorganism growth.

Table 2. pH values found in the stability tests performed with canned meal prepared with Atlantic horse mackerel and vegetables after 1 and 4 months of storage.

Storage Time (Months)	pH
1	6.4 ± 0.1^a , 5.9 ± 0.1 , 5.7 ± 0.5 (at ≈ 20 , 30 and 55 °C, respectively)
4 *	6.4 ± 0.1^a , 5.7 ± 0.1 , 5.5 ± 0.4 (at ≈ 20 , 30 and 55 °C, respectively)

Results are given as mean values $\pm$ standard deviations ($n = 3$). * Simulated 4-month aging period at 20 °C (1 month at ≈ 40 °C). For the glass jars maintained at ≈ 20 °C, the same letter (a) indicates absence of significant differences ($p < 0.05$) between storage times.

Although the pH was not considered stable, the results of the sterilization assessment showed that no microbial growth (mesophilic and thermophilic aerobes and anaerobes) was observed under the specified conditions of the tests, indicating that the developed meal could be considered sterile for at least 4 months at room temperature.

3.2. Proximate Composition, Energy Value, NaCl, and Macroelements (Na and K)

The proximate composition, energy value, and NaCl and macroelement (Na and K) contents of Atlantic horse mackerel and the canned meal are shown in Table 3. The values obtained for raw fish were in agreement with those reported by other studies with the same (caught in the spring season) or other pelagic species [39–44].

Additionally, there were no significant changes in the moisture content of the Atlantic horse mackerel canned meal after 1 and 4 months of storage at room temperature (≈ 79 g/100 g). Likewise, the protein (≈ 10 g/100 g) and fat (≈ 2 g/100 g) contents did not change either ($p > 0.05$). The values obtained also indicate that the canned meal developed in the present work can be nutritionally claimed as a “source of protein” and “low-fat” since more than 12% of the energy (i.e., approximately 45%) was supplied by proteins and it had an amount < 3 g fat/100 g, respectively [45]. The mean contents of ash and carbohydrates, as

well as the energy value, were also not significantly affected by the storage time (1 g/100 g, close to 8 g/100 g, and 93 kcal/100 g, respectively) ($p > 0.05$).

Table 3. Proximate composition, energy value, and NaCl and macroelement (Na and K) contents of the raw material (fish) and canned meal prepared with Atlantic horse mackerel and vegetables after 1 and 4 months of storage.

g/100 g	Atlantic Horse Mackerel	Canned Meal Storage Time (Months)	
		1	4 *
Moisture	76.7 ± 0.6	78.9 ± 0.1 ^a	79.1 ± 0.4 ^a
Protein	20.9 ± 0.8	10.2 ± 0.2 ^a	10.3 ± 0.6 ^a
Fat	1.9 ± 0.4	1.9 ± 0.2 ^a	1.8 ± 0.3 ^a
Ash	1.4 ± 0.1	1.0 ± 0.0 ^a	1.0 ± 0.0 ^a
Carbohydrates ¹	---	8.0 ± 0.3 ^a	7.9 ± 0.7 ^a
Energy value (kcal/100 g)	---	93.5 ± 0.9 ^a	92.4 ± 1.8 ^a
NaCl	0.2 ± 0.0	0.5 ± 0.0 ^a	0.5 ± 0.0 ^a
Na (mg/100 g)	54.9 ± 0.6	166.9 ± 12.0 ^a	162.6 ± 2.8 ^a
K (mg/100 g)	487.5 ± 11.7	155.3 ± 14.2 ^a	145.2 ± 3.8 ^a

Results are given as mean values ± standard deviations (n = 3). For each parameter, the same letter (a) indicates absence of significant differences ($p < 0.05$) between storage times. * Simulated 4-month aging period at 20 °C (1 month at ≈ 40 °C). ¹ Calculated by difference.

The NaCl and Na contents found in the canned meal (0.5 g/100 g and approximately 165 mg/100 g, respectively) were also not significantly influenced by the studied storage period ($p > 0.05$), being lower compared to those found in various canned fish products sold in supermarkets (NaCl: ≥ 1 g/100 g) [46]. The NaCl content was within the large variation (between 32 to 6000 mg/100 g) reported by [47] for canned fish products. Moreover, the Na value was below the global Na benchmark defined (360 mg/100 g) for canned fish (with the exception of canned anchovies) [19]. So, considering the values observed, the prepared canned meal can be considered a healthy product (with reduced NaCl/Na; [45]) whose consumption can contribute to achieve the objectives defined by the WHO (in terms of NaCl reduction) [48] and many seafood producers. Finally, the K content resulted not only from the amount naturally present in fish (e.g., Ref. [49]) but also from the quantities existing in the other ingredients used (mainly in chickpeas, around 270 mg/100 g after cooking) [16]. Storage for up to 4 months had no impact on K either (close to 150 mg/100 g) ($p > 0.05$). Hence, it appears that this new product can be stored at room temperature for up to 4 months without the alteration of its proximate composition and NaCl, Na, and K contents.

3.3. Estimated Nutritional Composition

The most abundant macroelement was Na, followed by P, K (with identical levels), Ca, and finally Mg. Within the microelements, Fe had the highest content followed by Zn, I, and lastly Se (Tables 3 and 4). The results shown in Table 4 support the possible nutritional claim “source of phosphorus, iron and selenium” since the meal recipe had amounts $> 15\%$ of the Nutrient Reference Values, NRVs (i.e., 700 mg/100 g, 14 mg/100 g, and 55 µg/100 g, respectively) [45,50]. Hence, the consumption of this meal based on fish and vegetables may be relevant, for instance, to avoid health constraints such as (i) muscle weakness and bone pain (due to P deficiency), (ii) anaemia (caused by P and Fe deficiencies), and (iii) organ and tissue degeneration (due to Se deficiency), which can lead to the appearance of the Keshan and Kashin–Beck diseases [36,51].

Table 4. Contents of some macro- and microelements and vitamins in the meal recipe with Atlantic horse mackerel and vegetables.

Content in the Recipe (mg/100 g of Drained Product) ¹	
Macroelements	
P	155.7
Ca	72.2
Mg	42.8
Microelements	
Fe	2.3
Zn	1.2
I	0.0142
Se	0.0100
Vitamins	
B12	0.0017
D	0.0012

¹ Nutrients contents were calculated with @Nutrisoft—@Nutrition Information Software version 1 (Registration n°. 3208/2012).

Additionally, the value of vitamin B12 was higher than 0.75 µg/100 g (>30% of the NRV—i.e., 2.5 µg/100 g), allowing the probable nutritional claim “high in vitamin B12” (Table 4) [45,50]. Thus, the intake of the prepared meal can provide a good amount of vitamin B12, which should help, for example, keep the body’s blood and nerve cells healthy in addition to reducing tiredness and fatigue [52,53]. In terms of vitamin D, the value found was higher than 0.75 µg/100 g (15% of the NRV—i.e., 5 µg/100 g), permitting the possible nutritional claim “source of vitamin D” (Table 4) [45,50]. This result shows that the consumption of this meal based on fish and vegetables may contribute to other health benefits such as the maintenance of normal bone, teeth, and muscle function, as well as the normal absorption/utilization of calcium and phosphorus and the proper functioning of the immune system [54,55]. Moreover, the vitamin D amount was well below the tolerable upper intake level/day (UL/day) set for children aged 7–10 years (50 µg/day) and adults/pregnant women (100 µg/day) [56].

It is well known that cooking techniques that require heat can affect the nutritional composition of food [57]. However, and according to the literature, the retention factors of the food constituents reported in Table 4 by cooking (boiling) fish and fish-based dishes (low fat fish; <5%) are high (i.e., between 80 and 100%) [16,57–59]. Moreover, vitamins D and B12 are heat stable and fat-soluble vitamins, meaning that both are not easily leached into cooking water [60,61]. So, it is expected that most of the estimated values presented in Table 4 are close to those that can be obtained after heat treatment (sterilization).

3.4. Nutritional Contribution

3.4.1. Proteins and Macro- and Microelements

The nutritional contribution of the meal based on fish and vegetables in terms of proteins and macro- and microelements is presented in Table 5. The consumption of 190 g corresponds to a greater protein contribution for children (74% of the PRI) and a lower one for pregnant women in the last trimester (22% of the PRI). In all cases, such contributions are valuable for tissue maintenance [36]. The nutritional contribution is at least 50% of the daily AI of P and Se for children. The same can be also observed in the case of P for adults. The 190 g meal also contributes with at least a quarter ($\geq 25\%$) of the DRVs (AIs or PRIs) in the following cases: Mg for children and women (adults, including pregnant women); I for children; Se for adults (including pregnant women); Fe for all population groups; and Zn for children and women (adults, including pregnant women). Assuming that the population

in Portugal consumes up to five meals a day [62], this percentage ($\geq 25\%$) can be a good contribution to reach the recommended amounts if the consumer eats other nutritious foods throughout the day. The contributions of Mg for male adults; I for adults, including pregnant women; Zn for male adults; and Na, K, and Ca for all population segments were less pronounced, ranging from 8% to 24% of the daily recommended amounts. Specifically, the nutritional contribution of Na corresponded to 16–19% for both groups, values far below the recommended limit proposed by [17].

Table 5. Nutritional contribution of the meal prepared with Atlantic horse mackerel and vegetables in terms of proteins and macro- and microelements for different population segments, considering a dose of 190 g.

Target Nutrient	Target Population	DRV: AI or PRI (mg/day) ¹	Nutritional Contribution (%)
Protein	Children (7–10 y)	26,261 ³	73.6 $\pm$ 1.1
	Adults (≥ 18 y)	52,539 ³	36.8 $\pm$ 0.6
	Pregnant women (3rd trimester)	86,515 ^{3,4}	22.3 $\pm$ 0.3
Macroelements			
Na	Children (7–10 y)	1700 ^{3,5}	18.7 $\pm$ 1.3
	Adults (≥ 18 y) ²	2000 ^{3,5}	15.9 $\pm$ 1.1
K	Children (7–10 y)	1800 ³	16.4 $\pm$ 1.5
	Adults (≥ 18 y) ²	3500 ³	8.4 $\pm$ 0.4
P	Children (7–10 y)	440 ³	67.2
	Adults (≥ 18 y) ²	550 ³	53.8
Ca	Children (7–10 y)	800 ³	17.1
	Adults (≥ 18 y) ²	950/1000 ³	13.7/14.4
Mg	Children (7–10 y)	230–250 ³ (W) ⁶ / 230–300 ³ (M) ⁶	27.1–35.4% (M) ⁶ / 32.5–35.4% (W) ⁶
	Adults (≥ 18 y) ²	300 ³ (W) ⁶ / 350 ³ (M) ⁶	23.2 (M) ⁶ / 27.1 (W) ⁶
Microelements			
I	Children (7–10 y)	0.09 ³	30.0
	Adults (≥ 18 y)	0.15 ³	18.0
	Pregnant women	0.2 ³	13.5
Se	Children (7–10 y)	0.035 ³	53.7
	Adults (≥ 18 y) ²	0.07 ³	26.9
Fe	Children (7–10 y)	11 ³	39.7
	Adults (≥ 18 y)	11 ³ (M, postmenopausal W) ⁶ / 16 ³ (premenopausal W) ⁶	27.3 (premenopausal W) ⁶ / 39.7 (M, postmenopausal W) ⁶
	Pregnant women	16 ⁷	27.3
Zn	Children (7–10 y)	7.4 ^{3,8}	30.8
	Adults (≥ 18 y)	7.5 ^{3,9} (W) ⁶ / 9.4 ^{3,9} (M) ⁶	24.3 (M) ⁶ / 30.4 (W) ⁶
	Pregnant women	9.1 ^{3,10}	25.1

Values of protein, Na, and K are means $\pm$ standard deviations ($n = 3$); other values were calculated from the contents obtained using the @Nutrisoft program; ¹ DRVs (Dietary Reference Values): AIs (Adequate Intakes) are presented in ordinary type and PRIs (Population Reference Intakes) in **bold type**; ² including pregnant women; ³ Ref. [36]; ⁴ +28,000 mg/day in addition to the PRI for protein of non-pregnant, non-lactating women. ⁵ Ref. [17]; ⁶ W—Women, M—Men. ⁷ The PRI covers the requirement of approximately 95% of premenopausal women. ⁸ The fractional absorption of Zn considered in setting PRIs for children was based on data from mixed diets expected to contain variable quantities of phytate; therefore, no adjustment for phytate intake was made; ⁹ considering a phytate intake level of 300 mg/day; ¹⁰ +1.6 mg/day—in addition to the PRIs for non-pregnant, non-lactating women.

3.4.2. Vitamins

The nutritional contributions of vitamins are shown in Table 6. The amounts of vitamin B12 provided by the meal were high for all segments. In other words, the meal (190 g) almost fulfils or fulfils the daily AIs of vitamin B12 for children (>100%) and adults including pregnant women (72–81%). Hence, this product can be a good option for vitamin B12 intake, especially for individuals that have a special need. The nutritional contribution of vitamin D for daily AIs was 15% for all population segments analyzed.

Table 6. Nutritional contribution of the meal prepared with Atlantic horse mackerel and vegetables in terms of vitamins for different segments of the population, considering a dose of 190 g.

Target Nutrient	Target Population	DRV: AI (mg/day) ¹	Nutritional Contribution (%)
B12	Children (7–10 y)	0.0025 ³	129.2
	Adults (≥18 y)	0.004 ³	80.8
	Pregnant women	0.0045 ³	71.8
D	Children (7–10 y)	0.015 ^{3,4}	15.2
	Adults (≥18 y) ²		

Values were calculated from the contents obtained with the @Nutrisoft program. ¹ DRVs (Dietary Reference Values): AIs (Adequate Intakes); ² including pregnant women; ³ Ref. [36]; ⁴ under conditions of assumed minimal cutaneous vitamin D synthesis. In the presence of endogenous cutaneous vitamin D synthesis, the requirement for dietary vitamin D is lower or may be even zero.

3.5. Histamine and TVB-N

Since some publications observed histamine poisoning from other fish species not considered in the EU legislation [63], the control of this parameter was performed in raw Atlantic horse mackerel because its ingestion may be responsible for cases of food intolerance, which may include allergies-like symptoms [64]. Additionally, histamine was only determined in the fish because this type of food has been identified as the predominant causative agent (over 90% of cases) of intoxication by this amine [63,64]. However, the histamine level found was below the detection limit (0.6 mg/kg), which means that the analyzed samples were in compliance with the current EU regulations [65] and were kept following the recommendations, i.e., adequate chilling conditions and hygienic measures [66]. Therefore, these levels (<10 mg/kg,) also indicated the utilization of good-quality fish [67,68]. For this reason, and due to the very low levels of histamine found in canned fish produced with species associated with histamine risk (probably related to the improving quality of the canning process over the years due to the widespread application of HACCP principles, i.e., from the fish caught in the vessel to the processed product) [63], this analysis was not carried out for the canned meal.

On the other hand, the TVB-N levels obtained in the canned meal prepared with Atlantic horse mackerel and vegetables were around 20 mg/100 g after 1 and 4 months of storage. Such contents were not significantly affected by the aging period ($p > 0.05$), being lower than (i) the satisfactory and acceptable range reported for this type of product (40–45 mg/100 g muscle; [39]) prepared from raw material of good quality (i.e., with contents ranging between 25–30 mg/100 g) and also (ii) the limits (25–35 mg N/100 g) set for the fish species or families mentioned in Commission Regulation (EC) No. 1022/2008 [69]. Thus, fish degradation could not be considered in all samples evaluated. Hence, it seems that the canned meal can be stored for up to 4 months while keeping its good quality.

3.6. Instrumental Color and pH

It is well known that chlorophylls and carotenoids, i.e., the pigments responsible for the characteristic colors (green and red/orange/yellow, respectively) of vegetables (e.g., green bean and carrot) are highly susceptible to degradation during storage, resulting

in color changes, which can be perceived by the consumer as a loss of quality [38,70]. However, in the present work, the color of the canned meal remained constant with storage ($p > 0.05$; Table 7). The L^* mean value ranged from 57 to 58 between the storage periods analyzed, being in a medium range of luminosity. The a^* , b^* , and $C^*_{a,b}$ mean values were close to 6.0, 27.0, and 27.0, respectively, at both times. The geometry of the CIELAB color model and a^* and b^* results showed that the color of the meal was in the orange-to-yellow range [71]. The absence of significant differences suggests that the time tested does not seem to cause any color degradation (e.g., associated to degradation of the pigments present in vegetables). So, it is not expected that there will be changes in the sensory perception and consumer acceptance (in terms of color) of the developed canned meal over a 4-month storage period at room temperature.

Table 7. Color parameters of the canned meal prepared with Atlantic horse mackerel and vegetables after 1 and 4 months of storage.

Color Parameters	Canned Meal Storage Time (Months)	
	1	4 *
L^*	57.2 ± 1.3^a	58.0 ± 0.5^a
a^*	6.0 ± 0.6^a	5.6 ± 0.1^a
b^*	26.7 ± 0.8^a	26.5 ± 1.3^a
Chroma ($C^*_{a,b}$)	27.4 ± 0.8^a	27.1 ± 1.2^a

Results are given as mean values $\pm$ standard deviations ($n = 3$). For each parameter, the same letter (a) indicates absence of significant differences ($p < 0.05$) between storage times; * simulated 4-month aging period at 20 °C (1 month at ≈ 40 °C).

On the other hand, accelerated aging (simulated 4 months at 20 °C) also did not promote significant changes in the pH of the meal (close to 6 at both times; $p > 0.05$), revealing a good maintenance of the initial quality (Table 2).

3.7. Sensory Evaluation

The results of the sensory properties (i.e., odor, color, taste, and texture) of the drained canned meal obtained after 1 and 4 months of storage can be found in Figure 1A,B. The meal was always scored by the panel as having a moderate canned food odor. The panelists also detected other odors (mainly sautéed or garlic), but with a slight intensity. Regarding color, the fish, chickpea, and courgette were scored with a moderate intensity while the green bean and carrot were described as having an intense color after both times. The filling medium was scored with a moderate-to-intense color (reported as yellow by most of the panelists) and a slightly thick consistency in the two periods. Furthermore, the prepared product was rated as having a slight-to-moderate salty taste regardless of the storage time evaluated. This result indicates that the NaCl content used was adequate to prepare a tasty and healthy meal with reduced NaCl/Na (Table 3). Other flavors (e.g., bitter and sweet) were considered imperceptible by the panel.

The fish fillet was classified as moderately firm and succulent after 1 month and intensely firm and slightly succulent after 4 months, which indicated a slight tendency of the product to become firmer and less succulent with storage. In both times, the oiliness of the meal was rated as moderate. The fish cooking degree was scored from moderate (1 month) to intense (4 months). The cooking degree of the chickpea was identified as moderate, that of the green bean and carrot as intense (i.e., slightly overcooked), and that of the courgette as extreme (i.e., overcooked) in both periods. Finally, the panel did not detect negative sensations such as astringency (at both times). Therefore, and according to Figure 1A,B, the sensory properties analyzed were not significantly influenced by storage

($p > 0.05$). These results indicate that the developed canned meal can be stored for up to 4 months at room temperature without sensory changes, although they also point out that the pre-cooking time of the fish (before canning) can be slightly reduced to obtain a product with a more appreciable texture. Additionally, it can also be noted that the recipe could be improved by replacing the courgette with another nutritionally rich ingredient since it was classified as overcooked and disintegrated.




Figure 1. Sensory profile of the canned meal prepared with Atlantic horse mackerel and vegetables after 1 and 4 * months of storage (* corresponds to a simulated 4-month aging period at 20 °C; 1 month at ≈ 40 °C) in terms of (A) odor, color, and taste and (B) textural properties. Results correspond to mean values ($0.4 \leq SD \leq 1.5$) ($n = 3$). Intensity scale: 0 (absent); 1 (slight); 2 (moderate); 3 (intense); 4 (extreme). For each parameter, no significant differences ($p < 0.05$) were found between storage times.

4. Conclusions

The findings allow us to conclude that the chosen ingredients (horse mackerel, chickpea, carrot, and green bean) are very suitable for preparing fish-based canned meals. As for the courgette, it ended up not being a good option because it did not provide a smooth consistency to the filling medium and contributed to a slight change in the meal's stability

(based on pH values), although it did not compromise sterility. Hence, further development is needed, especially in exploring vegetable alternatives, to ensure long-term stability.

Based on chemical (proximate composition, energy value, and NaCl, Na, and K contents and TVB-N values), physical (instrumental color and pH), and sensory properties (e.g., canned food odor, salty taste, fish firmness, and fish succulence), this prototype is stable for at least 4 months under normal storage conditions. Furthermore, it could possibly be labelled as “low-fat”, “reduced in NaCl/Na”, “source of protein, phosphorus, iron, selenium and vitamin D”, and “high in vitamin B12”. Moreover, its consumption (190 g; drained product) will contribute to meeting the daily needs of several nutrients (8- > 100%) essential for consumers’ health at all ages.

The present results provide valuable information for scaling up an innovative, nutritious, sustainable, and convenient prototype based on the studied formulation (without the courgette) but with a more appealing filling medium, for instance spiced tomato sauce, so that the consumer considers its consumption (as happens with canned fish in tomato sauces) and not its disposal. Therefore, the market application prospects are promising, considering the growing demand for this type of product. Potential consumers include conscious individuals concerned about health and sustainability issues and busy professionals and their children.

Supplementary Materials: The following supporting information can be downloaded at: <https://www.mdpi.com/article/10.3390/su17115050/su17115050/s1>, Table S1: Analytical quality control of chemical analyses (proximate composition, histamine, TVB-N, and macroelement contents).

Author Contributions: Conceptualization, H.O., A.M, A.G. and M.L.N.; software, C.V.; validation, H.O., M.C., A.N., C.C., M.S., C.P., H.M.L., M.H.F., M.J.F. (Maria J. Fernandes), M.J.F. (Maria J. Fraqueza), C.V., M.P.D., A.G. and M.L.N.; formal analysis, H.O., M.C. and C.C.; investigation, H.O., M.C., A.N., C.C., M.S., C.P., H.M.L., M.H.F., M.J.F. (Maria J. Fernandes), A.G. and M.L.N.; resources, C.V.; data curation, H.O., M.C. and A.N.; writing—original draft preparation, H.O. and M.C.; writing—review and editing, H.O., M.C., A.N., C.C., M.S., C.P., H.M.L., M.H.F., M.J.F. (Maria J. Fernandes), M.J.F. (Maria J. Fraqueza), H.C.A., C.V., M.P.D., A.M., A.G. and M.L.N.; visualization, H.O.; supervision, H.O., M.P.D., A.G. and M.L.N.; project administration, H.O., H.C.A., A.M. and M.L.N.; funding acquisition, H.O., H.C.A., A.M. and M.L.N. All authors have read and agreed to the published version of the manuscript.

Funding: This work was developed within the scope of “BLUE BIOECONOMY INNOVATION PACT” (Project N°. C644915664-00000026), funded by NextGenerationEU, under the incentive line “Agendas for Business Innovation” of the Recovery and Resilience Plan (PRR). It was also supported by the Portuguese Foundation for Science and Technology, I.P. (FCT) through the following strategic projects: UIDB/04077/2020, UIDP/04077/2020, UIDB/04423/2020, and UIDP/04423/2020.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: The original contributions presented in the study are included in the article/Supplementary Material. Further inquiries can be directed to the corresponding author.

Acknowledgments: M.S. is thankful for the doctoral scholarship (2023.02017.BDANA) provided by the Foundation for Science and Technology, I.P. The authors also wish to express their gratitude for the collaboration of the industrial unit COFISA—Conservas de Peixe da Figueira, S.A. Finally, the authors would like to thank the employees of IPMA, I.P. who constituted the sensory panel and Margarida Muro (from the bromatology laboratory) for her continuous and invaluable assistance whenever needed.

Conflicts of Interest: The authors declare no conflicts of interest.

References

- Food and Agriculture Organization of the United Nations (FAO). *The State of World Fisheries and Aquaculture 2024—Blue Transformation in Action*; FAO: Rome, Italy, 2024; 264p. [CrossRef]
- EFSA NDA Panel (EFSA Panel on Dietetic Products, Nutrition and Allergies). Scientific Opinion on health benefits of seafood (fish and shellfish) consumption in relation to health risks associated with exposure to methylmercury. *EFSA J.* **2014**, *12*, 3761. [CrossRef]
- European Commission. Health Promotion and Disease Prevention Knowledge Gateway. Available online: https://knowledge4policy.ec.europa.eu/health-promotion-knowledge-gateway_en (accessed on 1 October 2024).
- Jamioł-Milc, D.; Biernawska, J.; Liput, M.; Stachowska, L.; Domiszewski, Z. Seafood Intake as a Method of Non-Communicable Diseases (NCD) Prevention in Adults. *Nutrients* **2021**, *13*, 1422. [CrossRef] [PubMed]
- Maphosa, Y.; Jideani, V.A. The role of legumes in human nutrition. In *Functional Food—Improve Health Through Adequate Food*; Hueda, M.C., Ed.; IntechOpen: London, UK, 2017. [CrossRef]
- Mullins, A.P.; Arjmandi, B.H. Health Benefits of Plant-Based Nutrition: Focus on Beans in Cardiometabolic Diseases. *Nutrients* **2021**, *13*, 519. [CrossRef] [PubMed]
- CBI Ministry of Foreign Affairs. The European Market Potential for Canned Fish. Available online: <https://www.cbi.eu/market-information/fish-seafood/canned-fish/market-potential> (accessed on 11 November 2024).
- Akhbarizadeh, R.; Dobaradaran, S.; Nabipour, I.; Tajbakhsh, S.; Darabi, A.H.; Spitz, J. Abundance, composition, and potential intake of microplastics in canned fish. *Mar. Pollut. Bull.* **2020**, *160*, 111633. [CrossRef]
- Europe Canned Food Market. Europe Canned Food Market Research Report—Segmented By Type, Distribution Channel and Country (UK, France, Spain, Germany, Italy, Russia, Sweden, Denmark, Switzerland, Netherlands, Turkey, Czech Republic and Rest of Europe)—Industry Analysis on Size, Share, Trends, & Growth Forecast (2024 to 2032). Available online: <https://www.marketdataforecast.com/market-reports/europe-canned-food-market> (accessed on 11 November 2024).
- Fang, X.; Zhang, Y. The impact of climate change and economic development on the catches of small pelagic fisheries. *Mar. Policy* **2025**, *175*, 106631. [CrossRef]
- European Commission. Commission Proposes Fishing Opportunities for 2024 in the Atlantic, Kattegat and Skagerrak. Available online: https://ec.europa.eu/commission/presscorner/detail/en/ip_23_5279 (accessed on 11 October 2024).
- Fish Base. *Trachurus trachurus* (Linnaeus, 1758)—Atlantic Horse Mackerel. Available online: <https://www.fishbase.se/summary/SpeciesSummary.php?ID=1365&AT=horse+mackerel> (accessed on 12 October 2024).
- IPMA. Carapau-Branco—Trachurus Trachurus. Available online: <https://www.ipma.pt/pt/pescas/recursos/carapau/> (accessed on 15 October 2024).
- Bandarra, N.M.; Calhau, M.A.; Oliveira, L.; Ramos, M.; Dias, M.G.; Bártolo, H.; Faria, M.R.; Fonseca, M.C.; Gonçalves, J.; Batista, I.; et al. *Composition and Nutritional Value of Most Consumed Seafood Products in Portugal*; Publicações Avulsas Do IPIMAR 11; IPIMAR: Lisboa, Portugal, 2004. Available online: <https://www.ipma.pt/resources.www/docs/publicacoes.site/pescado/inicio.htm> (accessed on 20 December 2024).
- Costa, A.; Silva, A.; Rocha, A.; Pereira, B.; Maia, C.; Chaves, C.; Silva, C.; Rosa, D.; Feijó, D.; Mendes, H.; et al. *Estado dos Stocks em 2020 e Aconselhamento Científico para a sua Gestão em 2021*; Instituto Português do Mar e da Atmosfera (IPMA): Lisboa, Portugal, 2020; 133p. Available online: https://www.ipma.pt/pt/media/noticias/documentos/2021/ESTADO_DOS_RECursos_EXPLORADOS_2020-2.pdf (accessed on 20 November 2024).
- Instituto Nacional de Saúde Dr. Ricardo Jorge, I.P. (INSA). Food Composition Table. v 6.0—2023. Available online: <https://portfir-insa.min-saude.pt/> (accessed on 11 January 2025).
- EFSA NDA Panel (EFSA Panel on Nutrition, Novel Foods and Food Allergens); Turck, D.; Castenmiller, J.; de Henauw, S.; Hirsch-Ernst, K.-I.; Kearney, J.; Knutsen, H.K.; Maciuk, A.; Mangelsdorf, I.; McArdle, H.J.; et al. Scientific opinion on the dietary reference values for sodium. *EFSA J.* **2019**, *17*, 5778. [CrossRef]
- Oliveira, H.; Blocquel, C.; Santos, M.; Fretigny, M.; Correia, T.; Gonçalves, A.; Cabado, A.G.; López, L.B.; Raaholt, B.W.; Ferraris, F.; et al. Semi-industrial development of nutritious and healthy seafood dishes from sustainable species. *Food Chem. Toxicol.* **2021**, *155*, 112431. [CrossRef]
- World Health Organization (WHO). *WHO Global Sodium Benchmarks for Different Food Categories*; Licence: CC BY-NC-SA 3.0 IGO; WHO: Geneva, Switzerland, 2021; 24p. Available online: <https://www.who.int/publications/i/item/9789240025097> (accessed on 15 January 2025).
- Xiao, H.-W.; Pan, Z.; Deng, L.-Z.; El-Mashad, H.M.; Yang, X.-H.; Mujumdar, A.S.; Gao, Z.-J.; Zhang, Q. Recent developments and trends in thermal blanching—A comprehensive review. *Inf. Process. Agric.* **2017**, *4*, 101–127. [CrossRef]
- Warne, D. *Manual on Fish Canning*; FAO Fisheries Technical Paper: Rome, Italy, 1988; Volume 285, 72p.
- Labuza, T.P.; Fu, B. Growth kinetics for shelf-life prediction: Theory and practice. *J. Ind. Microbiol.* **1993**, *12*, 309–323. [CrossRef]

23. Instituto Português da Qualidade (IPQ). Norma Portuguesa NP 4404-1:2002—Microbiologia alimentar: Conservas. In *Parte 1: Apreciação da Estabilidade. Provas de Estufa (Método Corrente)*; IPQ: Caparica, Portugal, 2002; 9p.
24. Instituto Português da Qualidade (IPQ). Norma Portuguesa NP 2309-2:1998—Microbiologia alimentar: Conservas. In *Parte 2: Apreciação da Esterilidade*; IPQ: Caparica, Portugal, 1998; 16p.
25. Association of Official Analytical Chemists (AOAC) International. *Official Methods of Analysis*, 18th ed.; AOAC International: Gaithersburg, MD, USA, 2005.
26. Saint-Denis, T.; Goupy, J. Optimization of a nitrogen analyser based on the Dumas method. *Anal. Chim. Acta.* **2004**, *515*, 191–198. [[CrossRef](#)]
27. Food and Agriculture Organization of the United Nations (FAO). *Food Energy—Methods of Analysis and Conversion Factors*; FAO: Rome, Italy, 2003; 87p. Available online: https://www.fao.org/uploads/media/FAO_2003_Food_Energy_02.pdf (accessed on 15 October 2024).
28. Association of Official Analytical Chemists (AOAC) International. Salt (Chlorine as Sodium Chloride) in Seafood. In *Official Methods of Analysis*, 937.09, 11; AOAC: Washington, DC, USA, 1990; 870p.
29. Jorhem, L. Determination of metals in foods by atomic absorption spectrometry after dry ashing: NMKL collaborative study. *J. AOAC Int.* **2000**, *83*, 1204–1211. [[CrossRef](#)] [[PubMed](#)]
30. Veciana-Nogues, M.T.; Hernandez-Jover, T.; Marine-Font, A.; Vidal-Carou, M.C. Liquid chromatographic method for determination of biogenic amines in fish and fish products. *J. AOAC Int.* **1995**, *78*, 1045–1050. [[CrossRef](#)] [[PubMed](#)]
31. Cobb, B.F., III; Alaniz, I.; Thompson, C.A., Jr. Biochemical and microbial studies on shrimp: Volatile nitrogen and amino nitrogen analysis. *J. Food Sci.* **1973**, *38*, 431–436. [[CrossRef](#)]
32. Pathare, P.B.; Opara, U.L.; Al-Said, F.A.-J. Colour Measurement and Analysis in Fresh and Processed Foods: A Review. *Food Bioprocess Technol.* **2013**, *6*, 36–60. [[CrossRef](#)]
33. ISO 8589; Sensory Analysis—General Guidance for the Design of Test Rooms. ISO: Geneva, Switzerland, 2007.
34. Meilgaard, M.C.; Civille, G.V.; Carr, B.T. *Sensory Evaluation Techniques*, 5th ed.; CRC Press, Taylor & Francis Group: Boca Raton, FL, USA, 2016.
35. Viegas, C. @Nutrisoft—@Nutrition Information Software, Registo n. 3208/2012; Inspeção Geral das Atividades Culturais (IGAC): Lisboa, Portugal, 2012.
36. European Food Safety Authority (EFSA). Dietary Reference Values for nutrients. Summary report. *EFSA Support. Publ.* **2017**, *14*, e15121E. [[CrossRef](#)]
37. Zar, J.H. *Biostatistical Analysis*, 5th ed.; Prentice Hall Pearson: Upper Saddle River, NJ, USA, 2010.
38. Andrés-Bello, A.; Barreto-Palacios, V.; García-Segovia, P.; Mir-Bel, J.; Martínez-Monzó, J. Effect of pH on Color and Texture of Food Products. *Food Eng. Rev.* **2013**, *5*, 158–170. [[CrossRef](#)]
39. Aubourg, S.P. Review: Loss of Quality during the Manufacture of Canned Fish Products. *Food Sci. Technol. Int.* **2001**, *7*, 199–215. [[CrossRef](#)]
40. IPMA. Carapau (*Trachurus trachurus*). Available online: https://www.ipma.pt/resources.www/docs/publicacoes.site/pescado/site/carapau/carap_all.htm (accessed on 25 October 2024).
41. Celik, M. Seasonal changes in the proximate chemical compositions and fatty acids of chub mackerel (*Scomber japonicus*) and horse mackerel (*Trachurus trachurus*) from the north eastern Mediterranean Sea. *Int. J. Food Sci. Technol.* **2008**, *43*, 933–938. [[CrossRef](#)]
42. Durmuş, M. Evaluation of Nutritional and Mineral-Heavy Metal Contents of Horse Mackerel (*Trachurus trachurus*) in the Middle Black Sea in terms of Human Health. *Biol. Trace Elem. Res.* **2019**, *190*, 208–216. [[CrossRef](#)]
43. Arturpesca. The Atlantic Horse Mackerel. News. Available online: <https://asturpesca.com/en/2021/07/13/the-atlantic-horse-mackerel/> (accessed on 30 October 2024).
44. Rybicka, I.; Gonçalves, A.; Oliveira, H.; Marques, A.; Nunes, M.L. Salt reduction in seafood—A review. *Food Control* **2022**, *135*, 108809. [[CrossRef](#)]
45. European Union. Regulation (EC) No 1924/2006 of the European Parliament and of the Council of 20 December 2006 on nutrition and health claims made on foods. *Off. J. Eur. Union.* **2006**, *L404*, 9–25. Available online: <http://data.europa.eu/eli/reg/2006/1924/oj> (accessed on 30 October 2024).
46. Sousa, S.M.; Gregório, M.J.; Bernardino, F.; Fernandes, I.; Anjo, C.; Martins, S.; Bica, M.; Bandarra, N.; Carvalho, T.; Graça, P. *Receitas Com Enlatados (Can Recipes). Alimentação Saudável à Base de Conservas de Pescado “Made in Portugal”*; Estratégia Integrada para a Promoção da Alimentação Saudável (EIPAS): Lisboa, Portugal, 2015; 106p, ISBN 978-972-675-281-3.
47. Nielsen, T.; Mihnea, M.; Bâth, K.; Cunha, S.C.; Ferreira, R.; Fernandes, J.O.; Gonçalves, A.; Nunes, M.L.; Oliveira, H. New Formulation for Producing Salmon Pâté with Reduced Sodium Content. *Food Chem. Toxicol.* **2020**, *143*, 111546. [[CrossRef](#)] [[PubMed](#)]

48. World Health Organization. Sodium Reduction. Available online: <https://www.who.int/news-room/fact-sheets/detail/salt-reduction> (accessed on 31 October 2024).
49. Estévez, A.; Camacho, C.; Correia, T.; Barbosa, V.; Marques, A.; Lourenço, H.; Serrano, C.; Sapata, M.; Duarte, M.P.; Pires, C.; et al. Strategies to reduce sodium levels in European seabass sausages. *Food Chem. Toxicol.* **2021**, *153*, 112262. [CrossRef] [PubMed]
50. European Union. Regulation (EU) No 1169/2011 of the European Parliament and of the Council of 25 October 2011. *Off. J. Eur. Union.* **2011**, *L304*, 18–63. Available online: <http://data.europa.eu/eli/reg/2011/1169/oj> (accessed on 27 October 2024).
51. National Institutes of Health (NIH). Phosphorus. Fact Sheet for Consumers. Available online: <https://ods.od.nih.gov/factsheets/Phosphorus-Consumer/> (accessed on 30 November 2024).
52. EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA). Scientific Opinion on the substantiation of health claims related to vitamin B12 and contribution to normal neurological and psychological functions (ID 95, 97, 98, 100, 102, 109), contribution to normal homocysteine metabolism (ID 96, 103, 106), maintenance of normal bone (ID 104), maintenance of normal teeth (ID 104), maintenance of normal hair (ID 104), maintenance of normal skin (ID 104), maintenance of normal nails (ID 104), reduction of tiredness and fatigue (ID 108), and cell division (ID 212) pursuant to Article 13(1) of Regulation (EC) No 1924/2006. *EFSA J.* **2010**, *8*, 1756. [CrossRef]
53. National Institutes of Health (NIH). Vitamin B12. Fact Sheet for Consumers. Available online: <https://ods.od.nih.gov/factsheets/VitaminB12-Consumer/> (accessed on 28 October 2024).
54. EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA). Scientific Opinion on the substantiation of health claims related to vitamin D and maintenance of bone and teeth (ID 150, 151, 158), absorption and utilisation of calcium and phosphorus and maintenance of normal blood calcium concentrations (ID 152, 157), cell division (ID 153), and thyroid function (ID 156) pursuant to Article 13(1) of Regulation (EC) No 1924/2006 on request from the European Commission. *EFSA J.* **2009**, *7*, 1227. [CrossRef]
55. EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA). Scientific Opinion on the substantiation of health claims related to vitamin D and normal function of the immune system and inflammatory response (ID 154, 159), maintenance of normal muscle function (ID 155) and maintenance of normal cardiovascular function (ID 159) pursuant to Article 13(1) of Regulation (EC) No 1924/2006. *EFSA J.* **2010**, *8*, 1468. [CrossRef]
56. EFSA Panel on Dietetic Products, Nutrition and Allergies (NDA). Scientific opinion on the tolerable upper intake level of vitamin D. *EFSA J.* **2012**, *10*, 2813. [CrossRef]
57. Çın, N.N.A.; Yüce, M.; Özçelik, A.Ö. The iodine content of raw, boiled, pan-fried, and oven-baked fish samples using ICP-MS: Analysis and reliability of measurements. *Int. J. Food Sci. Technol.* **2024**, *59*, 3226–3234. [CrossRef]
58. Bógnar, A. *Tables on Weight Yield of Food and Retention Factors of Food Constituents for the Calculation of Nutrient Composition of Cooked Foods (Dishes)*; BFE-R--02-03; Berichte der Bundesforschungsanstalt für Ernährung: Karlsruhe, Germany, 2002; 98p. Available online: http://www.fao.org/uploads/media/bognar_bfe-r-02-03.pdf (accessed on 20 April 2019).
59. Mierke-Klemeyer, S.; Larsen, R.; Oehlenschläger, J.; Maehre, H.; Elvevoll, E.O.; Bandarra, N.M.; Parreira, R.; Andrade, A.M.; Nunes, M.L.; Schram, E.; et al. Retention of health-related beneficial components during household preparation of selenium-enriched African catfish (*Clarias gariepinus*) fillets. *Eur. Food Res. Technol.* **2008**, *227*, 827–833. [CrossRef]
60. University of Rochester Medical Center. Health Encyclopedia. Vitamin B12. Available online: <https://www.urmc.rochester.edu/encyclopedia/content.aspx?contenttypeid=19&contentid=vitaminb-12> (accessed on 20 September 2024).
61. University of Rochester Medical Center. Health Encyclopedia. Vitamin D. Available online: <https://www.urmc.rochester.edu/encyclopedia/content.aspx?contenttypeid=19&contentid=vitamind> (accessed on 20 September 2024).
62. devour. What Time Do People Eat in Portugal? Available online: <https://devourtours.com/blog/what-time-do-people-eat-in-portugal/?cnt=PT> (accessed on 20 September 2024).
63. Colombo, F.M.; Cattaneo, P.; Confalonieri, E.; Bernardi, C. Histamine food poisonings: A systematic review and meta-analysis. *Crit. Rev. Food Sci. Nutr.* **2018**, *58*, 1131–1151. [CrossRef] [PubMed]
64. Comas-Basté, O.; Sánchez-Pérez, S.; Veciana-Nogués, M.T.; Latorre-Moratalla, M.; Vidal-Carou, M.D.C. Histamine intolerance: The current state of the art. *Biomolecules* **2020**, *10*, 1181. [CrossRef] [PubMed]
65. European Union. Commission Regulation (EC) No 2073/2005 of 15 November 2005 on microbiological criteria for foodstuffs. *Off. J. Eur. Union.* **2005**, *L 338*, 1–26.
66. EFSA (European Food Safety Authority). Assessment of the incidents of histamine intoxication in some EU countries. *EFSA Support. Publ. EN-1301* **2017**, *14*, 1–37.
67. Muscarella, M.; Lo Magro, S.; Campaniello, M.; Armentano, A.; Stacchini, P. Survey of histamine levels in fresh fish and fish products collected in Puglia (Italy) by ELISA and HPLC with fluorimetric detection. *Food Control* **2013**, *31*, 211–217. [CrossRef]
68. Visciano, P.; Schirone, M.; Paparella, A. An Overview of Histamine and Other Biogenic Amines in Fish and Fish Products. *Foods* **2020**, *9*, 1795. [CrossRef]
69. European Union. Commission Regulation (EC) No 1022/2008 of 17 October 2008 amending Regulation (EC) No 2074/2005 as regards the total volatile basic nitrogen (TVB-N) limits (Text with EEA relevance). *Off. J. Eur. Union.* **2008**, *L 277*, 18–20.

70. King, V.A.; Liu, C.; Liu, Y. Chlorophyll stability in spinach dehydrated by freeze-drying and controlled low-temperature vacuum dehydration. *Food Res. Int.* **2001**, *34*, 167–175. [CrossRef]
71. MacEvoy, B. Modern Color Models. Available online: <https://www.handprint.com/HP/WCL/color7.html#hsv> (accessed on 6 December 2024).

Disclaimer/Publisher’s Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.